

Références bibliographiques citées dans le chapitre 1

- ACTA (2002). Recueil des effets non intentionnels des produits phytosanitaires, 8^{ème} ed., ACTA - UIPP, Paris, 492 p.
- ACTA (2008). Index Phytosanitaire 44, ACTA, Paris, 844 p.
- Adler P.B., Milchunas D.G., Iauenroth W.K., Sala O.E., Burke I.C. (2004). Functional traits of graminoids in semi-arid steppes: a test of grazing histories. *Journal of Applied Ecology* 41(4): 653-663.
- Adler P.B., Milchunas D.G., Sala O.E., Burke I.C., Iauenroth W.K. (2005). Plant traits and ecosystem grazing effects: comparison of US sagebrush steppe and Patagonian steppe. *Ecological Applications* 15(2): 774-792.
- Adler P.B., Raff D.A., Iauenroth W.K. (2001). The effect of grazing on the spatial heterogeneity of vegetation. *Oecologia* 128(4): 465-479.
- Adriaens D., Honnay O., Hermy M. (2006). No evidence of a plant extinction debt in highly fragmented calcareous grasslands in Belgium. *Biological Conservation* 133(2): 212-224.
- Aebischer N.J. (1990). Assessing pesticide effects on non-target invertebrates using long term monitoring and time-series modelling. *Functional Ecology* 4: 369-373.
- Aerts R., Chapin F.S.I. (2000). The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. *Advances in Ecological Research* 30: 1-67.
- Agreste (1998). Enquêtes sur la structure des vergers en 1997.
- Agreste (2006). Statistique Agricole Annuelle. <http://agreste.agriculture.gouv.fr/IMG/pdf/saa2008T3a.pdf>
- Agricola U., Barthel J., Laussmann H., Plachter H. (1996). Structure and dynamics of the fauna of an agricultural landscape in Southern Germany after change of landuse to ecological and integrated farming systems. *Verhandlungen der Gesellschaft für Ökologie* 26: 681-692.
- Agrios G.N. (1997). Plant Pathology, Fourth Edition ed., Academic Press, San Diego, 635 p.
- Ahmed M.T., Ismail S.M.M., Mabrouk S.S. (1998). Residues of some chlorinated hydrocarbon pesticides in rain water, soil and ground water, and their influence on some soil microorganisms. *Environment International* 24(5-6): 665-670.
- Aihartzia J.R., Garin I., Goiti U., Zabala J., Zuberogoitia I. (2003). Spring habitat selection by the Mediterranean Horseshoe Bat (*Rhinolophus euryale*) in the Urdaibai Biosphere Reserve (Basque Country). *Mammalia* 67(1): 25-32.
- Alabouvette C., Backhouse D., Steinberg C., Donovan N.J., Edel-Hermann V., Burgess L.W. (2004). Microbial diversity in soil - Effects on crop health. In *Managing Soil Quality: Challenges in Modern Agriculture*, (Emholt S., Christensen B.T., eds.), CABI Publishing, New York: 121-138.
- Alabouvette C., Olivain C., Steinberg C. (2006). Biological control of plant diseases: the European situation. *European Journal of Plant Pathology* 114(3): 329-341.
- Alabouvette C., Raaijmakers J., De Boer W., Notz R., Défago G., Steinberg C., Lemanceau P. (2006). Concepts and Methods to Assess the Phytosanitary Quality of Soils. In *Plant-microbe interactions and soil quality handbook*, (Bloem J. et al., eds.), CABI Publishing, Wallingford, UK: 257-270.
- Alabouvette C., Steinberg C. (2006). The soil as a reservoir for antagonists to plant diseases. In *An ecological and societal approach to biological control*, (Eilenberg J., Hokkanen H., eds.), Kluwer Academic Publishers, NL, Dordrecht: 123-144.
- Alard D., Bance J.F., Frileux P.N. (1994). Grassland vegetation as an indicator of the main agro-ecological factors in a rural landscape: consequences for biodiversity and wildlife conservation in central Normandy (France). *Journal of Environmental Management* 42(2): 91-109.
- Albrecht A., Mattheis A. (1998). The effects of organic and integrated farming on rare arable weeds on the Forschungsverbund Agrarökosysteme München (FAM) research station in southern Bavaria. *Biological Conservation* 86(3): 347-356.
- Albrecht H. (2003). Suitability of arable weeds as indicator organisms to evaluate species conservation effects of management in agricultural ecosystems. *Agriculture, Ecosystems & Environment* 98(1/3): 201-211.
- Al-Deeb M.A., Wilde G.E., Blair J.M., Todd T.C. (2003). Effect of Bt corn for corn rootworm control on nontarget soil microarthropods and nematodes. *Environmental Entomology* 32: 859-865.
- Alix A., Cortesero A.M., Nenon J.P., Anger J.P. (2001). Selectivity assessment of chlorfenvinphos reevaluated by including physiological and behavioral effects on an important beneficial insect. *Environmental Toxicology and Chemistry* 20(11): 2530-2536.
- Al-Mufti M.M., Sydes C.L., Furness S.B., Grime J.P., Band S.R. (1977). A quantitative analysis of shoot phenology and dominance in herbaceous vegetation. *Journal of Ecology* 65: 759-791.
- Altieri M.A. (1999). The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystems and Environment* 74(1-3): 19-31.
- Altieri M.A., Schmidt L.L. (1986). The dynamics of colonizing arthropod communities at the interface of abandoned, organic and commercial apple orchards and adjacent woodland habitats. *Agriculture, Ecosystems & Environment* 16(1): 29-43.
- Amiaud B., Bouzille J.B., Tournade F., Bonis A. (1998). Spatial patterns of soil salinities in old embanked marshlands in western France. *Wetlands* 18(3): 482-494.
- Anand M., Laurence S., Rayfield B. (2005). Diversity relationships among taxonomic groups in recovering and restored forests. *Conservation Biology* 19(3): 955-962.
- Andelman S.J., Fagan W.F. (2000). Umbrellas and flagships: efficient conservation surrogates or expensive mistakes? *Proceedings of the National Academy of Sciences of the United States of America* 97(11): 5954-5959.
- Andersen A.N., Hoffmann B.D., Muller W.J., Griffiths A.D. (2002). Using ants as bioindicators in land management: simplifying assessment of ant community responses. *Journal of Applied Ecology* 39(1): 8-17.
- Andersen A.N., Majer J.D. (2004). Ants show the way Down Under: invertebrates as bioindicators in land management. *Frontiers in Ecology and the Environment* 2(6): 291-298.
- Anderson T.H. (2003). Microbial eco-physiological indicators to assess soil quality. *Agriculture Ecosystems & Environment* 98(1-3): 285-293.
- Andow D.A. (1991). Vegetational diversity and arthropod population response. *Annual Review of Entomology* 36: 561-586.
- Andow D.A., Risch S.J. (1985). Predation in diversified agroecosystems: relations between a coccinellid predator *Coleomegilla maculata* and its food. *Journal of Applied Ecology* 22(2): 357-372.

- Andow D.A., Zwahlen C. (2006). Assessing environmental risks of transgenic plants. *Ecology Letters* 9(2): 196-214.
- Andrade D.D., Colozzi A., Balota E.L., Hungria M. (2003). Long-term effects of agricultural practices on microbial community. In *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-Economy, Policy*, KLUWER ACADEMIC PUBL, Dordrecht: 301-306.
- Andr en H. (1994). Effects of habitat fragmentation on birds and mammals in landscape with different proportions of suitable habitat: a review. *Oikos* 71: 355-366.
- Andren H., Delin A., Cuthill C. (1997). Population response to landscape changes depends on specialization to different landscape elements. *Forum. Oikos* 80: 193-196.
- Angeler D.G., Garcia G. (2005). Using emergence from soil propagule banks as indicators of ecological integrity in wetlands: advantages and limitations. *Journal of the North American Benthological Society* 24(4): 740-752.
- Angevin F., Klein E.K., Choimet C., Gauffreteau A., Lavigne C., Mess an A., Meynard J.M. (2008). Modelling impacts of cropping systems and climate on maize cross-pollination in agricultural landscapes: The MAPOD model. *European Journal of Agronomy* 28(3): 471-484.
- Arias M.E., Gonzalez-Perez J.A., Gonzalez-Vila F.J., Ball A.S. (2005). Soil health - a new challenge for microbiologists and chemists. *International Microbiology* 8(1): 13-21.
- Arlettaz R. (1996). Feeding behaviour and foraging strategy of free-living mouse-eared bats, *Myotis myotis* and *Myotis blythii*. *Animal Behaviour* 51(1): 1-11.
- Arlettaz R., Perrin N., Hausser J. (1997). Trophic Resource Partitioning and Competition between the Two Sibling Bat Species *Myotis myotis* and *Myotis blythii*. *Journal of Animal Ecology* 66(6): 897-911.
- Arnault I., Mondy N., Diwo S., Auger J. (2004). Soil behaviour of sulfur natural fumigants used as methyl bromide substitutes. *International Journal of Environmental Analytical Chemistry* 84(1-3): 75-82.
- Artois M. (2003). Wildlife infectious disease control in Europe. *Journal Mountain Ecology* 7(7): 89-97.
- Artois M., Biteau-Coroller F., Rossi S., Hars J. (2002). La surveillance et le contr le des maladies infectieuses de la faune sauvage en France et en Europe. *Bull. Soc. M d. v t. prat.* 86: 365-1.
- Auge R.M. (2004). Arbuscular mycorrhizae and soil/plant water relations. *Canadian Journal of Soil Science* 84(4): 373-381.
- Auger P., Bonafos R., Kreiter S. (2004). Mancozeb resistance patterns among *Kampimodromus aberrans* and *Typhlodromus pyri* (Acari : Phytoseiidae) strains from French vineyards. *Canadian Entomologist* 136(5): 663-673.
- Aveline A., Crozat Y., Cleyet-Marel J.C., Pinochet X. (2003). Bacterial growth rate and growth pouch nodulation profile differences as possible ways of *Bradyrhizobium japonicum* strain screening for low P soils. *Plant and Soil* 251(2): 199-209.
- Avenant N.L. (2000). Small mammal community characteristics as indicators of ecological disturbance in the Willem Pretorius Nature Reserve, Free State, South Africa. *South African Journal of Wildlife Research* 30(1): 26-33.
- Aviron S., Burel F., Baudry J., Schermann N. (2005). Carabid assemblages in agricultural landscapes: impacts of habitat features, landscape context at different spatial scales and farming intensity. *Agriculture, Ecosystems & Environment* 108(3): 205-217.
- Avrhami S., Conrad R. (2003). Patterns of community change among ammonia oxidizers in meadow soils upon long-term incubation at different temperatures. *Applied and Environmental Microbiology* 69(10): 6152-6164.
- Axelsen J.A., Kristensen K.T. (2000). Collembola and mites in plots fertilised with different types of green manure. *Pedobiologia* 44(5): 556-566.
- Aymonin G. (1965). Origines pr sum es et disparition progressive des adventices messicoles calcicoles en France, 2  Colloque sur la *Biologie des mauvaises herbes*, ENSA, Grignon, Guyot, ed.: 1-15.
- Aznar J.C., Dervieux A., Grillas P. (2003). Association between aquatic vegetation and landscape indicators of human pressure. *Wetlands* 23(1): 149-160.
- Baessler C., Klotz S. (2006). Effects of changes in agricultural land-use on landscape structure and arable weed vegetation over the last 50 years. *Agriculture, Ecosystems & Environment* 115(1-4): 43-50.
- Baguette M., Hance T. (1997). Carabid beetles and agricultural practices: Influence of soil ploughing. *Biological Agriculture & Horticulture* 15(1-4): 185-190.
- Bailey K.L., Gossen B.D., Derksen D.A., Watson P.R. (2000). Impact of agronomic practices and environment on diseases of wheat and lentil in southeastern Saskatchewan. *Canadian Journal of Plant Science* 80(4): 917-927.
- Bailey K.L., Lazarovits G. (2003). Suppressing soil-borne diseases with residue management and organic amendments. *Soil & Tillage Research* 72(2): 169-180.
- Bailey V.L., Smith J.L., Bolton H. (2002). Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration. *Soil Biology & Biochemistry* 34(7): 997-1007.
- Baker G.H. (1998). Recognising and responding to the influences of agriculture and other land-use practices on soil fauna in Australia. *Applied Soil Ecology* 9(1-3): 303-310.
- Bakker C., Blair J.M., Knapp A.K. (2003). Does resource availability, resource heterogeneity or species turnover mediate changes in plant species richness in grazed grasslands? *Oecologia* 137(3): 385-391.
- Bakker E.S., Ritchie M.E., Olf H., Milchunas D.G., Johannes M.H. (2006). Herbivore impact on grassland plant diversity depends on habitat productivity and herbivore size. *Ecology Letters* 9(7): 780-788.
- Bakker J.P. (1998). The impact of grazing on plant communities. In *Grazing and conservation management*, (Wallis De Vries M.F. et al., eds.), Kluwer Academic Publisher, Dordrecht: 137-184.
- Bakker J.P., Berendse F. (1999). Constraints in the restoration of ecological diversity in grassland and heathland communities. *Trends in Ecology & Evolution* 14(2): 63-68.
- Bal zs K., Jenser G., Buj ki G. (1996). Eight years' experiences of IPM in Hungarian apple orchards. *Bulletin OILB/SROP* 19(4): 95-101.
- Baldi A., Batary P., Erdos S. (2005). Effects of grazing intensity on bird assemblages and populations of Hungarian grasslands. *Agriculture, Ecosystems & Environment* 108(3): 251-263.
- Baldi A., Kisbenedek T. (1997). Orthopteran assemblages as indicators of grassland naturalness in Hungary. *Agriculture Ecosystems & Environment* 66(2): 121-129.
- Balent G., Allard D., Blanfort V., Gibon A. (1998). Activit s de p turage, paysages et biodiversit . *Annales de Zootechnie* 47: 419-429.

- Balmford A., Crane P., Dobson A., Green R.E., Mace G.M. (2005). The 2010 challenge: Data availability, information needs and extraterrestrial insights. *Philosophical Transactions of the Royal Society B-Biological Sciences* 360(1454): 221-228.
- Balmford A., Green M.J.B., Murray M.G. (1996). Using higher-taxon richness as a surrogate for species richness .1. Regional tests. *Proceedings of the Royal Society of London Series B-Biological Sciences* 263(1375): 1267-1274.
- Bamforth S.S. (1997). Protozoa: recyclers and indicators of agroecosystem quality. In *Fauna in soil ecosystems: recycling, processes, nutrient fluxes and agricultural production*, (Benckiser G., ed.), Marcel Dekker: 63-84.
- Bankhead S.B., Landa B.B., Lutton E., Weller D.M., Gardener B.B.M. (2004). Minimal changes in rhizobacterial population structure following root colonization by wild type and transgenic biocontrol strains. *Fems Microbiology Ecology* 49(2): 307-318.
- Bar A., Löffler J. (2007). Ecological process indicators used for nature protection scenarios in agricultural landscapes of SW Norway. *Ecological Indicators* 7(2): 396-411.
- Barbour J.D., Farrar R.R., Kennedy G.G. (1993). Interaction of *Manduca sexta* resistance in tomato with insect predators of *Helicoverpa zea*. *Entomologia Experimentalis et Applicata* 68: 143-155.
- Bardgett R.D., Cook R. (1998). Functional aspects of soil animal diversity in agricultural grasslands. *Applied Soil Ecology* 10(3): 263-276.
- Bardgett R.D., Jones A.C., Jones D.L., Kemmitt S.J., Cook R., Hobbs P.J. (2001). Soil microbial community patterns related to the history and intensity of grazing in sub-montane ecosystems. *Soil Biology & Biochemistry* 33(12-13): 1653-1664.
- Bardgett R.D., McAlister E. (1999). The measurement of soil fungal : bacterial biomass ratios as an indicator of ecosystem self-regulation in temperate meadow grasslands. *Biology and Fertility of Soils* 29(3): 282-290.
- Barnard D.R., Rogers G.D., Jones B.G. (1983). Strategic treatment of pastured beef cattle with acaricides: effect on populations of the lone star tick (Acari: Ixodidae). *Journal of Economic Entomology* 76(1): 99-102.
- Barr C.J., Gillespie M.K. (2000). Estimating hedgerow length and pattern characteristics in Great Britain using Countryside Survey data. *Journal of Environmental Management* 60(1): 23-32.
- Barralis G. (1978). Modifications de la flore adventice des agrocénoses résultant de l'emploi des herbicides et leurs conséquences, *Colloque sur les incidences secondaires des herbicides sur la flore et la faune*, Versailles (France), Société Française de Phytologie et de Phytopharmacie: 24-29.
- Barrett C.B., Gibson C.C., Hoffman B., McCubbins M.D. (2006). The complex links between governance and biodiversity. *Conservation Biology* 20(5): 1358-1366.
- Barrios E. (2007). Soil biota, ecosystem services and land productivity. *Ecological Economics* 64(2): 269-285.
- Barton D.R. (1996). The use of Percent Model Affinity to assess the effects of agriculture on benthic invertebrate communities in headwater streams of southern Ontario, Canada. *Freshwater Biology* 36(2): 397-410.
- Barton D.R., Farmer M.E.D. (1997). The effects of conservation tillage practices on benthic invertebrate communities in headwater streams in southwestern Ontario, Canada. *Environmental Pollution* 96(2): 207-215.
- Bass B., Hansell R., Choi J. (1998). Towards a simple indicator of biodiversity. *Environmental Monitoring and Assessment* 49(2-3): 337-347.
- Bastian O., Lutz M., Unger C., Koppen I., Roder M., Syrbe R.U. (2003). Framework for the development of local agri-environmental programmes in Europe - 1. The indicator approach. *Landnutzung und Landwirtschaft* 44(5): 229-237.
- Batary P., Baldi A., Szel G., Podlussany A., Rozner I., Erdos S. (2007). Responses of grassland specialist and generalist beetles to management and landscape complexity. *Diversity and Distributions* 13(2): 196-202.
- Bauernfeind E., Moog O. (2000). Mayflies (Insecta: Ephemeroptera) and the assessment of ecological integrity: a methodological approach. *Hydrobiologia* 422/423: 71-83.
- Baumgarte S., Tebbe C.C. (2005). Field studies on the environmental fate of the Cry1Ab Bt-toxin produced by transgenic maize (MON810) and its effect on bacterial communities in the maize rhizosphere. *Molecular Ecology* 14(8): 2539-2551.
- Baumgartner J., Hartmann J. (2001). The design and implementation of sustainable plant diversity conservation program for alpine meadows and pastures. *Journal of Agricultural & Environmental Ethics* 14(1): 67-83.
- Bazzaz F.A. (1975). Plant species diversity in old-field successional ecosystems in Southern Illinois. *Ecology* 56: 485-488.
- Bazzaz F.A. (1996). *Plants in Changing Environments. Linking Physiological, Population, and Community Ecology*, Cambridge University Press, Cambridge, 320 p.
- Beckie H.J., Harker K.N., Hall L.M., Warwick S.I., Légère A., Sikkema P.H., Clayton G.W., Thomas A.G., Leeson J.Y., Seguin-Swartz G., Simard M.-J. (2006). A decade of herbicide-resistant crops in Canada. *Canadian Journal of Plant Science* 86(4): 1243-1264.
- Beebee T.J.C., Griffiths R.A. (2005). The amphibian decline crisis: A watershed for conservation biology? *Biological Conservation* 125(3): 271-285.
- Behan-Pelletier V.M. (1999). Oribatid mite biodiversity in agroecosystems: role for bioindication. *Agriculture, Ecosystems & Environment* 74(1-3): 411-423.
- Behan-Pelletier V.M. (2003). Acari and Collembola biodiversity in Canadian agricultural soils. *Canadian Journal of Soil Science* 83(3): 279-288.
- Beintema A.J., Muskens G.J.D.M. (1987). Nesting success of birds breeding in Dutch agricultural grasslands. *Journal of Applied Ecology* 24(3): 743-758.
- Belanger L., Picard M. (1999). Cattle grazing and avian communities of the St. Lawrence River islands. *Journal of Range Management* 52(4): 332-338.
- Bellows T.S. (2001). Restoring population balance through natural enemy introductions. *Biological Control* 21: 199-205.
- Bel'skaya E.A., Eshyunin S.L. (2003). Arachnids (Arachnidae) in a spring wheat agrocenosis in Southern Sverdlovsk oblast and the effect of treatment with Decis, a pyrethroid insecticide, on their populations. *Russian Journal of Ecology* 34(5): 359-362.
- Beltman B., van der Broek T., Martin W., Ten Cate M., Gusewell S. (2003). Impact of mowing regime on species richness and biomass of a limestone hay meadow in Ireland. *Bulletin of the Geobotanical Institute ETH* 69(1): 17-30.
- Belyea L.R., Lancaster J. (1999). Assembly rules within a contingent ecology. *Oikos* 86: 402-416.
- Bengtsson J., Ahnstrom J., Weibull A.C. (2005). The effects of organic agriculture on biodiversity and abundance: a meta-analysis. *Journal of Applied Ecology* 42(2): 261-269.

- Benizri E., Amiaud B. (2005). Relationship between plants and soil microbial communities in fertilized grasslands. *Soil Biology & Biochemistry* 37(11): 2055-2064.
- Bennett A.E., Bever J.D. (2007). Mycorrhizal species differentially alter plant growth and response to herbivory. *Ecology* 88(1): 210-218.
- Bennett A.F., Radford J.Q., Haslem A. (2006). Properties of land mosaics: Implications for nature conservation in agricultural environments. *Biological Conservation* 133(2): 250-264.
- Bensusan N.R. (2000). Do we know how effective is the conservation of biodiversity in protected areas. *Boletim do Herbario Ezechias Paulo Heringer* 5: 113-123.
- Benton T.G., Bryant D.M., Cole L., Crick H.Q.P. (2002). Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology* 39(4): 673-687.
- Benton T.G., Vickery J.A., Wilson J.D. (2003). Farmland biodiversity: is habitat heterogeneity the key? *Trends in Ecology & Evolution* 18(4): 182-188.
- Berendse F., Aerts R. (1987). Nitrogen-use-efficiency: a biologically meaningful definition? *Functional Ecology* 1: 293-296.
- Berendse F., Elberse W.T., Geerts R.H.M.E. (1992). Competition and nitrogen loss from plants in grassland ecosystems. *Ecology* 73(1): 46-53.
- Berg G., Esselink P., Groeneweg M., Kiehl K. (1997). Micropatterns in *Festuca rubra*-dominated salt-marsh vegetation induced by sheep grazing. *Plant Ecology* 132(1): 1-14.
- Bernard J.-L., Gratadou P., Pindon G., Rodriguez A., Tisseur M., Decourtye A. (2006). Jachères et MAE : pour une gestion favorable à l'entomofaune prédatrice. *Phytoma-La défense des végétaux* 590: 10-16.
- Berny P. (2007). Pesticides and the intoxication of wild animals. *Journal of Veterinary Pharmacology and Therapeutics* 30: 93-100.
- Berrada S., Nguyen T.X., Fournier D. (1996). Comparative toxicities of some insecticides to *Cacopsylla pyri* L. (Hom., Psyllidae) and one of its important biological control agents, *Anthocoris nemoralis* F. (Het., Anthocoridae). *Journal of Applied Entomology* 120(3): 181-185.
- Bhatti M.A., Duan J., Head G.P., Jiang C.J., McKee M.J., Nickson T.E., Pilcher C.L., Pilcher C.D. (2005). Field evaluation of the impact of corn rootworm (Coleoptera : Chrysomelidae)-protected Bt corn on ground-dwelling invertebrates. *Environmental Entomology* 34(5): 1325-1335.
- Bhatti M.A., Duan J., Head G.P., Jiang C.J., McKee M.J., Nickson T.E., Pilcher C.L., Pilcher C.D. (2005). Field evaluation of the impact of corn rootworm (Coleoptera: Chrysomelidae)-protected Bt corn on foliage-dwelling arthropods. *Environmental Entomology* 34(5): 1336-1345.
- Biala K., Peeters A., Muys B., Hermy M., Brouckaert V., Garcia V., Veken B.v.d., Valckx J. (2005). Biodiversity indicators as a tool to assess sustainability levels of agro-ecosystems, with a special consideration of grassland areas. *Options Méditerranéennes. Serie A, Seminaires Méditerranéennes*(No.67): 439-443.
- Bianchi F., Booij C.J.H., Tschamtker T. (2006). Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B-Biological Sciences* 273(1595): 1715-1727.
- Bibby C.J. (1999). Making the most of birds as environmental indicators. *Ostrich* 70(1): 81-88.
- Biddinger D.J., Hull L.A. (1995). Effects of several types of insecticides on the mite predator, *Stethorus punctum* (Coleoptera: Coccinellidae), including insect growth regulators and abamectin. *Journal of Economic Entomology* 88(2): 358-366.
- Bilalis D., Efthimiadis P., Sidiras N. (2001). Effect of three tillage systems on weed flora in a 3-year rotation with four crops. *Journal of Agronomy and Crop Science-Zeitschrift Fur Acker Und Pflanzenbau* 186(2): 135-141.
- Billeter R., Liira J., Bailey D., Bugter R., Arens P., Augenstein I., Aviron S., Baudry J., Bukacek R., Burel F., Cerny M., De Blust G., De Cock R., Diekötter T., Dietz H., Dirksen J., Dormann C., Durka W., Frenzel M., Hamersky R., Hendrickx F., Herzog F., Klotz S., Koolstra B., Lausch A., Le Coeur D., Maelfait J.P., Opdam P., Roubalova M., Schermann A., Schermann N., Schmidt T., Schweiger O., Smulders M.J.M., Speelmans M., Simova P., Verboom J., van Wingerden W.K.R.E., Zobel M. (2008). Indicators for biodiversity in agricultural landscapes: a pan-European study. *Journal of Applied Ecology* 45(1): 141-150.
- Binet F., Hallaire V., Curmi P. (1997). Agricultural practices and the spatial distribution of earthworms in maize fields. Relationships between earthworm abundance, maize plants and soil compaction. *Soil Biology & Biochemistry* 29(3-4): 577-583.
- Bittman S., Forge T.A., Kowalenko C.G. (2005). Responses of the bacterial and fungal biomass in a grassland soil to multi-year applications of dairy manure slurry and fertilizer. *Soil Biology & Biochemistry* 37(4): 613-623.
- Blackstock T., Rimes C., Stevens D., Jefferson R., Robertson H., J. M., Hopkins J. (1999). The extent of semi-natural grassland communities in lowland England and Wales: a review of conservation surveys 1978-1996. *Grass & Forage Science* 54(1): 1-18.
- Blair J.M., Todd T.C., Callahan J. (2000). Responses of grassland soil invertebrates to natural and anthropogenic disturbances. In *Invertebrates as webmasters in ecosystems*, (Coleman D.C., Hendrix P.F., eds.), CABI International Press: 43-71.
- Blair R.B. (1999). Birds and butterflies along an urban gradient: Surrogate taxa for assessing biodiversity? *Ecological Applications* 9(1): 164-170.
- Bleeker W., Schmitz U., Ristow M. (2007). Interspecific hybridization between alien and native plant species in Germany and its consequences for native biodiversity. *Biological Conservation* 137(2): 248-253.
- Blondel J., Farre H. (1988). The convergent trajectories of bird communities along ecological successions in European forests. *Oecologia* 75(1): 83-93.
- Blonk H., Lindeijer E., Broers J. (1997). Towards a methodology for taking physical degradation of ecosystems into account in LCA. *International Journal of Life Cycle Assessment* 2(2): 91-98.
- Bobbink R., Hornung M., Roelofs J.M. (1998). The effects of air-borne pollutants on species diversity in natural and semi-natural European vegetation. *Journal of Ecology* 86(5): 717-738.
- Bockstaller C., Girardin P. (2003). How to validate environmental indicators. *Agricultural Systems* 76(2): 639-653.
- Bockstaller C., Girardin P., vanderWerf H.M.G. (1997). Use of agro-ecological indicators for the evaluation of farming systems. *European Journal of Agronomy* 7(1-3): 261-270.
- Bockstaller C., Guichard L., Makowski D., Aveline A., Girardin P., Plantureux S. (2008). Agri-environmental indicators to assess cropping and farming systems. A review. *Agronomy for Sustainable Development* 28(1): 139-149.

- Bockus W.W., Shroyer J.P. (1998). The impact of reduced tillage on soilborne plant pathogens. *Annual Review of Phytopathology* 36: 485-500.
- Bogya S., Markó V. (1999). Effect of pest management systems on ground-dwelling spider assemblages in an apple orchard in Hungary. *Agriculture, Ecosystems & Environment* 73(1): 7-18.
- Bogya S., Markó V., Szinétár C.S. (1999). Comparison of pome fruit orchard inhabiting spider assemblages at different geographical scales. *Agricultural and Forest Entomology* 1(4): 261-269.
- Bohanec M., Cortet J., Griffiths B., Znidarsic M., Debeljak M., Caul S., Thompson J., Krogh P.H. (2007). A qualitative multi-attribute model for assessing the impact of cropping systems on soil quality. *Pedobiologia* 51(3): 239-250.
- Bohm M., White P.C.L., Chambers J., Smith L., Hutchings M.R. (2007). Wild deer as a source of infection for livestock and humans in the UK. *Veterinary Journal* 174: 260-276.
- Bolduc E., Buddle C.M., Bostanian N.J., Vincent C. (2005). Ground-dwelling spider fauna (Araneae) of two vineyards in southern Quebec. *Environmental Entomology* 34(3): 635-645.
- Bonet A., Pausas J. (2004). Species richness and cover along a 60-year chronosequence in old-fields of southeastern Spain. *Plant Ecology* 174(2): 257-270.
- Bonkowski M., Roy J. (2005). Soil microbial diversity and soil functioning affect competition among grasses in experimental microcosms. *Oecologia* 143(2): 232-240.
- Bonn A., Rodrigues A.S.L., Gaston K.J. (2002). Threatened and endemic species: are they good indicators of patterns of biodiversity on a national scale? *Ecology Letters* 5(6): 733-741.
- Borie V. (1892). Les travaux des champs, 4ème ed., Bibliothèque du Cultivateur, Librairie Agricole de la Maison Rustique, Paris, 180 p.
- Bos D., Loonen M.J.J.E., Stock M., Hofeditz F., van der Graaf A.J., Bakker J.P. (2005). Utilisation of Wadden Sea salt marshes by geese in relation to livestock grazing. *Journal for Nature Conservation* 13(1): 1-15.
- Bossuyt H., Deneff K., Six J., Frey S.D., Merckx R., Paustian K. (2001). Influence of microbial populations and residue quality on aggregate stability. *Applied Soil Ecology* 16(3): 195-208.
- Bostrom U., Fogelfors, H. (1999). Type and time of autumn tillage with and without herbicides at reduced rates in southern Sweden 2. Weed flora and diversity. *Soil & Tillage Research* 50(3-4): 283-293.
- Bostrom U., Fogelfors, H. (2002). Long-term effects of herbicide-application strategies on weeds and yield in spring-sown cereals. *Weed Science* 50: 196-203.
- Boubakri H., Beuf M., Simonet P., Vogel T.M. (2006). Development of metagenomic DNA shuffling for the construction of a xenobiotic gene. *Gene* 375: 87-94.
- Bouché M.B. (1972). Lombriciens de France. Ecologie et systématique. *Annales de zoologie, écologie animale* 72(2)(n° spécial): 671 p.
- Bowen B.S., Kruse A.D. (1993). Effects of Grazing on Nesting by Upland Sandpipers in South-Central North-Dakota. *Journal of Wildlife Management* 57(2): 291-301.
- Braakhekke W.G., Hooftman D.A.P. (1999). The resource balance hypothesis of plant species diversity in grassland. *Journal of Vegetation Science* 10: 187-200.
- Braband D., Geier U., Kopke U. (2003). Bio-resource evaluation within agri-environmental assessment tools in different European countries. *Agriculture, Ecosystems & Environment* 98(1/3): 423-434
- Braband D., Van Elsen T. (2006). Arable plant species as ecological goods - development of a simple method to identify arablefields that are rich in arable plant species. *Journal of Plant Diseases and Protection*: 535-546.
- Brandenburg R.L. (1985). The effect of field applications of insecticides for variegated cutworm, *Peridroma saucia* (Hubner) (Noctuidae: Lepidoptera) control on non-target arthropods in alfalfa. *Journal of the Kansas Entomological Society* 58(3): 437-441.
- Brauer I. (2003). Money as an indicator: to make use of economic evaluation for biodiversity conservation. *Agriculture, Ecosystems & Environment* 98(1/3): 483-491.
- Bretagnolle V. (2006). Contribution des milieux prairiaux à rotation pluri-annuelle au maintien de la biodiversité en plaine céréalière intensive. Evaluation des surfaces et durées de rotations optimales par modélisation, et validation expérimentale par des mesures agri-environnementales. In *Diva: action publique, agriculture et biodiversité*, (Barre V., Baudry J., eds.), MEDD, Paris: 45-51.
- Bretagnolle V., Inchausti P. (2005). Modelling population reinforcement at a large spatial scale as a conservation strategy for the declining little bustard (*Tetrax tetrax*) in agricultural habitats. *Animal Conservation* 8: 59-68.
- Briemle G. (2006). Careful fertilisation increases species diversity of oligotrophic grassland - results of a study over 22 years in the low mountain range 'Schwabische Alb'. *Naturschutz und Landschaftsplanung* 38(2): 37-44.
- Brin A., Brustel H. (2006). Saproxylic beetles response to cork-oak forests heterogeneity in the Massif des Maures (France). *Revue D Ecologie-La Terre Et La Vie* 61(4): 327-342.
- Britschgi A., Spaar R., Arlettaz R. (2006). Impact of grassland farming intensification on the breeding ecology of an indicator insectivorous passerine, the Whinchat *Saxicola rubetra*: Lessons for overall Alpine meadowland management. *Biological Conservation* 130(2): 193-205.
- Britton A.J., Fisher J.M. (2007). Interactive effects of nitrogen deposition, fire and grazing on diversity and composition of low-alpine prostrate *Calluna vulgaris* heathland. *Journal of Applied Ecology* 44(1): 125-135.
- Brody S.D., Carrasco V., Highfield W. (2003). Evaluating ecosystem management capabilities at the local level in Florida: Identifying policy gaps using geographic information systems. *Environmental Management* 32(6): 661-681.
- Brooks D.R., Clark S.J., Perry J.N., Bohan D.A., Champion G.T., Firbank L.G., Haughton A.J., Hawes C., Heard M.S., Woiwod I.P. (2005). Invertebrate biodiversity in maize following withdrawal of triazine herbicides. *Proceedings of the Royal Society of London. Series B, Biological Sciences* 272(1571): 1497-1502.
- Brosset A., Barbe L., Beaucournu J.C., Faugier C., Salvayre H., Tupinier Y. (1988). La raréfaction du rhinolophe euryale (*Rhinolophus euryale* Blasius) en France: recherche d'une explication. *Mammalia* 52(1): 101-122.
- Brown G.G., Swift M.J., Bennack D.E., Bunning S., Montanez A., Brussaard L. (2007). Management of soil biodiversity in agricultural systems. In *Managing Biodiversity in agricultural systems*, (Jarvis D.I. et al., eds.), Columbia University Press: 224-268.
- Brown M.W. (1993). Resilience of the Natural Arthropod Community on Apple to External Disturbance. *Ecological Entomology* 18(3): 169-183.

- Brown M.W., Adler C.R.L. (1989). Community Structure of Phytophagous Arthropods on Apple. *Environmental Entomology* 18(4): 600-607.
- Brown M.W., Schmitt J.J. (2001). Seasonal and diurnal dynamics of beneficial insect populations in apple orchards under different management intensity. *Environmental Entomology* 30(2): 415-424.
- Brown M.W., Schmitt J.J., Abraham B.J. (2003). Seasonal and diurnal dynamics of spiders (Araneae) in West Virginia orchards and the effect of orchard management on spider communities. *Environmental Entomology* 32(4): 830-839.
- Brown P.D., Morra M.J. (1997). Control of soil-borne plant pest using glucosinolate-containing plants. *Advances in Agronomy* 61: 167-231.
- Broyer J., Lauranson-Broyer J. (1996). Sexual reproduction of plant species in floodplain meadows of the Saone valley (Ain). *Acta Botanica Gallica* 143(4/5): 353-365.
- Broyer J., Prudhomme J. (1995). Incidence de la fertilisation sur la diversité floristique des prairies de fauche inondables dans le val de Saône. *Ecologie* 26(1): 45-58.
- Bruinsma M., Kowalchuk G.A., van Veen J.A. (2003). Effects of genetically modified plants on microbial communities and processes in soil. *Biology and Fertility of Soils* 37(6): 329-337.
- Brussaard L., de Ruiter P.C., Brown G.G. (2007). Soil biodiversity for agricultural sustainability. *Agriculture, Ecosystems & Environment* 121(3): 233-244.
- Brussaard L., Kuiper T.W., Didden W.A.M., de Goede R.G.M., Bloem J. (2004). Biological soil quality from biomass to biodiversity - importance and resilience to management stress and disturbance. In *Managing soil quality: challenges in modern agriculture*, (Schjonning P. et al., eds.), CABI: 139-161.
- Buchanan G.M., Grant M.C., Sanderson R.A., Pearce-Higgins J.W. (2006). The contribution of invertebrate taxa to moorland bird diets and the potential implications of land-use management. *Ibis* 148(4): 615-628.
- Büchs W. (2003). Biotic indicators for biodiversity and sustainable agriculture—introduction and background. *Agriculture, Ecosystems & Environment* 98(1-3): 1-16.
- Büchs W., Harenberg A., Zimmermann J. (1997). The invertebrate ecology of farmland as a mirror of the intensity of the impact of man? - An approach to interpreting results of field experiments carried out in different crop management intensities of a sugar beet and an oil seed rape rotation including set-aside. *Biological Agriculture & Horticulture* 15(1-4): 83-107.
- Buckingham D.L., Peach W.J., Fox D.S. (2006). Effects of agricultural management on the use of lowland grassland by foraging birds. *Agriculture, Ecosystems & Environment* 112(1): 21-40.
- Buckland S.T., Magurran A.E., Green R.E., Fewster R.M. (2005). Monitoring change in biodiversity through composite indices. *Philosophical Transactions of the Royal Society B-Biological Sciences* 360(1454): 243-254.
- Bultmann H. (2005). Strategies and species richness of terricolous lichens in dry sand grasslands. *Tuexenia*(No.25): 425-443.
- Bünemann E.K., Schwenke G.D., Van Zwieten L. (2006). Impact of agricultural inputs on soil organisms - a review. *Australian Journal of Soil Research* 44(4): 379-406.
- Burel F., Baudry J., Butet A., Clergeau P., Delettre Y., le Coeur D., Dubs F., Morvan N., Paillat G., Petit S., Thenail C., Iefeuve J.C. (1998). Comparative biodiversity along a gradient of agricultural landscape. *Acta Oecologica* 19(1): 47-60.
- Burn A.J. (1988). Assessment of the impact of pesticides on invertebrate predation in cereal crops. *Aspects of Applied Biology* 17(1): 279-288.
- Butchart S.H.M., Stattersfield A.J., Bennun L.A., Shutes S.M., Akcakaya H.R., Baillie J.E.M., Stuart S.N., Hilton-Taylor C., Mace G.M. (2004). Measuring global trends in the status of biodiversity: Red list indices for birds. *PLoS Biology* 2(12): 2294-2304.
- Butet A., Leroux A.B.A. (2001). Effects of agriculture development on vole dynamics and conservation of Montagu's harrier in western French wetlands. *Biological Conservation* 100(3): 289-295.
- Butet A., Paillat G., Delettre Y. (2006). Factors driving small rodent assemblages from field boundaries in agricultural landscapes of western France. *Landscape Ecology* 21(3): 449-461.
- Butler S.J., Vickery J.A., Norris K. (2007). Farmland biodiversity and the footprint of agriculture. *Science* 315(5810): 381-384.
- Caballero-Mellado J., Martinez-Romero E. (1999). Soil fertilization limits the genetic diversity of Rhizobium in bean nodules. *Symbiosis* 26(2): 111-121.
- Caesar-TonThat T.C., Caesar A.J., Gaskin J.F., Sainju U.M., Busscher W.J. (2007). Taxonomic diversity of predominant culturable bacteria associated with microaggregates from two different agroecosystems and their ability to aggregate soil in vitro. *Applied Soil Ecology* 36(1): 10-21.
- Canals L.M.I., Clift R., Basson L., Hansen Y., Brandao M. (2006). Expert workshop on land use impacts in life cycle assessment (LCA). *International Journal of Life Cycle Assessment* 11(5): 363-368.
- Canals L.M.I., Romanya J., Cowell S.J. (2007). Method for assessing impacts on life support functions (LSF) related to the use of 'fertile land' in Life Cycle Assessment (LCA). *Journal of Cleaner Production* 15(15): 1426-1440.
- Caprio M.A., Hoy M.A. (1994). Metapopulation dynamics affect resistance development in the predatory mite, *Metaseiulus occidentalis* (Acari: Phytoseiidae). *Journal of Economic Entomology* 87(3): 525-534.
- Carignan V., Villard M.A. (2002). Selecting indicator species to monitor ecological integrity: A review. *Environmental Monitoring and Assessment* 78(1): 45-61.
- Caro T.M., O'Doherty G. (1999). On the use of surrogate species in conservation biology. *Conservation Biology* 13(4): 805-814.
- Carpenter-Boggs L., Kennedy A.C., Reganold J.P. (2000). Organic and biodynamic management: effects on soil biology. *Soil Science Society of America Journal* 64(5): 1651-1659.
- Carson R. (1962). *Silent Spring*, Houghton Mifflin, Boston, 400 p.
- Castello M., Skert N. (2005). Evaluation of lichen diversity as an indicator of environmental quality in the North Adriatic submediterranean region. *Science of the Total Environment* 336(1/3): 201-214.
- Catroux G., Hartmann A., Revellin C. (2001). Trends in rhizobial inoculant production and use. *Plant and Soil* 230(1): 21-30.
- Cavaco M., Goncalves M., Nave A., Santos J., Silvino P., Veiga C., Rodrigues R. (2003). Evaluation of the side effects of five insecticides on predatory mites (Acari: Phytoseiidae) in apple orchards in two different regions of Portugal. *Bulletin OILB/SROP* 26(5): 1-8.
- Ceballos G., Ehrlich P.R. (2002). Mammal population losses and the extinction crisis. *Science* 296(5569): 904-907.
- Cerdeira A.L., Duke S.O. (2006). The current status and environmental impacts of glyphosate-resistant crops: a review. *Journal of Environmental Quality* 35(5): 1633-1658.

- Chabert A., Beaufretton C. (2005). Impact of some agricultural practices on carabidae beetles. *Bulletin OILB/SROP* 28(6): 101-109.
- Chabert A., Gandrey J. (2005). Impact of some insecticides on Carabidae and consequences for slug populations. *Bulletin OILB/SROP* 28(6): 111-113.
- Chaboussou F. (1970). Influence des pesticides sur la plante : conséquences écologiques. *Bulletin de la Société Française d'Ecologie* 3: 146-158.
- Chambon J.P., Cocquempot C., Fougeroux A., Taupin P. (1987). Impact of an early insecticide treatment on the arthropods present in soft winter wheat. *Bulletin OILB/SROP* 10(1): 225-242.
- Chan K.Y. (2001). An overview of some tillage impacts on earthworm population abundance and diversity - implications for functioning in soils. *Soil & Tillage Research* 57(4): 179-191.
- Chape S., Harrison J., Spalding M., Lysenko I. (2005). Measuring the extent and effectiveness of protected areas as an indicator for meeting global biodiversity targets. *Philosophical Transactions of the Royal Society B-Biological Sciences* 360(1454): 443-455.
- Chapin F.S., III (1980). The mineral nutrition of wild plants. *Annual Review of Ecology and Systematics* 11: 233-260.
- Chapin F.S., III, Autumn K., Pugnaire F. (1993). Evolution of suites of traits in response to environmental stress. *American Naturalist* 142: S78-92.
- Chapman M.H., Hoy M.A. (1991). Relative toxicity of *Bacillus thuringiensis* var. *tenebrionis* to the two-spotted spider mite (*Tetranychus urticae* Koch) and its predator *Metaseiulus occidentalis* (Nesbitt) (Acari, Tetranychidae and Phytoseiidae). *Journal of Applied Entomology* 111: 147-154.
- Charleston D.S., Kfir R., Dicke M., Vet L.E.M. (2006). Impact of botanical extracts derived from *Melia azedarach* and *Azadirachta indica* on populations of *Plutella xylostella* and its natural enemies: a field test of laboratory findings. *Biological Control* 39(1): 105-114.
- Charron C.S., Sams C.E., Canaday C.H. (2002). Impact of glucosinolate content in broccoli (*Brassica oleracea* (Italica group)) on growth of *Pseudomonas marginalis*, a causal agent of bacterial soft rot. *Plant Disease* 86(6): 629-632.
- Charudattan R., Walker H.L. (1982). Biological control of weeds with plant pathogens, Wiley and Sons, New York, 293 p.
- Chase M.K., Kristan W.B., Lynam A.J., Price M.V., Rothenberry J.T. (2000). Single species as indicators of species richness and composition in California coastal sage scrub birds and small mammals. *Conservation Biology* 14(2): 474-487.
- Chauvel B., Barralis G., Chadoeuf R., Dessaint F. (1995). Développement de populations adventives en situation de jachère annuelle, 16. *Conférence du COLUMA-Journées Internationales sur la lutte contre les mauvaises herbes*, Reims, (France), ANPP, ed.: 725-734.
- Chen A., Chen Z., Luo K., Miao S. (2003). Effects of some insecticides on field population fluctuation and parasitism of the leafminer parasitoid. *Journal of Yunnan Agricultural University* 18(3): 249-252.
- Chen Y.H., Welter S.C. (2005). Crop domestication disrupts a native tritrophic interaction associated with the sunflower, *Helianthus annuus* (Asterales : Asteraceae). *Ecological Entomology* 30(6): 673-683.
- Chevassus-Au-Louis B. (2007). Biodiversity: A new look at the diversity of life I. Immensity and complexity. *Cahiers Agricultures* 16(3): 219-227.
- Chevre A.M., Eber F., Jenczewski E., Darmency H., Renard M. (2003). Gene flow from oilseed rape to weedy species. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science* 53: 22-25.
- Chu H., Lin X.G., Fujii T., Morimoto S., Yagi K., Hu J., Zhang J. (2007). Soil microbial biomass, dehydrogenase activity, bacterial community structure in response to long-term fertilizer management. *Soil Biology & Biochemistry* 39(11): 2971-2976.
- Chu H.Y., Fujii T., Morimoto S., Lin X.G., Yagi K., Hu J.L., Zhang J.B. (2007). Community structure of ammonia-oxidizing bacteria under long-term application of mineral fertilizer and organic manure in a sandy loam soil. *Applied and Environmental Microbiology* 73(2): 485-491.
- Chust G., Pretus J.L., Ducrot D., Bedos A., Deharveng L. (2003). Identification of landscape units from an insect perspective. *Ecography* 26(3): 257-268.
- Chust G., Pretus J.L., Ducrot D., Bedos A., Deharveng L. (2003). Response of soil fauna to landscape heterogeneity: Determining optimal scales for biodiversity modeling. *Conservation Biology* 17(6): 1712-1723.
- Cibien C., Bideau E., Boisauvert B., Biran H., Angibault J.M. (1995). Seasonal diet and habitat use in field Roe deer (*Capreolus capreolus*) in the Picardie region. *Gibier Faune Sauvage* 12(1): 37-49.
- Cingolani A.M., Cabido M., Gurvich D.E., Renison D., Díaz S. (2007). Filtering processes in the assembly of plant communities: Are species presence and abundance driven by the same traits? *Journal of Vegetation Science* 18: 911-920.
- Cingolani A.M., Noy-Meir I., Díaz S. (2005). Grazing effects on rangeland diversity: a synthesis of contemporary models. *Ecological Applications* 15(2): 757-773.
- Ciotola M., Watson, A.K, Hallet, S.G. (1995). Discovery of an isolate of *Fusarium oxysporum* with potential to control *Striga hermonthica* in Africa. *Weed Research* 35: 303-309.
- Clausen I.H.S. (1995). Review of results from the Danish EPA's Pesticide Research Programme concerning effects on flora and fauna. *SP Rapport - Statens Planteavlsvforsog*(3): 27-37.
- Clergeau P., Jokimaki J., Savard J.P.L. (2001). Are urban bird communities influenced by the bird diversity of adjacent landscapes? *Journal of Applied Ecology* 38(5): 1122-1135.
- Clergue B., Amiaud B., Pervanchon F., Lasserre-Joulin F., Plantureux S. (2005). Biodiversity: function and assessment in agricultural areas. A review. *Agronomy for Sustainable Development* 25: 1-15.
- Cloquell-Ballester V.A., Cloquell-Ballester V.A., Monerde-Diaz R., Santamarina-Siurana M.C. (2006). Indicators validation for the improvement of environmental and social impact quantitative assessment. *Environmental Impact Assessment Review* 26(1): 79-105.
- Cluzeau D., Hallaire V., Bodet J.-M. (2001). Le rôle des lombriciens sur le fonctionnement des sols. Impacts des pratiques agricoles. In Du labour au semis direct: enjeux agronomiques, INRA Dossiers, INRA, Paris: 10-13.
- Cocquempot C., Chambon J.P., Reynaud P., Fischer L. (1991). Incidence of insecticide treatments on the aranean fauna of a wheat field at flowering. *Agronomie* 11(5): 423-434.
- Codron J.M., Jacquet F., Habib R., Sauphanor B. (2003). Bilan et perspectives environnementales de la filière arboriculture fruitière. *Les Dossiers de l'Environnement de l'INRA* 23: 31-67.

- Cohen M.F., Yamasaki H., Mazzola M. (2005). Brassica napus seed meal soil amendment modifies microbial community structure, nitric oxide production and incidence of Rhizoctonia root rot. *Soil Biology & Biochemistry* 37(7): 1215-1227.
- Colbach N., Molinari N., Meynard J.M., Messéan A. (2005). Spatial aspects of gene flow between rapeseed varieties and volunteers. *Agronomy for Sustainable Development* 25(3): 355-368.
- Cole L., Bradford M.A., Shaw P.J.A., Bardgett R.D. (2006). The abundance, richness and functional role of soil meso- and macrofauna in temperate. *Agriculture, Ecosystems & Environment* 33(2): 186-198.
- Cole L., Buckland S.M., Bardgett R.D. (2005). Relating microarthropod community structure and diversity to soil fertility manipulations in temperate grassland. *Soil Biology & Biochemistry* 37(9): 1707-1717.
- Coll M., Bottrell D.G. (1996). Movement of an insect parasitoid in simple and diverse plant assemblages. *Ecological Entomology* 21(2): 141-149.
- Cook R., Baker K.F. (1983). The nature and practice of biological control of plant pathogens, Am. Phytopathol. Soc., St Paul, Minnesota, 539 p.
- Cooke A.S., Greig-Smith P.W., Jones S. (1992). Consequences for vertebrates wildlife of toxic residues in earthworm prey. In *Ecotoxicology of earthworms*, (Greig-Smith P.W. et al., eds.), Intercept, Andover, Hampshire: 139-155.
- Cortet J., Andersen M.N., Caul S., Griffiths B., Joffre R., Lacroix B., Sausse C., Thompson J., Krogh P.H. (2006). Decomposition processes under Bt (*Bacillus thuringiensis*) maize: Results of a multi-site experiment. *Soil Biology & Biochemistry* 38(1): 195-199.
- Cortet J., Griffiths B.S., Bohanec M., Demsar D., Andersen M.N., Caul S., Birch A.N.E., Pernin C., Tabone E., de Vaufléury A., Ke X., Krogh P.H. (2007). Evaluation of effects of transgenic Bt maize on microarthropods in a European multi-site experiment. *Pedobiologia* 51(3): 207-218.
- Cottrell T.E. (2005). Predation and cannibalism of lady beetle eggs by adult lady beetles. *Biological Control* 34(2): 159-164.
- Coulon A., Guillot G., Cosson J.F., Angibault J.M.A., Aulagnier S., Cargnelutti B., Galan M., Hewison A.J.M. (2006). Genetic structure is influenced by landscape features: empirical evidence from a roe deer population. *Molecular Ecology* 15(6): 1669-1679.
- Coulon J.B., Delacroix-Buchet A., Martin B., Pirisi A. (2004). Relationships between ruminant management and sensory characteristics of cheeses: a review. *Lait* 84(3): 221-241.
- Coulson S.J., Bullock J.M., Stevenson M.J., Pywell R.F. (2001). Colonization of grassland by sown species: dispersal versus microsite limitation in responses to management. *Journal of Applied Ecology* 38(1): 204-216.
- Cousins S., Lavorel S., Davies I.D. (2003). Modelling the effects of landscape pattern and grazing regimes on the persistence of plant species with high conservation value in grasslands in south-eastern Sweden. *Landscape Ecology* 18(3): 315-332.
- Cousins S.A.O. (2006). Plant species richness in midfield islets and road verges - The effect of landscape fragmentation. *Biological Conservation* 127(4): 500-509.
- Cousins S.A.O., Eriksson O. (2001). Plant species occurrences in a rural hemiboreal landscape: effects of remnant habitats, site history, topography and soil. *Ecography* 24(4): 461-469.
- Couty A., Vina G.d.I., Clark S.J., Kaiser L., Pham-Delegue M.H., Poppy G.M. (2001). Direct and indirect sublethal effects of *Galanthus nivalis* agglutinin (GNA) on the development of a potato-aphid parasitoid, *Aphelinus abdominalis* (Hymenoptera: Aphelinidae). *Journal of Insect Physiology* 47(6): 553-561.
- Coventry E., Noble R., Mead A., Whipps J.M. (2005). Suppression of *Allium* white rot (*Sclerotium cepivorum*) in different soils using vegetable wastes. *European Journal of Plant Pathology* 111(2): 101-112.
- Crawley M.J., Johnston A.E., Silvertown J., Dodd M., de Mazencourt C., Heard M.S., Henman D.F., Edwards G.R. (2005). Determinants of species richness in the Park Grass Experiment. *American Naturalist* 165: 179-192.
- Cremene C., Groza G., Rakosy L., Schileyko A.A., Baur A., Erhardt A., Baur B. (2005). Alterations of steppe-like grasslands in Eastern Europe: a threat to regional biodiversity hotspots. *Conservation Biology* 19(5): 1606-1618.
- Crist T.O., Veech J.A., Gering J.C., Summerville K.S. (2003). Partitioning species diversity across landscapes and regions: a hierarchical analysis of α , β , and γ -diversity. *American Naturalist* 162: 734-743.
- Critchley C.N.R., Walker K.J., Pywell R.F., Stevenson M.J. (2007). The contribution of English agri-environment schemes to botanical diversity in arable field margins. *Aspects of Applied Biology* No.81: 293-300.
- Cross J.V., Solomon M.G., Babandriker D., Blommers L., Easterbrook M.A., Jay C.N., Jenser G., Jolly R.L., Kuhlmann U., Lilley R., Olivella E., Toepfer S., Vidal S. (1999). Biocontrol of pests of apples and pears in northern and central Europe: 2. Parasitoids. *Biocontrol Science and Technology* 9(3): 277-314.
- Crossley D.A., Mueller B.R., Perdue J.C. (1992). Biodiversity of Microarthropods in Agricultural Soils - Relations to Processes. *Agriculture, Ecosystems & Environment* 40(1-4): 37-46.
- Cugnasse J.M., Gibert P., Bouchet J. (1998). Hybridation, en nature, de mouflons méditerranéens mâles avec des brebis domestiques. *Bulletin mensuel O.N.C.* 236: 24-29.
- Cunningham H.M., Chaney K., Bradbury R.B., Wilcox A. (2004). Non-inversion tillage and farmland birds: a review with special reference to the UK and Europe. *Ibis* 146(2): 192-202.
- Cureton A.N., Newbury H.J., Raybould A.F., Ford-Lloyd B.V. (2006). Genetic structure and gene flow in wild beet populations: the potential influence of habitat on transgene spread and risk assessment. *Journal of Applied Ecology* 43: 1203-1212.
- Curry J.P. (1994). Grassland invertebrates. Ecology, influence on soil fertility and effects on plant growth, Chapman & Hall, London, 437 p.
- Curry J.P., Good J.A. (1992). Soil faunal degradation and restoration. In *Advances in soil science*, (Lal R., Steward B.A., eds.), 17, Springer-Verlag, New York: 171-215.
- Dalmastri C., Chiarini L., Cantale C., Bevivino A., Tabacchioni S. (1999). Soil type and maize cultivar affect the genetic diversity of maize root-associated *Burkholderia cepacia* populations. *Microbial Ecology* 38(3): 273-284.
- Danell K., Huss-Danell K. (1985). Feeding by insects and hares on birches earlier affected by moose browsing. *Oikos* 44(1): 75-81.
- Danks H.V. (1997). Assessing insect biodiversity - without wasting your time. *Global Biodiversity* 7(3): 17-21.
- Darmency H., Gasquez, J. (1990). Résistance aux herbicides chez les mauvaises herbes. *Agronomie* 6: 457-472.
- Darmency H. (2005). Effets non intentionnels des cultures transgéniques. In *Enjeux phytosanitaires pour l'Agriculture et l'Environnement*, (Regnault-Roger C., ed.), Tec et Doc Lavoisier, Paris: 411-430.

- Darmency H., Lefol E., Fleury A. (1998). Spontaneous hybridizations between oilseed rape and wild radish. *Molecular Ecology* 7: 1467-1473.
- Dauber J., Bengtsson J., Lenoir L. (2006). Evaluating effects of habitat loss and land-use continuity on ant species richness in seminatural grassland remnants. *Conservation Biology* 20(4): 1150-1160.
- Davis A.L.V., Scholtz C.H., Dooley P.W., Bham N., Kryger U. (2004). Scarabaeine dung beetles as indicators of biodiversity, habitat transformation and pest control chemicals in agro-ecosystems. *South African Journal of Science* 100(9/10): 415-424.
- de Bello F. (2005). Predictive value of plant traits to grazing along a climatic gradient in the Mediterranean. *Journal of Applied Ecology* 42(5): 824-833.
- de Bello F., Lepš J., Sebastia M.-T. (2006). Variations in species and functional plant diversity along climatic and grazing gradients. *Ecography* 29: 801-810.
- de Goede R.G.M., Brussaard L., Akkermans A.D.L. (2003). On-farm impact of cattle slurry manure management on biological soil quality. *Njas-Wageningen Journal of Life Sciences* 51(1-2): 103-133.
- de Gournay X. (1963). La lutte contre le vulpin des champs (*Alopecurus myosuroides* Huds.) dans les cultures de blé d'hiver. *Annales de Physiologie Végétale* 5(3): 229-247.
- de la Poza M., Pons X., Farinos G.P., Lopez C., Ortego F., Eizaguirre M., Castanera P., Albajes R. (2005). Impact of farm-scale Bt maize on abundance of predatory arthropods in Spain. *Crop Protection* 24(7): 677-684.
- de Snoo G., R. (1997). Arable flora in sprayed and unsprayed crop edges. *Agriculture Ecosystems and Environment* 66: 223-230.
- de Snoo G.R. (1999). Unsprayed field margins: effects on environment, biodiversity and agricultural practice. *Landscape and Urban Planning* 46(1-3): 151-160.
- de Vaufléury A., Kramarz P.E., Binet P., Cortet J., Caul S., Andersen M.N., Plumey E., Coeurdassier M., Krogh P.H. (2007). Exposure and effects assessments of Bt-maize on non-target organisms (gastropods, microarthropods, mycorrhizal fungi) in microcosms. *Pedobiologia* 51(3): 185-194.
- Deacon L.J., Pryce-Miller E.J., Frankland J.C., Bainbridge B.W., Moore P.D., Robinson C.H. (2006). Diversity and function of decomposer fungi from a grassland soil. *Soil Biology & Biochemistry* 38(1): 7-20.
- DeBach P., Rosen D. (1991). Biological control by natural enemies, 2 ed., Cambridge University Press, Cambridge, 456 p.
- Debaeke P., Orlando D. (1991). Simplification du travail du sol et évolution de la flore adventice : conséquences pour le désherbage à l'échelle de la rotation. In Simplification du travail du sol., (INRA, ed.): 35-62.
- Debeljak M., Cortet J., Demsar D., Krogh P.H., Dzeroski S. (2007). Hierarchical classification of environmental factors and agricultural practices affecting soil fauna under cropping systems using Bt maize. *Pedobiologia* 51(3): 229-238.
- Debinski D., Holt R.D. (2000). A survey and overview of habitat fragmentation experiments *Conservation Biology* 14(2): 342-355.
- Debras J.F., Torre F., Rieux R., Kreiter S., Garcin M.S., Van Helden M., Buisson E., Dutoit T. (2006). Discrimination between agricultural management and the hedge effect in pear orchards (south-eastern France). *Annals of Applied Biology* 149(3): 347-355.
- Debussche M., Escaré J., Lepart J., Houssard C., Lavorel S. (1996). Changes in Mediterranean plant succession: old-fields revisited. *Journal of Vegetation Science* 7: 519-526.
- Debussche M., Lepart J., Dervieux A. (1999). Mediterranean landscape changes: evidence from old postcards. *Global Ecology and Biogeography* 8: 3-15.
- Decaens T., Dutoit T., Alard D. (1997). Earthworm community characteristics during afforestation of abandoned chalk grasslands (Upper Normandy, France). *European Journal of Soil Biology* 33(1): 1-11.
- Decaens T., Dutoit T., Alard D., Lavelle P. (1998). Factors influencing soil macrofaunal communities in post-pastoral successions of western France. *Applied Soil Ecology* 9(1-3): 361-367.
- Decaens T., Jimenez J.J., Gioia C., Measey G.J., Lavelle P. (2006). The values of soil animals for conservation biology. *European Journal of Soil Biology* 42: S23-S38.
- Deckers B., De Becker P., Honnay O., Hermy M., Muys B. (2005). Sunken roads as habitats for forest plant species in a dynamic agricultural landscape: effects of age and isolation. *Journal of Biogeography* 32(1): 99-109.
- Deckers B., Hermy M., Muys B. (2004). Factors affecting plant species composition of hedgerows: relative importance and hierarchy *Acta Oecologica* 26(1): 23-37.
- Decourtye A., Lecompte P., Pierre J., Chauzat M.-P., Thiébeau P. (2007). Introduction de jachères florales en zones de grandes cultures : comment mieux concilier agriculture, biodiversité et apiculture ? *Le Courrier de l'environnement de l'INRA* 54: 33-56.
- deGroot R.S., Ketner P., Ova A.H. (1995). Selection and use of bio-indicators to assess the possible effects of climate change in Europe. *Journal of Biogeography* 22(4-5): 935-943.
- Delbaere B., Serradilla A.N.E. (2004). Environmental risks from agriculture in Europe: Locating environmental risk zones in Europe using agri-environmental indicators, EC NC-European Centre for Nature Conservation, Tilburg, The Netherlands, 184 p.
- Delpuech J.M., Meyet J. (2003). Reduction in the sex ratio of the progeny of a parasitoid wasp (*Trichogramma brassicae*) surviving the insecticide chlorpyrifos. *Archives of Environmental Contamination and Toxicology* 45(2): 203-208.
- Demanou J., Sharma S., Dorfler U., Schroll R., Pritsch K., Njine T., Bausenwein U., Monkiedje A., Munch J.C., Schloter M. (2006). Structural and functional diversity of soil microbial communities as a result of combined applications of copper and mefenoxam. *Soil Biology & Biochemistry* 38(8): 2381-2389.
- Dennis P., Doering J., Stockan J.A., Jones J.R., Rees M.E., Vale J.E., Sibbald A.R. (2004). Consequences for biodiversity of reducing inputs to upland temperate pastures: effects on beetles (Coleoptera) of cessation of nitrogen fertilizer application and reductions in stocking rates of sheep. *Grass & Forage Science* 59(2): 121-135.
- Dennis P., Fry G.L.A., Thomas M.B. (1993). The effects of reduced doses of insecticide on aphids and their natural enemies in oats. *Norwegian Journal of Agricultural Sciences* 7(3/4): 311-325.
- Dennis P., Young M.R., Bentley C. (2001). The effects of varied grazing management on epigeal spiders, harvestmen and pseudoscorpions of *Nardus stricta* grassland in upland Scotland. *Agriculture, Ecosystems & Environment* 86(1): 39-57.
- Dennis P., Young M.R., Gordon I.J. (1998). Distribution and abundance of small insects and arachnids in relation to structural heterogeneity of grazed, indigenous grasslands. *Ecological Entomology* 23(3): 253-264.

- Dennis P., Young M.R., Howard C.L., Gordon I.J. (1997). The response of epigeal beetles (Col: Carabidae, Staphylinidae) to varied grazing regimes on upland *Nardus stricta* grasslands. *Journal of Applied Ecology* 34(2): 433-443.
- Denys C., Tschamtké T. (2002). Plant-insect communities and predator-prey ratios in field margin strips, adjacent crop fields, and fallows. *Oecologia* 130(2): 315-324.
- Depret G., Houot S., Allard M.R., Breuil M.C., Nouaim R., Laguerre G. (2004). Long-term effects of crop management on *Rhizobium leguminosarum* biovar *viciae* populations. *FEMS Microbiology Ecology* 51(1): 87-97.
- Desneux N., Fauvergue X., Dechaume-Moncharmont F.X., Kerhoas L., Ballanger Y., Kaiser L. (2005). *Diaeretiella rapae* limits *Myzus persicae* populations after applications of deltamethrin in oilseed rape. *Journal of Economic Entomology* 98(1): 9-17.
- Desriers M. (2007). L'agriculture française depuis 50 ans: des petites exploitations familiales aux droits à paiement unique. *Agreste Cahiers* 2: 3-14.
- Devare M., Londono-R L.M., Thies J.E. (2007). Neither transgenic Bt maize (MON863) nor tefluthrin insecticide adversely affect soil microbial activity or biomass: A 3-year field analysis. *Soil Biology & Biochemistry* 39(8): 2038-2047.
- Devers M., Henry S., Hartmann A., Martin-Laurent F. (2005). Horizontal gene transfer of atrazine-degrading genes (*atz*) from *Agrobacterium tumefaciens* St96-4 pADP1 : Tn5 to bacteria of maize-cultivated soil. *Pest Management Science* 61(9): 870-880.
- Devictor V., Jiguet F. (2007). Community richness and stability in agricultural landscapes : The importance of surrounding habitats. *Agriculture, Ecosystems & Environment* 120(2-4): 179-184.
- Devillers J., Farret R., Girardin P., Rivière J.-L., Soulas G. (2005). Indicateurs pour évaluer les risques liés à l'utilisation des pesticides, Tec & Doc, Lavoisier, 278 p.
- Di Giulio M., Edwards P.J., Meister E. (2001). Enhancing insect diversity in agricultural grasslands: the roles of management and landscape structure. *Journal of Applied Ecology* 38(2): 310-319.
- Díaz S., Cabido M., Casanoves F. (1999). Functional implications of trait-environment linkages in plant communities. In *Ecological Assembly Rules*, (Weiher E., Keddy P., eds.), 12, Cambridge University Press: 338-362.
- Díaz S., Fargione J., Chapin F.S., Tilman D. (2006). Biodiversity loss threatens human well-being. *Plos Biology* 4(8): e277.
- Díaz S., Lavorel S., McIntyre S., Falczuk V., Casanoves F., Milchunas D.G., Skarpe C., Rusch G., Sternberg M., Noy-Meir I., Landsberg J., Zhang W., Clark H., Campbell B.D. (2007). Plant trait responses to grazing - A global synthesis. *Global Change Biology* 13: 313-341.
- Díaz S., Noy-Meir I., Cabido M. (2001). Can grazing response of herbaceous plants be predicted from simple vegetative traits? *Journal of Applied Ecology* 38: 497-508.
- Dicke M. (1999). Direct and indirect effects of plants on performance of beneficial organisms. In *Handbook of Pest Management* (Ruberson J.R., ed.), Marcel Dekker, New York: 105-153.
- Didden W., Fründ H.C., Graefe U. (1997). Enchytraeids. In *Fauna in soil ecosystems: recycling, processes, nutrient fluxes and agricultural production*, (Benckiser G., ed.), Marcel Dekker: 135-172.
- Didden W., Rombke J. (2001). Enchytraeids as indicator organisms for chemical stress in terrestrial ecosystems. *Ecotoxicology and Environmental Safety* 50(1): 25-43.
- Dierschke J., Bairlein F. (2004). Habitat selection of wintering passerines in salt marshes of the German Wadden Sea. *Journal of Ornithology* 145(1): 48-58.
- Dierssen K. (2006). Indicating botanical diversity - Structural and functional aspects based on case studies from Northern Germany. *Ecological Indicators* 6(1): 94-103.
- Dietschi S. (2005). Botanical Diversität in Mähwiesen in der Berglandwirtschaft, Ergebnisse und Bewertung handlungsorientierter Vertragsbewirtschaftung, *Phil.Alp 2005, Die Alpen aus der Sicht junger Forschender*, Berne, 2005/06/09-10, Commission interacadémique de recherche alpine C., ed., PNR 49.
- Ding P., Liu F.X., Xu C.X., Wang K.R. (2007). Transmission of *Cryphonectria hypovirus* to protect chestnut trees from chestnut blight disease. *Biological Control* 40(1): 9-14.
- Dinter A., Wiles J.A. (2000). Safety of the new DuPont insecticide "Indoxacarb" to beneficial arthropods: an overview. *Bulletin OILB/SROP* 23(9): 149-156.
- Dollacker A. (2006). Conserving biodiversity alongside agricultural profitability through integrated R&D approaches and responsible use of crop protection products. *Pflanzenschutz-Nachrichten Bayer* 59(1): 117-134.
- Dolphin K., Quicke D.L.J. (2001). Estimating the global species richness of an incompletely described taxon: an example using parasitoid wasps (Hymenoptera : Braconidae). *Biological Journal of the Linnean Society* 73(3): 279-286.
- Donnelly A., Jennings E., Mooney P., Finnan J., Lynn D., Jones M., O'Mahony T., Therivel R., Byrne G. (2006). Workshop approach to developing objectives, targets and indicators for use in SEA. *Journal of Environmental Assessment Policy and Management* 8(2): 135-156.
- Donnison L.M., Griffith G.S., Hedger J., Hobbs P.J., Bardgett R.D. (2000). Management influences on soil microbial communities and their function in botanically diverse haymeadows of northern England and Wales. *Soil Biology & Biochemistry* 32(2): 253-263.
- Doring T.F., Hiller A., Wehke S., Schulte G., Broll G. (2003). Biotic indicators of carabid species richness on organically and conventionally managed arable fields. *Agriculture, Ecosystems & Environment* 98(1/3): 133-139.
- Dorrough J., Moll J., Crosthwaite J. (2007). Can intensification of temperate Australian livestock production systems save land for native biodiversity? *Agriculture, Ecosystems & Environment* 121(3): 222-232.
- Dover J., Sparks T., Clarke S., Gobbett K., Glossop S. (2000). Linear features and butterflies: the importance of green lanes. *Agriculture, Ecosystems & Environment* 80(3): 227-242.
- Downs N.C., Racey P.A. (2006). The use by bats of habitat features in mixed farmland in Scotland. *Acta Chiropterologica* 8(1): 169-185.
- Duelli P., Obrist M.K. (1998). In search of the best correlates for local organismal biodiversity in cultivated areas. *Biodiversity and Conservation* 7: 297-309.
- Duelli P., Obrist M.K. (2003). Biodiversity indicators: the choice of values and measures. *Agriculture, Ecosystems & Environment* 98(1-3): 87-98.
- Duffield S. (1991). Does the size of area you spray with insecticides influence the extent of side-effects on invertebrate populations? *Game Conservancy Review*(22): 64-65.

- Duffield S.J., Bryson R.J., Young J.E.B., Sylvester-Bradley R., Scott R.K. (1997). The influence of nitrogen fertiliser on the population development of the cereal aphids *Sitobion avenae* (F.) and *Metopolophium dirhodum* (Wlk.) on field grown winter wheat. *Annals of Applied Biology* 130(1): 13-26.
- Duijff B.J., Recorbet G., Bakker P.A.H.M., Loper J.E., Lemanceau P. (1999). Microbial antagonism at the root level is involved in the suppression of *Fusarium* wilt by the combination of nonpathogenic *Fusarium oxysporum* Fo47 and *Pseudomonas putida* WCS358. *Phytopathology* 89(11): 1073-1079.
- Dunfield K.E., Germida J.J. (2001). Diversity of bacterial communities in the rhizosphere and root interior of field-grown genetically modified *Brassica napus*. *Fems Microbiology Ecology* 38(1): 1-9.
- Dupré C., Ehlén J. (2002). Habitat configuration, species traits and plant distributions. *Journal of Ecology* 90(5): 796-805.
- Durant D., Tichit M., Kerneis E., Fritz H. (2008). Management of agricultural grasslands from breeding waders: integrating ecological and livestock system perspectives - a review. *Biodiversity and Conservation* 17(9): 2275-2295.
- Duro D., Coops N.C., Wulder M.A., Han T. (2007). Development of a large area biodiversity monitoring system driven by remote sensing. *Progress in Physical Geography* 31(3): 235-260.
- Eckert H., Breitschuh G., Sauerbeck D.R. (2000). Criteria and standards for sustainable agriculture. *Journal of Plant Nutrition and Soil Science-Zeitschrift Fur Pflanzenernahrung Und Bodenkunde* 163(4): 337-351.
- Edwards C.A., Loft J.R. (1977). *Biology of earthworms* (2nd edition), Chapman & Hall, London, 352 p.
- EEA (2002). An inventory of biodiversity indicators in Europe, 2002, Technical report, n°92: 2 p. http://reports.eea.europa.eu/technical_report_2004_92/en/Technical92_for_web.pdf
- EEA (2007). Halting the loss of biodiversity by 2010: proposal for a first set of indicators to monitor progress in Europe, Technical report No 11/2007: 186 p. http://reports.eea.europa.eu/technical_report_2007_11/en/Tech_report_11_2007_SEBI.pdf
- EEA, OPOCE (2005). Agriculture et Environment in EU 15 - The IRENA indicator report, EEA Report, n°6/2005: 128 p. http://reports.eea.europa.eu/eea_report_2005_6/en/EEA_report_6_2005.pdf
- Eggleton P., Vanbergen A.J., Jones D.T., Lambert M.C., Rockett C., Hammond P.M., Beccaloni J., Marriott D., Ross E., Giusti A. (2005). Assemblages of soil macrofauna across a Scottish land-use intensification gradient: influences of habitat quality, heterogeneity and area. *Journal of Applied Ecology* 42(6): 1153-1164.
- El Titi A. (2003). Effects of tillage on invertebrates in soil ecosystems. In *Soil tillage in Agroecosystems*, (El-Titi A., ed.), CRC Press, London: 261-296.
- El Titi A., Ipach U. (1989). Soil fauna in sustainable agriculture: results of an integrated farming system at Lautenbach, F.R.G. *Agriculture, Ecosystems & Environment* 27(1-4): 561-572.
- El-Ghar G.E.S.A., El-Sayed A.E.M., El-Shiekh A.E., Radwan H.S.A. (1994). Field tests with insecticides and insect growth regulators to control insect pests of cowpea and its effects on certain beneficial insects. *Archives of Phytopathology and Plant Protection* 28(6): 531-543.
- Elhottova D., Szili-Kovacs T., Triska J. (2002). Soil microbial community of abandoned sand fields. *Folia Microbiologica* 47(4): 435-440.
- Elisseou G.K., Veresoglou D.S., Mamolos A.P. (1995). Vegetation productivity and diversity of acid grasslands in Northern Greece as influenced by winter rainfall and limiting nutrients. *Acta Oecologica* 16(6): 687-702.
- Epstein D.L., Zack R.S., Brunner J.F., Gut L., Brown J.J. (2000). Effects of broad-spectrum insecticides on epigeal arthropod biodiversity in Pacific Northwest apple orchards. *Environmental Entomology* 29(2): 340-348.
- Eriksson O., Cousins S.A.O., Bruun H.H. (2002). Land use history and fragmentation of traditionally managed grasslands in Scandinavia. *Journal of Vegetation Science* 13: 743-748.
- Erkilić L., Uygun N. (1997). Studies on the effects of some pesticides on white peach scale, *Pseudaulacaspis pentagona* (Targ.-Tozz.) (Homoptera: Diaspididae) and its side-effects on two common scale insect predators. *Crop Protection* 16(1): 69-72.
- Ernoul A., Tremauville Y., Cellier D., Margerie P., Langlois E., Alard D. (2006). Potential landscape drivers of biodiversity components in a flood plain: Past or present patterns? *Biological Conservation* 127(1): 1-17.
- Escarré J., Houssard C., Debussche M. (1983). Evolution de la végétation et du sol après abandon cultural en région méditerranéenne : étude de succession dans les Garrigues du Montpelliérais (France). *Acta Oecologica, Oecologia Plantarum* 4(18): 221-239.
- Escher N., Käch B., Nentwig W. (2000). Decomposition of transgenic *Bacillus thuringiensis* maize by microorganisms and woodlice *Porcellio scaber* (Crustacea: Isopoda). *Basic and Applied Ecology* 1(2): 161-169.
- Esselink P., Zijlstra W., Dijkema K.S., van Diggelen R. (2000). The effects of decreased management on plant-species distribution patterns in a salt marsh nature reserve in the Wadden Sea. *Biological Conservation* 93(1): 61-76.
- Evans D.M., Redpath S.M., Elston D.A., Evans S.A., Mitchell R.J., Dennis P. (2006). To graze or not to graze? Sheep, voles, forestry and nature conservation in the British uplands. *Journal of Applied Ecology* 43(3): 499-505.
- Evans D.M., Redpath S.M., Evans S.A., Elston D.A., Dennis P. (2005). Livestock grazing affects the egg size of an insectivorous passerine. *Biology Letters* 1(3): 322-325.
- Evans D.M., Redpath S.M., Evans S.A., Elston D.A., Gardner C.J., Dennis P., Pakeman R.J. (2006). Low intensity, mixed livestock grazing improves the breeding abundance of a common insectivorous passerine. *Biology Letters* 2(4): 636-638.
- Ewald J., Aebischer T. (2000). Trends in pesticide use and efficacy during 26 years of changing agriculture in Southern England. *Environmental Monitoring and Assessment* 64: 493-529.
- Fabre B., Kockmann F. (2006). Effects of liming on permanent pastures and on long-duration pastures. A bibliographical review. *Fourrages*(No.185): 103-122.
- Fabre F., Pierre J.S., Dedryver C.A., Plantegenest M. (2006). Barley yellow dwarf disease risk assessment based on Bayesian modelling of aphid population dynamics. *Ecological Modelling* 193(3-4): 457-466.
- Fahrig L. (2003). Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology Evolution and Systematics* 34: 487-515.
- Fahrig L., Merriam H.G. (1985). Habitat patch connectivity and population survival. *Ecology* 66: 1762-1768.
- Fairbanks D.H.K. (2004). Regional land-use impacts affecting avian richness patterns in Southern Africa-insights from historical avian atlas data. *Agriculture, Ecosystems & Environment* 101(2/3): 269-288.

- Farrar R.R., Jr., Barbour J.D., Kennedy G.G. (1994). Field evaluation of insect resistance in a wild tomato and its effects on insect parasitoids. *Entomologia Experimentalis et Applicata* 71(3): 211-226.
- Fauvel G. (1999). Diversity of Heteroptera in agroecosystems: role of sustainability and bioindication. *Agriculture, Ecosystems & Environment* 74(1-3): 275-303.
- Favreau J.M., Drew C.A., Hess G.R., Rubino M.J., Koch F.H., Eschelbach K.A. (2006). Recommendations for assessing the effectiveness of surrogate species approaches. *Biodiversity and Conservation* 15: 3949-3969.
- Fédoroff E., Ponge J.F., Dubs F., Fernandes-Gonzalez F., Lavelle P. (2005). Small-scale response of plant species to land-use intensification. *Agriculture, Ecosystems & Environment* 105(1-2): 283-290.
- Ferris H., Carlson H.L., Westerdahl B.B. (1994). Nematode Population-Changes under Crop-Rotation Sequences - Consequences for Potato Production. *Agronomy Journal* 86(2): 340-348.
- Ferro D.N. (1993). Potential for resistance to *Bacillus thuringiensis*: Colorado potato beetle (Coleoptera: Chrysomelidae) - a model system. *American Entomologist* 39(1): 38-44.
- Filser J., Fromm H., Nagel R.F., Winter K. (1995). Effects of Previous Intensive Agricultural Management on Microorganisms and the Biodiversity of Soil Fauna. *Plant and Soil* 170(1): 123-129.
- Finke D.L., Denno R.F. (2006). Spatial refuge from intraguild predation: implications for prey suppression and trophic cascades. *Oecologia* 149(2): 265-275.
- Firbank L.G., Barr C.J., Bunce R.G.H., Furse M.T., Haines-Young R., Hornung M., Howard D.C., Sheail J., Sier A., Smart S.M. (2003). Assessing stock and change in land cover and biodiversity in GB: an introduction to Countryside Survey 2000. *Journal of Environmental Management* 67(3, Special issue): 207-218.
- Firbank L.G., Smart S.M., Crabb J., Critchley C.N.R., Fowbert J.W., Fuller R.J., Gladders P., Green D.B., Henderson I., Hill M.O. (2003). Agronomic and ecological costs and benefits of set-aside in England. *Agriculture Ecosystems & Environment* 95(1): 73-85.
- Fischer M., Wipf S. (2002). Effect of low-intensity grazing on the species-rich vegetation of traditionally mown subalpine meadows. *Biological Conservation* 104(1): 1-11.
- Flather C.H., Bevers M. (2002). Patchy reaction-diffusion and population abundance: the relative importance of habitat amount and arrangement. *American Naturalist* 159(1): 40-56.
- Fleishman E., Thomson J.R., Mac Nally R., Murphy D.D., Fay J.P. (2005). Using indicator species to predict species richness of multiple taxonomic groups. *Conservation Biology* 19(4): 1125-1137.
- Fliessbach A., Oberholzer H.R., Gunst L., Mader P. (2007). Soil organic matter and biological soil quality indicators after 21 years of organic and conventional farming. *Agriculture Ecosystems & Environment* 118(1-4): 273-284.
- Floate K.D. (2006). Endectocide use in cattle and fecal residues: environmental effects in Canada. *Canadian Journal of Veterinary Research-Revue Canadienne De Recherche Veterinaire* 70(1): 1-10.
- Floate K.D., Colwell D.D., Fox A.S. (2002). Reductions of non-pest insects in dung of cattle treated with endectocides: a comparison of four products. *Bulletin of Entomological Research* 92(6): 471-481.
- Flores S., Saxena D., Stotzky G. (2005). Transgenic Bt plants decompose less in soil than non-Bt plants. *Soil Biology & Biochemistry* 37(6): 1073-1082.
- Foissner W. (1997). Protozoa as bioindicators in agroecosystems, with emphasis on farming practices, biocides, and biodiversity. *Agriculture, Ecosystems & Environment* 62(2-3): 93-103.
- Foley J.A., DeFries R., Asner G.P., Barford C., Bonan G., Carpenter S.R., Chapin F.S.I., Coe M.T., Daily G.C., Gibbs H.K., Helkowski J.H., Holloway T., Howard E.A., Kucharik C.J., Monfreda C., Patz J.A., Prentice I.C., Ramankutty N., Snyder P.K. (2005). Global consequences of land use. *Science* 309: 570-574.
- Folgarait P.J. (1998). Ant biodiversity and its relationship to ecosystem functioning: a review. *Biodiversity and Conservation* 7(9): 1221-1244.
- Fontaine A.L., Kennedy P.L., Johnson D.H. (2004). Effects of distance from cattle water developments on grassland birds. *Journal of Range Management* 57(3): 238-242.
- Foote A.L., Hornung C.L.R. (2005). Odonates as biological indicators of grazing effects on Canadian prairie wetlands. *Ecological Entomology* 30(3): 273-283.
- Forti D., Ioriatti C., Angeli G., Catoni M. (1994). Two new acaricides, pyridaben and fenpyroximate: evaluation of their efficacy on *Panonychus ulmi* (Koch) and *Aculus schlechtendali* (Nal.) and their nontarget effects on beneficial arthropods. *Informatore Fitopatologico* 44(7-8): 38-42.
- Foster B.L., Gross K.L. (1998). Species richness in a successional grassland: effects of nitrogen enrichment and plant litter. *Ecology* 79(8): 2593-2602.
- Fournier-Chambillon C., Berny P.J., Coiffier P., Barbedienne P., Dasse B., Delas G., Galineau H., Mazet A., Pouzenc P., Rosoux R., Fournier P. (2004). Evidence of secondary poisoning of free-ranging riparian mustelids by anticoagulant rodenticides in France: implications for conservation of European Mink (*Mustela lutreola*). *Journal of Wildlife Diseases* 40(4): 688-695.
- Frampton G.K. (2000). Recovery responses of soil surface Collembola after spatial and temporal changes in long-term regimes of pesticide use. *Pedobiologia* 44(3-4): 489-501.
- Frampton G.K. (2002). Long-term impacts of an organophosphate-based regime of pesticides on field and field-edge Collembola communities. *Pest Management Science* 58(10): 991-1001.
- Frampton G.K., Dome J. (2007). The effects on terrestrial invertebrates of reducing pesticide inputs in arable crop edges: a meta-analysis. *Journal of Applied Ecology* 44(2): 362-373.
- Frampton G.K., Langton S.D., Greig-Smith P.W., Hardy A.R. (1992). Changes in the soil fauna at Boxworth. In *Pesticides, cereal farming and the environment. Boxworth project*, (Greig-Smith P. *et al.*, eds.), Ministry of Agriculture, Fisheries and Food: 132-143.
- Frantzen J., Rossi F. (2002). Integration of biological control of common groundsel (*Senecio vulgaris*) and chemical control. *Weed Science* 50: 787-793.
- Fraser P.M., Piercy J.E. (1998). The effects of cereal straw management practices on lumbricid earthworm populations. *Applied Soil Ecology* 9(1-3): 369-373.

- Freemark K., Bert D., Villard M.-A. (2002). Patch-, landscape-, and regional-scale effects on biota. *In* Applying Landscape Ecology in Biological Conservation, (Gutzwiller K.J., ed.), Springer-Verlag, New York: 58-83.
- Freléchoux F., Meisser M., Gillet F. (2007). Succession secondaire et perte de diversité végétale après réduction du broutage dans un pâturage boisé des Alpes centrales suisses. *Botanica Helvetica* 117: 37-56.
- Friberg H., Lagerlof J., Ramert B. (2005). Influence of soil fauna on fungal plant pathogens in agricultural and horticultural systems. *Biocontrol Science and Technology* 15(7): 641-658.
- Frick R., Fluri P. (2001). Bienenverluste beim Mahem mit Rotationsmäherwerken. *Agrarforschung* 8: 196-201.
- Fried G., Chauvel B., Reboud X. (in press). A functional analysis of large scale temporal shifts in the sunflower weed assemblages in France between 1970 and 2000. *Journal of Vegetation Science*.
- Frouz J. (1999). Use of soil dwelling Diptera (Insecta, Diptera) as bioindicators: a review of ecological requirements and response to disturbance. *Agriculture Ecosystems & Environment* 74(1-3): 167-186.
- Funderburk J.E., Rhoads F.M., Teare I.D. (1994). Modifying Soil Nutrient Level Affects Soybean Insect Predators. *Agronomy Journal* 86(3): 581-585.
- Gabriel D., Roschewitz I., Tschamtk T., Thies C. (2006). Beta diversity at different spatial scales: plant communities in organic and conventional agriculture *Ecological Applications* 16(5): 2011-2021.
- Gadoum S., Terzo M., Rasmont P. (2007). Jachères apicoles et jachères fleuries : la biodiversité au menu de quelles abeilles ? *Le Courrier de l'environnement de l'INRA* 54: 57-63.
- Gaisler J., Hejzman M., Pavlu V. (2004). Effect of different mulching and cutting regimes on the vegetation of upland meadow. *Plant, Soil & Environment* 7: 324-331.
- Gallo L., Siverio F., Rodriguez-Perez A.M. (2007). Thermal sensitivity of *Phytophthora cinnamomi* and long-term effectiveness of soil solarisation to control avocado root rot. *Annals of Applied Biology* 150(1): 65-73.
- Garbeva P., van Elsas J.D., van Veen J.A. (2008). Rhizosphere microbial community and its response to plant species and soil history. *Plant and Soil*. DOI 10.1007/s11104-007-9432-0.
- Garbeva P., van Veen J.A., van Elsas J.D. (2004). Assessment of the diversity, and antagonism towards *Rhizoctonia solani* AG3, of *Pseudomonas* species in soil from different agricultural regimes. *FEMS Microbiology Ecology* 47(1): 51-64.
- Garcia F., Carrère P., Soussana J.F., Baumont R. (2003). How do severity and frequency of grazing affect sward characteristics and the choices of sheep during the grazing season? *Grass & Forage Science* 58(2): 138-150.
- Garcin A., Demarle O., Soldati F. (2004). Carabids, indicators of biodiversity and generalist auxiliaries. *Infos-Citifi*(No.199): 42-47.
- Gardiner M.M., Landis D.A. (2007). Impact of intraguild predation by adult *Harmonia axyridis* (Coleoptera: Coccinellidae) on *Aphis glycines* (Hemiptera: Aphididae) biological control in cage studies. *Biological Control* 40(3): 386-395.
- Gardner S.M., Hartley S.E., Davies A., Palmer S.C.F. (1997). Carabid communities on heather moorlands in northeast Scotland: The consequences of grazing pressure for community diversity. *Biological Conservation* 81(3): 275-286.
- Garnier A., Deville A., Lecomte J. (2006). Stochastic modelling of feral plant populations with seed immigration and road verge management. *Ecological Modelling* 197: 373-382.
- Garnier A., Lecomte J. (2006). Using a spatial and stage-structured invasion model to assess the spread of feral populations of transgenic oilseed rape. *Ecological Modelling* 194(1-3): 141-149.
- Garnier E., Aronson J. (1998). Nitrogen use efficiency from leaf to stand level: Clarifying the concept. *In* Inherent Variation in Plant Growth. Physiological Mechanisms and Ecological Consequences, (Lambers H. et al., eds.), Backhuys Publishers, Leiden: 515-538.
- Garnier E., Cortez J., Billès G., Navas M.-L., Roumet C., Debussche M., Laurent G., Blanchard A., Aubry D., Bellmann A., Neill C., Toussaint J.-P. (2004). Plant functional markers capture ecosystem properties during secondary succession. *Ecology* 85(9): 2630-2637.
- Garnier E., Lavorel S., Ansquer P., Castro H., Cruz P., Dolezal J., Eriksson O., Fortunel C., Freitas H., Golodets C., Grigulis K., Jouany C., Kazakou E., Kigel J., Kleyer M., Lehsten V., Lepš J., Meier T., Pakeman R., Papadimitriou M., Papanastasis V.P., Quested H., Quéfier F., Robson M., Roumet C., Rusch G., Skarpe M., Sternberg M., Theau J.-P., Thébault A., Vile D., Zarovali M. (2007). Assessing the effects of land use change on plant traits, communities and ecosystem functioning in grasslands: a standardized methodology and lessons from an application to 11 European sites. *Annals of Botany* 99: 967-985.
- Garola C.V. (1919). La pratique des travaux de la ferme, 3ème ed., Hachette et Cie, Paris, 220 p.
- Garrett K.A., Nelson R.J., Mundt C.C., Chacón G., Jaramillo R.E., Forbes G.A. (2001). The effects of host diversity and other management components on epidemics of potato late blight in the humid highland tropics. *Phytopathology* 91(10): 993-1000.
- Gasquez J. (1994). Les adventices en jachère. *In* Jachères, (Fraval A., ed.), Dossier de l'Environnement de l'INRA, 9, INRA, Paris: 101-104.
- Gauthier D., Gibert P., Hars J. (1992). Sanitary consequences of mountain cattle breeding on wild ungulates. , *Ongulés / Ungulates* 91, Paris-Toulouse, Spitz F. et al., eds., S.F.E.P.M.-I.R.G.M.: 621-630.
- Gayubo S.F., Gonzalez J.A., Asis J.D., Tormos J. (2005). Conservation of European environments: the Spheciformes wasps as biodiversity indicators (Hymenoptera: Apoidea: Ampulicidae, Sphecidae and Crabronidae). *Journal of Natural History* 39(29): 2705-2714.
- Gelsomino A., Cacco G. (2006). Compositional shifts of bacterial groups in a solarized and amended soil as determined by denaturing gradient gel electrophoresis. *Soil Biology & Biochemistry* 38(1): 91-102.
- Gendron A.D., Marcogliese D.J., Barbeau S., Christin M.S., Brousseau P., Ruby S., Cyr P., Fournier M. (2003). Exposure of leopard frogs to a pesticide mixture affects life history characteristics of the lungworm *Rhabdias ranae*. *Oecologia* 135(3): 469-476.
- Genthner F.J., Middaugh D.P. (1992). Effects of *Beauveria bassiana* on embryos on inland silverside fish (*Menidia beryllina*). *Applied Environmental Microbiology* 58(9): 2840-2845.
- Georgieva S.S., McGrath S.P., Hooper D.J., Chambers B.S. (2002). Nematode communities under stress: the long-term effects of heavy metals in soil treated with sewage sludge. *Applied Soil Ecology* 20(1): 27-42.
- Georgis R., Kaya H.K., Gaugler R. (1991). Effect of steinernematid and heterorhabditid nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) on nontarget arthropods. *Environmental Entomology* 20(3): 815-822.

- Ghiorbani R., Seel, W, Litterick, A, Leifert, C, (2000). Evaluation of *Alternaria alternata* for biological control of *Amaranthus retroflexus*. *Weed Science* 48: 474-480.
- Ghosheh H.Z. (2005). Constraints in implementing biological weed control: a review. *Weed Biology and Management* 5: 83-92.
- Giacometti M., Roganti R., De Tann D., Stahlberger-Saitbekova N., Obexer-Ruff G. (2004). Alpine ibex *Capra ibex ibex* x domestic goat *C-aegagrus domestica* hybrids in a restricted area of southern Switzerland. *Wildlife Biology* 10(2): 137-143.
- Gibert P., Appolinaire J. (2004). Intoxication d'isards au lindane dans les Hautes Pyrénées. *Faune Sauvage* 261: 42-47.
- Gibon A. (2005). Managing grassland for production, the environment and the landscape. Challenges at the farm and the landscape level. *Livestock Production Science* 96(1): 11-31.
- Giller P.S. (1996). The diversity of soil communities, the 'poor man's tropical rainforest'. *Biodiversity and Conservation* 5(2): 135-168.
- Girardin P., Bockstaller C., Van der Werf H. (1999). Indicators: Tools to evaluate the environmental impacts of farming systems. *Journal of Sustainable Agriculture* 13(4): 5-21.
- Girvan M.S., Campbell C.D., Killham K., Prosser J.I., Glover L.A. (2005). Bacterial diversity promotes community stability and functional resilience after perturbation. *Environmental Microbiology* 7(3): 301-313.
- Gleeson S.K., Tilman D. (1994). Plant allocation, growth rate and successional status. *Functional Ecology* 8: 543-550.
- Goiti U., Aihartza J.R., Garin I., Zabala J. (2003). Influence of habitat on the foraging behaviour of the Mediterranean horseshoe bat, *Rhinolophus euryale*. *Acta Chiropterologica* 5(1): 75-84.
- Gonnet J.M. (2001). Influence of cattle grazing on population density and species richness of granivorous birds (Emberizidae) in the and plain of the Monte, Argentina. *Journal of Arid Environments* 48(4): 569-579.
- Gonod L.V., Martin-Laurent F., Chenu C. (2006). 2,4-D impact on bacterial communities, and the activity and genetic potential of 2,4-D degrading communities in soil. *Fems Microbiology Ecology* 58(3): 529-537.
- Gonzalez-Zamora J.E., Leira D., Bellido M.J., Avilla C. (2004). Evaluation of the effect of different insecticides on the survival and capacity of *Eretmocerus mundus* Mercet to control *Bemisia tabaci* (Gennadius) populations. *Crop Protection* 23(7): 611-618.
- Goodlass G., Halberg N., Verschuur G. (2003). Input output accounting systems in the European community - an appraisal of their usefulness in raising awareness of environmental problems. *European Journal of Agronomy* 20(1-2): 17-24.
- Gordon C.E. (2000). Fire and cattle grazing on wintering sparrows in Arizona grasslands. *Journal of Range Management* 53(4): 384-389.
- Gordon I.J. (1989). Vegetation community selection by ungulates on the Isle of Rhum. II. Vegetation community selection. *Journal of Applied Ecology* 26(1): 53-64.
- Gordon I.J. (1989). Vegetation community selection by ungulates on the Isle of Rhum. III. Determinants of vegetation community selection. *Journal of Applied Ecology* 26(1): 65-79.
- Gordon P.L., McEwen F.L. (1984). Insecticide-stimulated reproduction of *Myzus persicae*, the green peach aphid (Homoptera: Aphididae). *Canadian Entomologist* 116(5): 783-784.
- Gorman M.L., Reynolds P. (1993). The Impact of Land-Use Change on Voles and Raptors. *Mammal Review* 23(3-4): 121-126.
- Gormsen D., Hedlund K., Korhals G.W., Mortimer S.R., Pizl V., Smilauerova M., Sugg E. (2004). Management of plant communities on set-aside land and its effects on earthworm communities. *European Journal of Soil Biology* 40(3-4): 123-128.
- Gormsen D., Hedlund K., Wang H.F. (2006). Diversity of soil mite communities when managing plant communities on set-aside arable land. *Applied Soil Ecology* 31(1-2): 147-158.
- Gortazar C., Ferroglio E., Hofle U., Frolich K., Vicente J. (2007). Diseases shared between wildlife and livestock: a European perspective. *European Journal of Wildlife Research* 53: 241-256.
- Govaerts B., Mezzalama M., Unno Y., Sayre K.D., Luna-Guido M., Vanherck K., Dendooven L., Deckers J. (2007). Influence of tillage, residue management, and crop rotation on soil microbial biomass and catabolic diversity. *Applied Soil Ecology* 37(1-2): 18-30.
- Grafton-Cardwell E.E., Ouyang Y. (1993). Toxicity of four insecticides to various populations of the predacious mite, *Euseius tularensis* Congdon (Acarina: Phytoseiidae) from San Joaquin Valley California citrus. *Journal of Agricultural Entomology* 10(1): 21-29.
- Grandchamp A.C., Bergamini A., Stofer S., Niemela J., Duelli P., Scheidegger C. (2005). The influence of grassland management on ground beetles (Carabidae, Coleoptera) in Swiss montane meadows. *Agriculture, Ecosystems & Environment* 110(3-4): 307-317.
- Grandval P. (1988). Approche écologique de la gestion de l'espace rural: des besoins de la bécasse (*Scolopax rusticola* L.) à la qualité des milieux. Thèse doctorat : Ecologie, Aménagement Université de Rennes 1, 179 p.
- Gras R., Benoit M., Deffontaines J.P., Duru M., Lafarge M., Langlet A., Osty P.L. (1989). Le fait technique en agronomie. Activite agricole, concepts et methodes d'etude, Inra, L'Harmattan, 184 p.
- Graveland J., van der Wal R., van Balen J.H., van Noordwijk A.J. (1994). Poor reproduction in forest passerines from decline of snail abundance on acidified soils. *Nature* 368: 446-448.
- Grayston S.J., Campbell C.D., Bardgett R.D., Mawdsley J.L., Clegg C.D., Ritz K., Griffiths B.S., Rodwell J.S., Edwards S.J., Davies W.J., Elston D.J., Millard P. (2004). Assessing shifts in microbial community structure across a range of grasslands of differing management intensity using CLPP, PLFA and community DNA techniques. *Applied Soil Ecology* 25(1): 63-84.
- Grayston S.J., Griffith G.S., Mawdsley J.L., Campbell C.D., Bardgett R.D. (2001). Accounting for variability in soil microbial communities of temperate upland grassland ecosystems. *Soil Biology & Biochemistry* 33(4-5): 533-551.
- Gregory R.D., van Strien A., Vorisek P., Meyling A.W.G., Noble D.G., Foppen R.P.B., Gibbons D.W. (2005). Developing indicators for European birds. *Philosophical transactions of the Royal Society B* 360(1454): 269-288.
- Grémillet X., Boireau J. (2004). Intoxication mortelle par le plomb et par les fongicides P.C.P. des juvéniles dans un gîte de parturition de Grands Rhinolophes, *Rhinolophus ferrumequinum*, dans le Finistère : difficultés du diagnostic et réalisation et d'un gîte alternatif. *Symbioses* 10: 59-61.
- Grevilliot F., Krebs L., Muller S. (1998). Comparative importance and interference of hydrological conditions and soil nutrient gradients in floristic biodiversity in flood meadows. *Biodiversity and Conservation* 7(11): 1495-1520.
- Grieve I.C., Davidson D.A., Bruneau P.M.C. (2005). Effects of liming on void space and aggregation in an upland grassland soil. *Geoderma* 125(1-2): 39-48.
- Griffiths B.S., Caul S., Thompson J., Birch A.N.E., Cortet J., Andersen M.N., Krogh P.H. (2007). Microbial and microfaunal community structure in cropping systems with genetically modified plants. *Pedobiologia* 51(3): 195-206.

- Griffiths B.S., Caul S., Thompson J., Birch A.N.E., Scrimgeour C., Andersen M.N., Cortet J., Messean A., Sausse C., Lacroix B., Krogh P.H. (2005). A comparison of soil microbial community structure, protozoa and nematodes in field plots of conventional and genetically modified maize expressing the *Bacillus thuringiensis* CryIAb toxin. *Plant and Soil* 275(1-2): 135-146.
- Griffiths B.S., Caul S., Thompson J., Birch A.N.E., Scrimgeour C., Cortet J., Foggo A., Hackett C.A., Krogh P.H. (2006). Soil microbial and faunal community responses to Bt maize and insecticide in two soils. *Journal of Environmental Quality* 35(3): 734-741.
- Grime J.P. (1973). Competitive exclusion in herbaceous vegetation. *Nature* 242: 344-347.
- Grime J.P. (1979). *Plant strategies and vegetation processes*, John Wiley & Sons, Chichester, 222 p.
- Grime J.P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology* 86: 902-906.
- Grime J.P. (2001). *Plant strategies, vegetation processes, and ecosystem properties*, 2nd Edition ed., John Wiley & Sons, Chichester, 456 p.
- Grime J.P. (2003). Plants hold the key. Ecosystems in a changing world. *Biologist* 50(2): 87-91.
- Grime J.P. (2006). Trait convergence and trait divergence in herbaceous plant communities: Mechanisms and consequences. *Journal of Vegetation Science* 17: 255-260.
- Groot A.T., Dicke M. (2002). Insect-resistant transgenic plants in a multi-trophic context. *Plant Journal* 31(4): 387-406.
- Gross K.L., Mittelbach G.G., Reynolds H.L. (2005). Grassland invasibility and diversity: responses to nutrients, seed input, and disturbance. *Ecology* 86: 476-486.
- Grossi J.L., Chenavier L., Delcros P., Brun J.J. (1995). Effects of Landscape Structure on Vegetation and Some Animal Groups after Agriculture Abandonment. *Landscape and Urban Planning* 31(1-3): 291-301.
- Grynja M., Kryszak A., Grzelak M. (1996). Species diversity in the meadow communities of the middle Odra river. *Prace z Zakresu Nauk Rolniczych* 81: 81-86.
- Gudleifsson B.E. (2005). Beetle species (Coleoptera) in hayfields and pastures in northern Iceland. *Agriculture, Ecosystems & Environment* 109(3-4): 181-186.
- Guggenberger G., Frey S.D., Six J., Paustian K., Elliott E.T. (1999). Bacterial and fungal cell-wall residues in conventional and no-tillage agroecosystems. *Soil Science Society of America Journal* 63(5): 1188-1198.
- Gurr G.M., Thwaite W.G., Nicol H.I. (1999). Field evaluation of the effects of the insect growth regulator tebufenozide on entomophagous arthropods and pests of apples. *Australian Journal of Entomology* 38(2): 135-140.
- Gustavsson E., Lennartsson T., Emanuelsson M. (2007). Land use more than 200 years ago explains current grassland plant diversity in a Swedish agricultural landscape. *Biological Conservation* 138(1-2): 47-59.
- Haberl H., Plutzer C., Erb K.H., Gaube V., Pollheimer M., Schulz N.B. (2005). Human appropriation of net primary production as determinant of avifauna diversity in Austria. *Agriculture, Ecosystems & Environment* 110(3/4): 119-131.
- Hager H.A., Gorman R.M., Nudds T.D. (2006). The relative performance of umbrella species for biodiversity conservation in island archipelagos of the Great Lakes, North America. *Ecoscience* 13(4): 475-484.
- Hagn A., Engel M., Kleikamp B., Munch J., Schloter M., Bruns C. (2007). Microbial community shifts in *Pythium ultimum*-inoculated suppressive substrates. *Biology and Fertility of Soils* 44(3): 481-490.
- Hagn A., Pritsch K., Schloter M., Munch J.C. (2003). Fungal diversity in agricultural soil under different farming management systems, with special reference to biocontrol strains of *Trichoderma* spp. *Biology and Fertility of Soils* 38(4): 236-244.
- Hails R.S. (2000). Genetically modified plants - the debate continues. *Trends in Ecology & Evolution* 15(1): 14-18.
- Hairton N.G., Hairton N.G. (1993). Cause-Effect Relationships in Energy-Flow, Trophic Structure, and Interspecific Interactions. *American Naturalist* 142(3): 379-411.
- Halffter G. (1998). A strategy for measuring landscape biodiversity. *Biology International*(No. 36): 3-17.
- Häni F.J., Boller E.F., Keller S. (1998). Natural regulation at the farm level. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Pickett C.H., Bugg R.L., eds.), University of California Press, Berkeley - Los Angeles - London: 161-210.
- Hansen S. (1996). Effects of manure treatment and soil compaction on plant production of a dairy farm system converting to organic farming practice. *Agriculture, Ecosystems & Environment* 56(3): 173-186.
- Hansson M., Fogelfors H. (2000). Management of a semi-natural grassland; results from a 15-year-old experiment in southern Sweden. *Journal of Vegetation Science* 11(1): 31-38.
- Harper J.L., Hawksworth D.L. (1994). Biodiversity: measurement and estimation. *Philosophical Transactions of the Royal Society of London Series B* 345(1311): 5-12.
- Hars J., Boué F., Boireau J., Garin-Bastuji B., Le Potier M.F., Mesplede A., Rossi S., Saint-Andrieux C., Toma B. (2004). Impact sanitaire de l'augmentation des effectifs de sangliers sauvages (*Sus scrofa*) en France / Sanitary impact of wild-boar (*Sus scrofa*) population number increase in France. *Game & Wildlife Science* 21(4): 513-528.
- Hart J.D., Milsom T.P., Baxter A., Kelly P.F., Parkin W.K. (2002). The impact of livestock on Lapwing *Vanellus vanellus* breeding densities and performance on coastal grazing marsh. *Bird Study* 49: 67-78.
- Hartley S.E., Gardner S.M., Mitchell R.J. (2003). Indirect effects of grazing and nutrient addition on the hemipteran community of heather moorlands. *Journal of Applied Ecology* 40(5): 793-803.
- Hartnett D., Hickman K., Walter L. (1996). Effects of bison grazing, fire, and topography on floristic diversity in tallgrass prairie. *Journal of Range Management* 49(5): 413-420.
- Haskell D.G. (2000). Effects of Forest Roads on Macroinvertebrate Soil Fauna of the Southern Appalachian Mountains. *Conservation Biology* 14(1): 57-63.
- Hasken K.H., Poehling H.M. (1995). Effects of different intensities of fertilisers and pesticides on aphids and aphid predators in winter wheat. *Agriculture, Ecosystems & Environment* 52(1): 45-50.
- Hattingh V., Tate B. (1995). Effects of field-weathered residues of insect growth regulators on some Coccinellidae (Coleoptera) of economic importance as biocontrol agents. *Bulletin of Entomological Research* 85(4): 489-493.
- Havet P. (2002). Agriculture et biodiversité. Du concept scientifique à la pratique. *Faune Sauvage* 256: 25-32.

- Hayes T.B., Paola Case P., Chui S., Chung D., Haeffele C., Haston K., Lee M., Phoung Mai V., Marjuoa Y., Parker J., Tsui M. (2006). Pesticide Mixtures, Endocrine Disruption, and Amphibian Declines: Are We Underestimating the Impact? *Environmental Health Perspectives* 114(S1): 40-50.
- Heard M.S., Clark S.J., Rothery P., Perry J.N., Bohan D.A., Brooks D.R., Champion G.T., Dewar A.M., Hawes C., Haughton A.J., May M.J., Scott R.J., Stuart R.S., Squire G.R., Firbank L.G. (2003). Weeds in fields with contrasting conventional and genetically modified herbicide-tolerant crops. I. Effects on abundance and diversity. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 358: 1819-1832.
- Heard M.S., Clark S.J., Rothery P., Perry J.N., Bohan D.A., Brooks D.R., Champion G.T., Dewar A.M., Hawes C., Haughton A.J., May M.J., Scott R.J., Stuart R.S., Squire G.R., Firbank L.G. (2006). Effects of successive seasons of genetically modified herbicide-tolerant maize cropping on weeds and invertebrates. *Annals of Applied Biology* 149: 249-254.
- Heard M.S., Hawes C., Champion G.T., Clark S.J., Firbank L.G., Haughton A.J., Parish A.M., Perry J.N., Rothery P., Roy D.B., Scott R.J., Skellern M.P., Squire G.R., Hill M.O. (2003). Weeds in fields with contrasting conventional and genetically modified herbicide-tolerant crops. II. Effects on individual species. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 358: 1833-1846.
- Heard M.S., Rothery P., Perry J.N., Firbank L.G. (2005). Predicting longer-term changes in weed populations under GMHT crop management. *Weed Research* 45(5): 331-338.
- Heckmann L.H., Griffiths B.S., Caul S., Thompson J., Pusztai-Carey M., Moar W.J., Andersen M.N., Krogh P.H. (2006). Consequences for *Protaphorura armata* (Collembola : Onychiuridae) following exposure to genetically modified *Bacillus thuringiensis* (Bt) maize and non-Bt maize. *Environmental Pollution* 142(2): 212-216.
- Hector A., Bagchi R. (2007). Biodiversity and ecosystem multifunctionality. *Nature* 448: 188-191.
- Hedde M. (2006). Etude de la relation entre la biodiversité des macro-invertébrés et la dynamique de la matière organique des sols limoneux de Haute-Normandie. Thèse de doctorat, Université de Rouen.
- Hedlund K. (2002). Soil microbial community structure in relation to vegetation management on former agricultural land. *Soil Biology & Biochemistry* 34(9): 1299-1307.
- Hedlund K., Griffiths B., Christensen S., Scheu S., Setälä H., Tschamtké T., Verhoef H. (2004). Trophic interactions in changing landscapes: responses of soil food webs. *Basic and Applied Ecology* 5(6): 495-503.
- Hedlund K., Regina I.S., Van der Putten W.H., Leps J., Diaz T., Korthals G.W., Lavorel S., Brown V.K., Gormsen D., Mortimer S.R., Barrueco C.R., Roy J., Smilauer P., Smilauerova M., Van Dijk C. (2003). Plant species diversity, plant biomass and responses of the soil community on abandoned land across Europe: idiosyncrasy or above-belowground time lags. *Oikos* 103(1): 45-58.
- Hejman M., Klaudivsova M., Sturba J., Pavlu V., Schellberg J., Hejmanova P., Hakl J., Rauch O., Vacek S. (2007). Revisiting a 37 years abandoned fertilizer experiment on *Nardus* grassland in the Czech Republic *Agriculture, Ecosystems & Environment* 118(1-4): 231-236
- Hellmich R.L., Siegfried B.D., Sears M.K., Stanley-Horn D.E., Daniels M.J., Mattila H.R., Spencer T., Bidne K.G., Lewis L.C. (2001). Monarch larvae sensitivity to *Bacillus thuringiensis*-purified proteins and pollen, *Proc. Natl Acad. Sci. USA*, 98: 11925-11930.
- Henault C., English L.C., Halpin C., Andreux F., Hopkins D.W. (2006). Microbial community structure in soils with decomposing residues from plants with genetic modifications to lignin biosynthesis. *Fems Microbiology Letters* 263(1): 68-75.
- Hendrickx F., Maelfait J.P., Van Wingerden W., Schweiger O., Speelmans M., Aviron S., Augenstein I., Billeter R., Bailey D., Bukacek R., Burel F., Diekötter T., Dirksen J., Herzog F., Liira J., Roubalova M., Vandomme V., Bugter R. (2007). How landscape structure, land-use intensity and habitat diversity affect components of total arthropod diversity in agricultural landscapes. *Journal of Applied Ecology* 44(2): 340-351.
- Hernandez L.M., Ibanez C., Fernandez M.A., Guillen A., Gonzalez M.J., Perez J.L. (1993). Organochlorine Insecticide and Pcb Residues in 2 Bat Species from 4 Localities in Spain. *Bulletin of Environmental Contamination and Toxicology* 50(6): 871-877.
- Heroldova M., Bryja J., Zejda J., Tkadlec E. (2007). Structure and diversity of small mammal communities in agriculture landscape. *Agriculture, Ecosystems & Environment* 120(2-4): 206-210.
- Herrmann S., Dabbert S., Schwarz-von Raumer H.G. (2003). Threshold values for nature protection areas as indicators for bio-diversity - a regional evaluation of economic and ecological consequences. *Agriculture, Ecosystems & Environment* 98(1/3): 493-506.
- Herzog F., Richner W., Walter T. (2006). Mesures écologiques: un effet modérément positif. *Revue Suisse d'Agriculture* 38(2): 63-68.
- Herzog F., Steiner B., Bailey D., Baudry J., Billeter R., Bukacek R., de Blust G., de Cock R., Dirksen J., Dormann C.F., de Filippi R., Frossard E., Liira J., Schmidt T., Stockli R., Thenail C., van Wingerden W., Bugter R. (2006). Assessing the intensity of temperate European agriculture at the landscape scale. *European Journal of Agronomy* 24(2): 165-181.
- Herzon I., O'Hara R.B. (2007). Effects of landscape complexity on farmland birds in the Baltic States. *Agriculture, Ecosystems & Environment* 118(1-4): 297-306.
- Hester A.J., Bergman M., Iason G.R., Moen J. (2006). Impact of large herbivores on plant community structure and dynamics. In *Large Herbivore Ecology, Ecosystem Dynamics and Conservation* (Danell K. et al., eds.), Cambridge University Press: 97-141.
- Hewison A.J.M., Vincent J.P., Joachim J., Angibault J.M., Cargnelutti B., Cibien C. (2001). The effects of woodland fragmentation and human activity on roe deer distribution in agricultural landscapes. *Canadian Journal of Zoology* 79(4): 679-689.
- Hiddink G.A., van Bruggen A.H.C., Termorshuizen A.J., Raaijmakers J.M., Semenov A.V. (2005). Effect of organic management of soils on suppressiveness to *Gaeumannomyces graminis* var. *tritici* and its antagonist, *Pseudomonas fluorescens*. *European Journal of Plant Pathology* 113(4): 417-435.
- Hietala-Koivu R., Jarvenpää T., Helenius J. (2004). Value of semi-natural areas as biodiversity indicators in agricultural landscapes. *Agriculture, Ecosystems & Environment* 101(1): 9-19.
- Hilbeck A. (2001). Implications of transgenic, insecticidal plants for insect and plant biodiversity. *Perspectives in Plant Ecology Evolution and Systematics* 4(1): 43-61.
- Hilbeck A., Moar W.J., Pusztai-Carey M., Filippini A., Bigler F. (1999). Prey-mediated effects of Cry1Ab toxin and protoxin and Cry2A protoxin on the predator *Chrysoperla carnea*. *Entomologia Experimentalis et Applicata* 91: 305-316.
- Hirano S.S., Upper C.D. (2000). Bacteria in the leaf ecosystem with emphasis on *Pseudomonas syringae* - a pathogen, ice nucleus, and epiphyte. *Microbiology and Molecular Biology Reviews* 64(3): 624-653.

- Hobbs R.J., McIntyre S. (2005). Categorizing Australian landscapes as an aid to assessing the generality of landscape management guidelines. *Global Ecology and Biogeography* 14(1): 1-15.
- Hodde M.S. (2004). Restoring balance: Using exotic species to control invasive exotic species. *Conservation Biology* 18(1): 38-49.
- Hodgson J.G., Montserrat-Martí G., Tallowin J., Thompson K., Diaz S., Cabido M., Grime J.P., Wilson P.J., Band S.R., Bogard A., Cabido R., Caceres D., Castro-Díez P., Ferrer C., Maestro-Martínez M., Perez-Rontome M.C., Charles M., Cornelissen J.H.C., Dabbert S., Perez-Harguindeguy N., Krimly T., Sijsma F.J., Strijker D., Vendramini F., Guerrero-Campo J., Hynd A., Jones G., Romo-Díez A., de Torres Espuny L., Villar-Salvador P., Zak M.R. (2005). How much will it cost to save grassland diversity? *Biological Conservation* 122(2): 263-273.
- Hofer G., Herzog F., Spiess M., Birrer S. (2002). Ecological measures: vegetation and breeding birds. *Agrarforschung* 9(4): 152-157.
- Hoffmann J., Greef J.M. (2003). Mosaic indicators--theoretical approach for the development of indicators for species diversity in agricultural landscapes. *Agriculture, Ecosystems & Environment* 98(1-3): 387-394.
- Hokkanen H., Husberg G.B., Soderblom M. (1988). Natural enemy conservation for the integrated control of the rape blossom beetle *Meligethes aeneus* F. *Annales Agriculturae Fenniae* 27(4): 281-194.
- Hole D.G., Perkins A.J., Wilson J.D., Alexander I.H., Grice F., Evans A.D. (2005). Does organic farming benefit biodiversity? *Biological Conservation* 122(1): 113-130.
- Holland J.M., Drysdale A., Hewitt M.V., Turley D. (1996). The LINK IFS project - the effect of crop rotations and cropping systems on Carabidae. *Aspects of Applied Biology* 47: 119-126.
- Holland J.M., Hutchison M.A.S., Smith B., Aebischer N.J. (2006). A review of invertebrates and seed-bearing plants as food for farmland birds in Europe. *Annals of Applied Biology* 148: 49-71.
- Holzschuh A., Steffan-Dewenter I., Kleijn D., Tscharntke T. (2007). Diversity of flower-visiting bees in cereal fields: effects of farming system, landscape composition and regional context. *Journal of Applied Ecology* 44(1): 44-49.
- Hondru N., Margarit G., Picu I., Chireceanu C. (1998). Effect of crop rotation and fertilization on the invertebrate complex from the soil of an irrigated maize crop. *Romanian Agricultural Research*(9/10): 97-105.
- Honnay O., Adriaens D., Coart E., Jacquemyn H., Roldan-Ruiz I. (2007). Genetic diversity within and between remnant populations of the endangered calcareous grassland plant *Globularia bisnagarica* L. *Conservation Genetics* 8(2): 293-303.
- Hopkins A., Wilkins R.J. (2006). Temperate grassland: key developments in the last century and future perspectives. *Journal of Agricultural Science* 144: 503-523.
- Houssard C., Escarré J., Romane F. (1980). Development of species diversity in some mediterranean plant communities. *Vegetatio* 43: 59-72.
- Hovd H., Skogen A. (2005). Plant species in arable field margins and road verges of central Norway. *Agriculture, Ecosystems & Environment* 110(3-4): 257-265.
- Howarth F.G. (2000). Non-target effects of biological control agents. In *Biological Control: Measures of success*, (Gurr G.M., Wratten S.D., eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands: 369-403.
- Hoy M.A., Groot J.J.R., van de Baan H.E. (1985). Influence of aerial dispersal on persistence and spread of pesticide-resistant *Metaseiulus occidentalis* in California almond orchards. *Entomologia Experimentalis et Applicata* 37(1): 17-31.
- Hubbell S.P. (2001). A unified neutral theory of biodiversity and biogeography, *Monographs in population biology*, 32, Princeton University Press, 396 p.
- Huelsbergen K.-J., Kuestermann B. (2005). Development of an environmental management system for organic farms and its introduction into practice, *Researching Sustainable Systems - International Scientific Conference on Organic Agriculture*, Adelaide, Australia, 2005/09/21-23, 4375. Organic eprints.
- Hülsbergen K.J. (2003). *Entwicklung und Anwendung eines Bilanzierungsmodells zur Bewertung der Nachhaltigkeit andwirtschaftlicher Systeme*, Shaker Verlag, 292 p.
- Hummel R.L., Walgenbach J.F., Hoyt G.D., Kennedy G.G. (2002). Effects of production system on vegetable arthropods and their natural enemies. *Agriculture Ecosystems & Environment* 93(1-3): 165-176.
- Hunter Jr M.L. (1996). *Fundamentals of conservation biology*, Blackwell Publishing, Cambridge, 547 p.
- Huston M.A. (1979). A general hypothesis of species diversity. *American Naturalist* 113: 81-101.
- Huston M.A. (1994). *Biological Diversity. The Coexistence of Species in changing landscapes*, Cambridge University Press, 681 p.
- Huston M.A. (1999). Local processes and regional patterns: appropriate scales for understanding variation in the diversity of plants and animals. *Oikos* 86: 393-401.
- Huston M.A., Smith T. (1987). Plant succession: Life history and competition. *American Naturalist* 130: 168-198.
- Hutchings M.R., Harris S. (1996). The current status of the brown hare (*Lepus europaeus*) in Britain, Joint Nature Conservation Committee. <http://www.jncc.gov.uk/page-2816>
- Hutton S.A., Giller P.S. (2003). The effects of the intensification of agriculture on northern temperate dung beetle communities. *Journal of Applied Ecology* 40(6): 994-1007.
- Hyvönen T. (2007). Can conversion to organic farming restore the species composition of arable weed communities? *Biological Conservation* 137(3): 382-390.
- Hyvönen T., Ketoja E., Salonen J., Jalli H., Tiainen J. (2003). Weed species diversity and community composition in organic and conventional cropping of spring cereals. *Agriculture, Ecosystems & Environment* 97: 131-149.
- Ibekwe A.M. (2004). Effects of fumigants on non-target organisms in soils. In *Advances in Agronomy*, Vol 83, Advances in Agronomy, 83, ELSEVIER ACADEMIC PRESS INC, San Diego: 1-35.
- Ibrahim M.A., Kainulainen P., Aflatuni A., Tiilikka K., Holopainen J.K. (2001). Insecticidal, repellent, antimicrobial activity and phytotoxicity of essential oils: with special reference to limonene and its suitability for control of insect pests. *Agricultural and Food Science in Finland* 10(3): 243-259.
- IFEN (2006). L'agriculture et la sylviculture, Les synthèses, IFEN: 63-82. http://www.ifen.fr/uploads/media/agriculture_ree2006_01.pdf
- Inouye R.S., Huntly N.J., Tilman D., Tester J.R., Stillwell M., Zinnel K.C. (1987). Old-field succession on a Minnesota sand plain. *Ecology* 68: 12-26.

- Iqbal J., Wright D. (1997). Effects of nitrogen supply on competition between wheat and three annual weed species. *Weed Research*, 37(6): 391-400.
- Ismail S.M.M. (1997). Selectivity and joint action of *Melia azedarach* L. fruit extracts with certain acaricides to *Tetranychus urticae* Koch and *Stethorus gilvifrons* Mulsant. *Annals of Agricultural Science, Moshtohor* 35(1): 605-618.
- Isselstein J., Jeangros B., Pavlu V. (2005). Agronomic aspects of biodiversity targeted management of temperate grasslands in Europe – a review. *Agronomy Research* 3(2): 139-151.
- Jansen J.P. (1999). Selectivity of plant protection products used in wheat in summer on the major cereal aphid natural enemies. *Mededelingen - Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, Universiteit Gent* 64(3a): 25-40.
- Janson J., P., de Fontanges A., Deneufbourg F., Jouy L. (2004). Nuisibilité des adventices en production de semences fourragères : cas du vulpin dans le ray-grass anglais, *XIIème Colloque International sur la Biologie des Mauvaises Herbes*, Dijon (France): 81-88.
- Janssens F., Peeters A., Tallowin J.R.B., Bakker J.P., Bekker R.M., Fillat F., Oomes M.J.M. (1998). Relationship between soil chemical factors and grassland diversity. *Plant and Soil* 202(1): 69-78.
- Jauzein P. (1996). Les souchets tubéreux. «Comestibles» ou «ronds» il faut apprendre à les connaître., *Phytoma, la défense des végétaux* 484: 27-21.
- Jauzein P. (2001). Biodiversité des champs cultivés : l'enrichissement floristique. In *Dossiers de l'Environnement de l'INRA - Agriculture et biodiversité des plantes*, (Le Perchec S. et al., eds.), 21: 43-64.
- Jauzein P. (2001). L'appauvrissement floristiques des champs cultivés. In *Dossiers de l'Environnement de l'INRA - Agriculture et biodiversité des plantes*, (Le Perchec S. et al., eds.), 21: 65-78.
- Jauzein P., Montégut J. (1983). Graminées (Poaceae) nuisibles en agriculture, Ecole Nationale Supérieure d'Horticulture de Versailles, Aubervilliers (France), 537 p.
- Jeangros B., Bertola C. (2002). Long-term evolution of an intensively managed meadow after cessation of fertilisation and reduction of cutting frequency. In *Multi-function grasslands: quality forages, animal products and landscapes*. Proceedings of the 19th General Meeting of the European Grassland Federation, La Rochelle, France, 27-30 May 2002, Organizing Committee of the European Grassland Federation, Versailles cedex France: 794-795.
- Jeanneret P., Schupbach B., Steiger J., Waldburger M., Bigler F. (2000). Evaluation of ecological measures: biodiversity. Spiders and butterflies. *Agrarforschung* 7(3): 112-116.
- Jefferson R.G. (2005). The conservation management of upland hay meadows in Britain: a review. *Grass and Forage Science* 60(4): 322-331.
- Jelenic S. (2003). Controversy associated with the common component of most transgenic plants - Kanamycin resistance marker gene. *Food Technology and Biotechnology* 41(2): 183-190.
- Jelinski D.E. (1997). On genes and geography: a landscape perspective on genetic variation in natural plant populations *Landscape & Urban Planning* 39(1): 11-23.
- Jenser G. (2005). The judgement of the population regulating effect of *Zetzellia mali* (Stigmaeidae). *Bulletin OILB/SROP* 28(7): 239-244.
- Jervis M.A. (1997). Parasitoids as limiting and selective factors: can biological be evolutionarily stable? *Trends in Ecology & Evolution* 12(10): 378-380.
- Johansen A., Knudsen I.M.B., Binnerup S.J., Winding A., Johansen J.E., Jensen L.E., Andersen K.S., Svenning M.M., Bonde T.A. (2005). Non-target effects of the microbial control agents *Pseudomonas fluorescens* DR54 and *Clonostachys rosea* IK726 in soils cropped with barley followed by sugar beet: a greenhouse assessment. *Soil Biology & Biochemistry* 37(12): 2225-2239.
- Johnson M.W., Krugner R. (2004). Impact of legally compliant organic pesticides on natural enemies. In *California Conference on Biological Control IV*, Berkeley, California, USA, 13-15 July, 2004, (Hodde M.S., ed.), Berkeley USA: 69-75.
- Jollands N. (2006). How to aggregate sustainable development indicators: a proposed framework and its application. *International Journal of Agricultural Resources, Governance and Ecology* 5(1): 18-34.
- Jones C.G., Lawton J.H., Shachak M. (1997). Positive and negative effects of organisms as physical ecosystem engineers. *Ecology* 78(7): 1946-1957.
- Jones N.E., Naylor R.E.L. (1992). Significance of seed rain from set-aside. *BCPC Monograph* 50: 91-96.
- Joseph C.A., Phillips D.A. (2003). Metabolites from soil bacteria affect plant water relations. *Plant Physiology and Biochemistry* 41(2): 189-192.
- Joshi J., Stoll P., Rusterholz H.-P., Schmid B., Dolt C., Baur B. (2006). Small-scale experimental habitat fragmentation reduces colonization rates in species-rich grasslands. *Oecologia* 148(1): 144-152.
- Jourdain E., Gibert P., Gauthier D., Fromont E., Jullien J.M., Hars J. (2005). Sondage sur les maladies abortives chez les ongulés sauvages et domestiques en alpage. *Faune Sauvage* 268: 24-32.
- Joyce C. (2001). The sensitivity of a species-rich flood-meadow plant community to fertilizer nitrogen: the Luznice river floodplain, Czech Republic. *Plant Ecology* 155(1): 47-60.
- Jules E.S., Shahani P. (2003). A broader ecological context to habitat fragmentation: why matrix habitat is more important than we thought. *Journal of Vegetation Science* 14(3): 459-464.
- Juttersonke B., Arit K. (2002). Influence of extensification in crop husbandry on the diversity of plant species in fields and field margins. *Zeitschrift Fur Pflanzenkrankheiten Und Pflanzenschutz-Journal of Plant Diseases and Protection*. 305-309.
- Kaaya G.P. (1992). Non-chemical agents and factors capable of regulating tick populations in nature: a mini review. *Insect Science and its Application* 13(4): 587-594.
- Kajita Y., Takano F., Yasuda H., Agarwala B.K. (2000). Effects of indigenous ladybird species (Coleoptera: Coccinellidae) on the survival of an exotic species in relation to prey abundance. *Applied Entomology and Zoology* 35(4): 473-479.
- Katan J. (1996). Soil solarization : integrated control aspects. In *Principles and Practice of Managing Soilborne Plant Pathogens*, (Hall R., ed.), APS Press, St. Paul, Minnesota: 250-278.
- Kautz T., Lopez-Fando C., Ellmer F. (2006). Abundance and biodiversity of soil microarthropods as influenced by different types of organic manure in a long-term field experiment in Central Spain. *Applied Soil Ecology* 33(3): 278-285.
- Keddy P. (1992). Assembly and response rules: two goals for predictive community ecology. *Journal of Vegetation Science* 3: 157-164.
- Keddy P.A. (1989). *Competition*, Chapman and Hall, London, 202 p.

- Keith D.A., Burgman M.A. (2004). The Lazarus effect: can the dynamics of extinct species lists tell us anything about the status of biodiversity? . *Biological Conservation* 117(1): 41-48.
- Kendall D.A., Chinn N.E., Smith B.D., Tidboald C., Winstone L., Western N.M. (1991). Effects of straw disposal and tillage on spread of barley yellow dwarf virus in winter barley. *Annals of Applied Biology* 119(2): 359-364.
- Kerr J.T., Sugar A., Packer L. (2000). Indicator taxa, rapid biodiversity assessment, and nestedness in an endangered ecosystem. *Conservation Biology* 14(6): 1726-1734.
- Ketoh G.K., Glietho A.I., Huignard J. (2002). Susceptibility of the bruchid *Callosobruchus maculatus* (Coleoptera: Bruchidae) and its parasitoid *Dinarmus basalis* (Hymenoptera: Pteromalidae) to three essential oils. *Journal of Economic Entomology* 95(1): 174-182.
- Kidd P., Prieto-Fernandez A., Monterroso C., Acea M. (2007). Rhizosphere microbial community and hexachlorocyclohexane degradative potential in contrasting plant species. *Plant and Soil* 302(1-2): 233-247.
- Kie J.G., Bowyer R.T., Nicholson M.C., Boroski B.B., Loft E.R. (2002). Landscape heterogeneity at differing scales: Effects on spatial distribution of mule deer. *Ecology* 83(2): 530-544.
- Kiehl K., Eischeid I., Gettner S., Walter J. (1996). Impact of different sheep grazing intensities on salt marsh vegetation in northern Germany. *Journal of Vegetation Science* 7(1): 99-106.
- King K.L., Hutchinson K.J. (2007). Pasture and grazing land: assessment of sustainability using invertebrate bioindicators. *Australian Journal of Experimental Agriculture* 47(4): 392-403.
- Kinkel L.L., Wilson M., Lindow S.E. (2000). Plant species and plant incubation conditions influence variability in epiphytic bacterial population size. *Microbial Ecology* 39(1): 1-11.
- Kirkegaard J.A., Sarwar M. (1998). Biofumigation potential of brassicas - I. Variation in glucosinolate profiles of diverse field-grown brassicas. *Plant and Soil* 201(1): 71-89.
- Kishimoto H. (2002). Species composition and seasonal occurrence of spider mites (Acari : Tetranychidae) and their predators in Japanese pear orchards with different agrochemical spraying programs. *Applied Entomology and Zoology* 37(4): 603-615.
- Kladivko E.J. (2001). Tillage systems and soil ecology. *Soil & Tillage Research* 61(1-2): 61-76.
- Klamer M., Hedlund K. (2004). Fungal diversity in set-aside agricultural soil investigated using terminal-restriction fragment length polymorphism. *Soil Biology & Biochemistry* 36(6): 983-988.
- Klein E.K., Lavigne C., Devaux C., Gouyon P.H. (2003). Modélisation de la dispersion à longue distance, *Impact sur l'environnement des cultures de colza génétiquement modifié tolérant à un herbicide*, Novembre 2003, 43, Commission du Génie Biomoléculaire: 12-19.
- Klimek S., Richter gen. Kemmermann A., Hofmann M., Isselstein J. (2007). Plant species richness and composition in managed grasslands: The relative importance of field management and environmental factors. *Biological Conservation* 134(4): 559-570.
- Klimes F. (2000). Dynamics of species richness of floodplain meadows. *Roslinna Vyroba* 46(5): 199-208.
- Klopper J.W., Rodriguez-Kabana R., Zehnder G.W., Murphy J.F., Sikora E., Fernandez C. (1999). Plant root-bacterial interactions in biological control of soilborne diseases and potential extension to systemic and foliar diseases. *Australasian Plant Pathology* 28(1): 21-26.
- Klose S., Acosta-Martinez V., Ajwa H.A. (2006). Microbial community composition and enzyme activities in a sandy loam soil after fumigation with methyl bromide or alternative biocides. *Soil Biology & Biochemistry* 38(6): 1243-1254.
- Koch B. (1996). Extensification of intensively managed grassland. *Agrarforschung* 3(4): 149-152.
- Koehler H.H. (1997). Mesostigmata (Gamasina, Uropodina), efficient predators in agroecosystems. *Agriculture, Ecosystems & Environment* 62(2): 105-117.
- Koehler H.H. (1999). Predatory mites (Gamasina, Mesostigmata). *Agriculture, Ecosystems & Environment* 74(1-3): 395-410.
- Kolliker R., Stadelmann F.J., Reidy B., Nosberger J. (1998). Fertilization and defoliation frequency affect genetic diversity of *Festuca pratensis* Huds. in permanent grasslands. *Molecular Ecology* 7(11): 1557-1567.
- Kong W.D., Zhu Y.G., Fu B.J., Marschner P., He J.Z. (2006). The veterinary antibiotic oxytetracycline and Cu influence functional diversity of the soil microbial community. *Environmental Pollution* 143(1): 129-137.
- Korthals G.W., Smilauer P., van Dijk C., van der Putten W.H. (2001). Linking above- and below-ground biodiversity: abundance and trophic complexity in soil as a response to experimental plant communities on abandoned arable land. *Functional Ecology* 15(4): 506-514.
- Koss A.M., Jensen A.S., Schreiber A., Pike K.S., Snyder W.E. (2005). Comparison of predator and pest communities in Washington potato fields treated with broad-spectrum, selective, or organic insecticides. *Environmental Entomology* 34(1): 87-95.
- Kowalchuk G.A., Bruinsma M., van Veen J.A. (2003). Assessing responses of soil microorganisms to GM plants. *Trends in Ecology & Evolution* 18(8): 403-410.
- Kozár F. (1992). Organization of arthropod communities in agroecosystems, *Acta phytopathol. Entomol. Hung.*, 27: 365-373.
- Kozdroj J., Trevors J.T., van Elsland J.D. (2004). Influence of introduced potential biocontrol agents on maize seedling growth and bacterial community structure in the rhizosphere. *Soil Biology & Biochemistry* 36(11): 1775-1784.
- Kramarz P., Stark J.D. (2003). Population level effects of cadmium and the insecticide imidacloprid to the parasitoid, *Aphidius ervi* after exposure through its host, the pea aphid, *Acyrtosiphon pisum* (Harris). *Biological Control* 27(3): 310-314.
- Kramer I. (1996). Biodiversity of arthropods in wandering fallows and its evaluation on the basis of ground beetles, hoverflies and aculeate Hymenoptera. In "Agrarökologie", pp. 149 pp.
- Krauss J., Klein A.-M., Steffan-Dewenter I., Tschantke T. (2004). Effects of habitat area, isolation, and landscape diversity on plant species richness of calcareous grasslands. *Biodiversity and Conservation* 13(8): 1427-1439.
- Kreiter S., Brian F. (1987). Les Phytoseiidæ de la vigne en France, *Proc. Intern. Conf. on "Pests in Agriculture"*, Paris, 1-3 décembre 1987, Ann. ANPP, 6 (3), ANPP, ed.: 241-249.
- Krogh P.H., Griffiths B., Demsar D., Bohanec M., Debeljak M., Andersen M.N., Sausse C., Birch A.N.E., Caul S., Holmstrup M., Heckmann L.H., Cortet J. (2007). Responses by earthworms to reduced tillage in herbicide tolerant maize and Bt maize cropping systems. *Pedobiologia* 51(3): 219-227.
- Kromp B. (1999). Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agriculture, Ecosystems & Environment* 74(1-3): 187-228.

- Krooss S., Schaefer M. (1998). The effect of different farming systems on epigeic arthropods: a five-year study on the rove beetle fauna (Coleoptera: Staphylinidae) of winter wheat. *Agriculture, Ecosystems & Environment* 69(2): 121-133.
- Kruess A., Tscharntke T. (1994). Habitat fragmentation, species loss and biological control. *Science* 264(5165): 1581-1584.
- Kruess A., Tscharntke T. (2002). Contrasting responses of plant and insect diversity to variation in grazing intensity. *Biological Conservation* 106(3): 293-302.
- Kruess A., Tscharntke T. (2002). Grazing intensity and the diversity of grasshoppers, butterflies, and trap-nesting bees and wasps. *Conservation Biology* 16(6): 1570-1580.
- Kumar S., Stohlgren T.J., Chong G.W. (2006). Spatial heterogeneity influences native and nonnative plant species richness. *Ecology* 87(12): 3186-3199.
- Kumarasinghe N.C. (1991). Studies on the damage by termites to seed sets and to post-harvest stubbles of sugarcane plantations in the dry zone of Sri Lanka. In *Advances in management and conservation of soil fauna*, (Veeresh G.K. et al., eds.), Oxford and IBH, New Delhi: 147-154.
- Kwon Y.W., Kim D.S. (2001). Herbicide-resistant genetically-modified crop: its risks with an emphasis on gene flow. *Weed Biology and Management* 1(1): 42-52.
- Lagerlof J., Goffre B., Vincent C. (2002). The importance of field boundaries for earthworms (Lumbricidae) in the Swedish agricultural landscape. *Agriculture, Ecosystems & Environment* 89(1/2): 91-103
- Lagerlof J., Wallin H. (1993). The Abundance of Arthropods Along 2 Field Margins with Different Types of Vegetation Composition - an Experimental-Study. *Agriculture, Ecosystems & Environment* 43(2): 141-154.
- Lalande R., Gagnon B., Simard R.R., Cote D. (2000). Soil microbial biomass and enzyme activity following liquid hog manure application in a long-term field trial. *Canadian Journal of Soil Science* 80(2): 263-269.
- Lamarque F., Artois M., Berny P., Hatier C. (1999). Réseau SAGIR : douze ans de toxico-vigilance. *Bulletin mensuel O.N.C.* 246: 18-25.
- Lamarque F., Barrat J., Hatier C., Artois M. (1999). Causes of mortality in Roe deer (*Capreolus capreolus*) diagnosed by an epidemiological surveillance network in France. *Gibier Faune Sauvage* 16(2): 101-122.
- LaMondia J.A., Elmer W.H., Mervosh T.L., Cowles R.S. (2002). Integrated management of strawberry pests by rotation and intercropping. *Crop Protection* 21(9): 837-846.
- Lande R. (1996). Statistics and partitioning of species diversity, and similarity among multiple communities. *Oikos* 76: 5-13.
- Landis D.A., Menalled F.D., Costamagna A.C., Wilkinson T.K. (2005). Manipulating plant resources to enhance beneficial arthropods in agricultural landscapes. *Weed Science* 53(6): 902-908.
- Landis D.A., Wratten S.D., Gurr G.M. (2000). Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology* 45: 175-201.
- Landsberg J., Crowley G. (2004). Monitoring rangeland biodiversity: plants as indicators. *Austral Ecology* 29(1): 59-77.
- Langanke T., Rossner G., Vrscaj B., Lang S., Mitchley J. (2005). Selection and application of spatial indicators for nature conservation at different institutional levels. *Journal for Nature Conservation* 13(2/3): 101-114.
- Langellotto G.A., Denno R.F. (2004). Responses of invertebrate natural enemies to complex-structured habitats: a meta-analytical synthesis. *Oecologia* 139(1): 1-10.
- Langhof M., Gathmann A., Poehling H.M. (2003). Field study on the effects of insecticide drift on non-target arthropods: residual toxicity, effect on populations and recolonisation processes. *Bulletin OILB/SROP* 26(5): 73-82.
- Lannou C. (2001). Intrapathotype Diversity for Aggressiveness and Pathogen Evolution in Cultivar Mixtures. *Phytopathology* 91(5): 500-510.
- Lannou C., Vallavieille-Pope C., Goyeau H. (1995). Induced resistance in host mixtures and its effect on disease control in computer-simulated epidemics. *Plant Pathology* 44(3): 478-489.
- Lapointe S., Giroux J.F., Belanger L., Filion B. (2000). Benefits of rotational grazing and dense nesting cover for island-nesting waterfowl in southern Quebec. *Agriculture, Ecosystems & Environment* 78(3): 261-272.
- Larink O. (1997). Springtails and mites: important knots in the food web of soils. In *Fauna in soil ecosystems : recycling, processes, nutrient fluxes and agricultural production*, (Benckiser G., ed.), Marcel Dekker, New York: 225-264.
- Larkin R.P. (2003). Characterization of soil microbial communities under different potato cropping systems by microbial population dynamics, substrate utilization, and fatty acid profiles. *Soil Biology & Biochemistry* 35(11): 1451-1466.
- Larkin R.P., Griffin T.S. (2007). Control of soilborne potato diseases using Brassica green manures. *Crop Protection* 26(7): 1067-1077.
- Larkin R.P., Honeycutt C.W. (2006). Effects of different 3-year cropping systems on soil microbial communities and rhizoctonia diseases of potato. *Phytopathology* 96(1): 68-79.
- Lassau S.A., Cassis G., Flemons P.K.J., Wilkie L., Hochuli D.F. (2005). Using high-resolution multi-spectral imagery to estimate habitat complexity in open-canopy forests: can we predict ant community patterns? *Ecography* 28(4): 495-504.
- Lavelle P., Blanchart E., Martin A., Martin S., Spain A., Toutain F., Barois I., Schaefer R. (1993). A Hierarchical Model for Decomposition in Terrestrial Ecosystems - Application to Soils of the Humid Tropics. *Biotropica* 25(2): 130-150.
- Lavelle P., Spain A.V. (2001). Soil ecology, Kluwer Academic Publishers, Netherlands, 654 p.
- Lavoisier E., Léon F., Havet P., Granval P. (1996). Intérêt de la jachère pour réduire les dégâts occasionnés par la grande faune sauvage. *Bulletin mensuel O.N.C.* 124: 50-55.
- Lavorel S., Garnier E. (2002). Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology* 16: 545-556.
- Lavorel S., Quetier F., Gaucherand S., Choler P. (2004). Contribution of the functional characteristics of plants to the ecological assessment of the evolutive patterns of management in grassland environments. *Fourrages*(No.178): 179-191.
- Lazzeri L., Baruzzi G., Malaguti L., Antoniaci L. (2003). Replacing methyl bromide in annual strawberry production with glucosinolate-containing green manure crops. *Pest Management Science* 59(9): 983-990.
- Le Coeur D., Baudry J., Burel F., Thenail C. (2002). Why and how we should study field boundaries biodiversity in an agrarian landscape context. *Agriculture, Ecosystems & Environment* 89: 23-40.
- Leblanc M.L., Cloutier D.C. (1996). Effet de la technique du faux semis sur la levée des adventives annuelles, X. *Colloque International sur la Biologie des mauvaises herbes*, Dijon (France), Septembre 1996, AFPP, ed.: 29-34.

- Legere A., Samson N. (2004). Tillage and weed management effects on weeds in barley-red clover cropping systems. *Weed Science* 52: 881-885.
- Lena F. (2002). Etude des cohabitations entre ongulés domestiques et sauvages dans le Parc National de la Vanoise et caractérisation des risques de transmission des maladies. *Bulletin d'Informations sur la Pathologie des Animaux Sauvages en France* 25: 119-130.
- Lenné J.M., Wood D. (1999). Optimizing biodiversity for productive agriculture. In *Agrobiodiversity: Characterization, Utilization and Management*, (Wood D., Lenné J.M., eds.), CABI Publishing, Oxon UK, New-York USA: 447-470.
- Lepart J., Escarré J. (1983). La succession végétale, mécanismes et modèles: analyse bibliographique. *Bulletin d'Ecologie* 14(3): 133-178.
- Lepš J. (2005). Diversity and ecosystem function. In *Vegetation Ecology*, (van der Maarel E., ed.), Blackwell Publishing, Malden: 199-237.
- Leslie T.W., Hoheisel G.A., Biddinger D.J., Rohr J.R., Fleischer S.J. (2007). Transgenes sustain epigeal insect biodiversity in diversified vegetable farm systems. *Environmental Entomology* 36(1): 234-244.
- Lester P.J., Thistlewood H.M.A., Marshall D.B., Harmsen R. (1999). Assessment of *Amblyseius fallacis* (Acari: Phytoseiidae) for biological control of tetranychid mites in an Ontario peach orchard. *Experimental & Applied Acarology* 23(12): 995-1009.
- Letourneau D.K. (1988). Soil management for pest control: a critical appraisal of the concepts, *Global Perspectives on Agroecology and Sustainable Agricultural Systems. Proceedings of the Sixth International Scientific Conference of IFOAM*, Santa Cruz: 581-587.
- Levrel H. (2007). Quels indicateurs pour la gestion de la biodiversité ? , Cahier de l'IFB (Weber J., ed.), 1, Institut français de la biodiversité, Paris, 99 p.
- Liira J., Schmidt T., Aavik T., Arens P., Augenstein I.I., Bailey D., Billeter R., Bukacek R., Burel F., De Blust G., De Cock R., Dirksen J., Edwards P.J., Hamersky R., Herzog F., Klotz S., Kühn I., Le Coeur D., Miklova P., Roubalova M., Schweiger O., Smulders M.J.M., van Wingerden W.K.R.E., Bugter R., Zobel M. (2008). Plant functional group composition and large-scale species richness in the agricultural landscapes of Europe. *Journal of Vegetation Science* 19(1): 3-14.
- Limpens H.J.G.A., Kapteyn K. (1991). Bats, their behaviour and linear landscape elements. *Myotis* 29: 39-48.
- Lindborg R., Eriksson O. (2004). Historical landscape connectivity affects present plant species diversity. *Ecology* 85(7): 1840-1845.
- Lindenmayer D.B., Fischer J. (2007). Incorporating edge effects into landscape management and design. In *Managing and designing landscapes for conservation: moving from perspectives to principles* (Lindenmayer D.B., Hobbs R.J., eds.), Blackwell Publishing, Oxford: 165-178.
- Liss W.J., Gut L.J., Westgard P.H., Warren C.E. (1986). Perspectives on Arthropod Community Structure, Organization, and Development in Agricultural Crops. *Annual Review of Entomology* 31: 455-478.
- Loe L.E., Myserud A., Stien A., Steen H., Evans D.M., Austrheim G. (2007). Positive short-term effects of sheep grazing on the alpine avifauna. *Biology Letters* 3(1): 109-111.
- LopezFando C., Bello A. (1995). Variability in soil nematode populations due to tillage and crop rotation in semi-arid Mediterranean agrosystems. *Soil & Tillage Research* 36(1-2): 59-72.
- Lopez-Ridaura S., van Keulen H., van Ittersum M.K., Lefelaar P.A. (2005). Multiscale methodological framework to derive criteria and indicators for sustainability evaluation of peasant natural resource management systems. *Environment Development and Sustainability* 7(1): 51-69.
- Losey J.E., Rayor L.S., Carter M.E. (1999). Transgenic pollen harms monarch larvae. *Nature* 399: 214.
- Louault F., Pillar V.D., Aufrère J., Garnier E., Soussana J.-F. (2005). Plant traits and functional types in response to reduced disturbance in a semi-natural grassland. *Journal of Vegetation Science* 16: 151-160.
- Loucougaray G., Bonis A., Bouzille J.B. (2004). Effects of grazing by horses and/or cattle on the diversity of coastal grasslands in western France. *Biological Conservation* 116(1): 59-71.
- Louda S.M., Kendall D., Conner J., Simberloff D. (1997). Ecological effects of an insect introduced for the biological control of weeds. *Science* 277: 1088-1090.
- Louvet J., Rouxel F., Alabouvette C. (1976). Recherches sur la résistance des sols aux maladies. I - Mise en évidence de la nature microbiologique de la résistance d'un sol au développement de la fusariose vasculaire du melon. *Annales de Phytopathologie* 8(4): 425-436.
- Lowery D.T., Isman M.B. (1994). Effects of Neem and Azadirachtin on Aphids and Their Natural Enemies. In *Bioregulators for Crop Protection and Pest Control*, Acs Symposium Series, 557, AMER CHEMICAL SOC, Washington: 78-91.
- Ludy C., Lang A. (2006). A 3-year field-scale monitoring of foliage-dwelling spiders (Araneae) in transgenic Bt maize fields and adjacent field margins. *Biological Control* 38(3): 314-324.
- Luka H., Uehlinger G., Pfiffner L., Muhlethaler R., Blick T. (2006). Extended field margins - a new element of ecological compensation in farmed landscapes - deliver positive impacts for Articulata. *Agrarforschung* 13(9): 386-391.
- Lumaret J.P., Errouissi F. (2002). Use of anthelmintics in herbivores and evaluation of risks for the non target fauna of pastures. *Veterinary Research* 33(5): 547-562.
- Lumaret J.P., Errouissi F., Galtier P., Alvinerie M. (2005). Pour on formulation of eprinomectin for cattle: Fecal elimination profile and effects on the development of the dung-inhabiting diptera *Neomyia cornicina* (L.) (Muscidae). *Environmental Toxicology and Chemistry* 24(4): 797-801.
- Luoto M., Pykala J., Kuussaari M. (2003). Decline of landscape-scale habitat and species diversity after the end of cattle grazing. *Journal for Nature Conservation* 11(3): 171-178.
- Lupwayi N.Z., Arshad M.A., Rice W.A., Clayton G.W. (2001). Bacterial diversity in water-stable aggregates of soils under conventional and zero tillage management. *Applied Soil Ecology* 16(3): 251-261.
- Lupwayi N.Z., Clayton G.W., O'Donovan J.T., Harker K.N., Turkington T.K., Rice W.A. (2004). Soil microbiological properties during decomposition of crop residues under conventional and zero tillage. *Canadian Journal of Soil Science* 84(4): 411-419.
- Lupwayi N.Z., Hanson K.G., Harker K.N., Clayton G.W., Blackshaw R.E., O'Donovan J.T., Johnson E.N., Gan Y., Irvine R.B., Monreal M.A. (2007). Soil microbial biomass, functional diversity and enzyme activity in glyphosate-resistant wheat-canola rotations under low-disturbance direct seeding and conventional tillage. *Soil Biology & Biochemistry* 39(7): 1418-1427.
- Lupwayi N.Z., Monreal M.A., Clayton G.W., Grant C.A., Johnston A.M., Rice W.A. (2001). Soil microbial biomass and diversity respond to tillage and sulphur fertilizers. *Canadian Journal of Soil Science* 81(5): 577-589.

- Lupwayi N.Z., Rice W.A., Clayton G.W. (1998). Soil microbial diversity and community structure under wheat as influenced by tillage and crop rotation. *Soil Biology & Biochemistry* 30(13): 1733-1741.
- Mac Nally R., Bennett A.F., Brown G.W., Lumsden L.F., Yen A., Hinkley S., Lillywhite P., Ward D.A. (2002). How well do ecosystem-based planning units represent different components of biodiversity? *Ecological Applications* 12(3): 900-912.
- Mac Nally R., Ellis M., Barrett G. (2004). Avian biodiversity monitoring in Australian rangelands. *Austral Ecology* 29(1): 93-99.
- Madge D.S. (1981). Influence of agricultural practice on soil invertebrate animals. In *Biological Husbandry: a scientific approach to organic farming*, (Stonehouse B., ed.), Butterworths, London: 79-98.
- Maes D., Bauwens D., De Bruyn L., Anselin A., Vermeersch G., Van Landuyt W., De Knijf G., Gilbert M. (2005). Species richness coincidence: conservation strategies based on predictive modelling. *Biodiversity and Conservation* 14(6): 1345-1364.
- Maestre F.T., Reynolds J.F. (2006). Spatial heterogeneity in soil nutrient supply modulates nutrient and biomass responses to multiple global change drivers in model grassland communities. *Global Change Biology* 12(12): 2431-2441.
- Maherali H., Klironomos J.N. (2007). Influence of Phylogeny on fungal community assembly and ecosystem functioning. *Science* 316(5832): 1746-1748.
- Maillard A., Cuendet G. (1997). Résultats d'un essai de culture sans labour depuis plus de 20 ans à Changins - Propriétés biologiques du sol et vers de terre. *Revue suisse d'Agriculture* 29(5): 225-230.
- Mair J., Port G.R. (2002). The influence of mucus production by the slug, *Deroceras reticulatum*, on predation by *Pterostichus madidus* and *Nebria brevicollis* (Coleoptera: Carabidae). *Biocontrol Science and Technology* 12(3): 325-335.
- Makeschin F. (1997). Earthworms (Lumbricidae: Oligochaeta): Important promoters of soil development and soil fertility. In *Fauna in soil ecosystems: recycling, processes, nutrient fluxes and agricultural production*, (Benckiser G., ed.), Marcel Dekker: 173-223.
- Malone L.A., Pham-Delègue M.-H. (2001). Effects of transgene products on honey bees (*Apis mellifera*) and bumblebees (*Bombus* sp.). *Apidologie* 32: 287-304.
- Mandrillon A.L., Saglio P. (2005). Une revue des effets des pesticides sur la morphologie, le comportement et les traits d'histoire de vie des amphibiens. *Bulletin de la Société herpétologique de France* 116: 5-29.
- Manne L.L., Williams P.H. (2003). Building indicator groups based on species characteristics can improve conservation planning. *Animal Conservation* 6: 291-297.
- Marchandeu F. (1992). Faune sauvage et faune domestique en milieu pastoral : une synthèse bibliographique. *Gibier Faune sauvage* 9(2): 167-186.
- Marhan S., Scheu S. (2005). The influence of mineral and organic fertilisers on the growth of the endogeic earthworm *Octolasion tytaeum* (Savigny). *Pedobiologia* 49(3): 239-249.
- Marini L., Scotton M., Klimek S., Isselstein J., Pecile A. (2007). Effects of local factors on plant species richness and composition of Alpine meadows. *Agriculture, Ecosystems & Environment* 119(3-4): 281-288.
- Marinissen J.C.Y. (1992). Population dynamics of earthworms in a silt loam soil under conventional and "integrated" arable farming during two years with different weather patterns. *Soil Biology & Biochemistry* 24(12): 1647-1654.
- Markovic M., Zivanovic M. (1988). Ecological selectivity of some acaricides to the predators of *Panonychus ulmi* Koch. *Zastita Bilja* 39(2): 183-195.
- Marriott C.A., Fothergill M., Jeangros B., Scotton M., Louault F. (2004). Long-term impacts of extensification of grassland management on biodiversity and productivity in upland areas. A review. *Agronomie* 24(8): 447-462.
- Marrs R.H., Lowday J.E. (1992). Control of bracken and the restoration of heathland. II. Regeneration of the heathland community. *Journal of Applied Ecology* 29(1): 204-211.
- Marshall E., J.P. (2001). Biodiversity, herbicides and non target plants, *The BCPC Conference - Weeds 2001*: 855-862.
- Marshall E., J. P., Moonen A., C. (2002). Field margins in northern Europe: their functions and interactions with agriculture. *Agriculture Ecosystems and Environment* 89: 5-21.
- Marshall E.J., Smith B.D. (1987). Field margin flora and fauna; interaction with agriculture. In *British Crop Protection Council (BCPC) Monograph, 35*, British Crop Protection Council: 23-32.
- Marshall E.J.P. (2004). Agricultural landscapes: field margin habitats and their interaction with crop production. *Journal of Crop Improvement* 12(1/2): 365-404.
- Marshall E.J.P. (2005). Field margins in northern Europe: integrating agricultural, environmental and biodiversity functions. . In *Field boundary habitats: implications for weed, insect and disease management*, (Thomas A.G., ed.), Topics in Canadian Weed Science, 1, Canadian Weed Science Society: 39-67.
- Marshall E.J.P., West T.M., Kleijn D. (2006). Impacts of an agri-environment field margin prescription on the flora and fauna of arable farmland in different landscapes. *Agriculture, Ecosystems & Environment* 113(1-4): 36-44.
- Martin-Laurent F., Barres B., Wagschal I., Piutti S., Devers M., Soulas G., Philippot L. (2006). Impact of the maize rhizosphere on the genetic structure, the diversity and the atrazine-degrading gene composition of cultivable atrazine-degrading communities. *Plant and Soil* 282(1-2): 99-115.
- Martin-Laurent F., Cornet L., Ranjard L., Lopez-Gutierrez J.C., Philippot L., Schwartz C., Chaussod R., Catroux G., Soulas G. (2004). Estimation of atrazine-degrading genetic potential and activity in three French agricultural soils. *FEMS Microbiology Ecology* 48(3): 425-435.
- Marulanda A., Azcon R., Ruiz-Lozano J.M. (2003). Contribution of six arbuscular mycorrhizal fungal isolates to water uptake by *Lactuca sativa* plants under drought stress. *Physiologia Plantarum* 119(4): 526-533.
- Marvier M., McCreedy C., Regetz J., Kareiva P. (2007). A Meta-Analysis of Effects of Bt Cotton and Maize on Nontarget Invertebrates. *Science* 316: 1475-1477.
- Maud J., Edwards-Jones G., Quin F. (2001). Comparative evaluation of pesticide risk indices for policy development and assessment in the United Kingdom. *Agriculture Ecosystems & Environment* 86(1): 59-73.
- Maurizi B., Verrel J.-L. (2002). Des indicateurs pour des actions de maîtrise des pollutions d'origine agricole. *Ingénieries* 30: 3-14.
- Mazzola M. (2004). Assessment and management of soil microbial community structure for disease suppression. *Annual Review of Phytopathology* 42: 35-59.
- Mazzola M. (2007). Manipulation of rhizosphere bacterial communities to induce suppressive soils. *Journal of Nematology* 39(3): 213-220.

- McCloskey C., Arnason J.T., Donskov N., Chenier R., Kaminski J., Philogene B.J.R. (1993). Third trophic level effects of azadirachtin. *Canadian Entomologist* 125(1): 163-165.
- McCrea A.R., Trueman I.C., Fullen M.A. (2004). Factors relating to soil fertility and species diversity in both semi-natural and created meadows in the West Midlands of England. *European Journal of Soil Science* 55(2): 335-348.
- McEvoy P.B. (1996). Host specificity and biological pest control. *BioScience* 46: 401-405.
- McGeoch M.A. (1998). The selection, testing and application of terrestrial insects as bioindicators. *Biological Reviews of the Cambridge Philosophical Society* 73(2): 181-201.
- McIntyre S., Heard K.M., Martin T.G. (2003). The relative importance of cattle grazing in subtropical grasslands: does it reduce or enhance plant biodiversity? *Journal of Applied Ecology* 40: 445-457
- McIntyre S., Lavorel S. (2001). Livestock grazing in subtropical pastures: steps in the analysis of attribute response and plant functional types. *Journal of Ecology* 89: 209-226.
- McIntyre S., Lavorel S., Landsberg J., Forbes T.D.A. (1999). Disturbance response in vegetation - towards a global perspective on functional traits. *Journal of Vegetation Science* 10: 621-630.
- McKinley V.L., Peacock A.D., White D.C. (2005). Microbial community PLFA and PHB responses to ecosystem restoration in tallgrass prairie soils. *Soil Biology & Biochemistry* 37(10): 1946-1958.
- McLaughlin A., Mineau P. (1995). The impact of agricultural practices on biodiversity. *Agriculture, Ecosystems & Environment* 55(3): 201-212.
- McSorley R., Dickson D.W., Debrito J.A., Hochmuth R.C. (1994). Tropical Rotation Crops Influence Nematode Densities and Vegetable Yields. *Journal of Nematology* 26(3): 308-314.
- Meerts P. (1993). La régression des plantes messicoles en Belgique, *Faut-il sauver les mauvaises herbes ?*, Gap (France), Conservatoire Botanique de Gap-Charance: 49-55.
- Mellinger M.V., McNaughton S.J. (1975). Structure and function of successional vascular plant communities in Central New York. *Ecological Monographs* 45: 161-182.
- Menalled F.D., Smith R.G., Dauer J.T., Fox T.B. (2007). Impact of agricultural management on carabid communities and weed seed predation. *Agriculture, Ecosystems & Environment* 118(1-4): 49-54.
- Mendel Z., Blumberg D., Zehavi A., Weissenberg M. (1992). Some polyphagous homoptera gain protection from their natural enemies by feeding on the toxic plants *Spartium junceum* and *Erythrina corallodendrum* (Leguminosae). *Chemoecology* 3: 118-124.
- Mendras H. (1984). *La fin des paysans*, Babel, Actes-Sud, Arles, 436 p.
- Mercier J., Lindow S.E. (2001). Field performance of antagonistic bacteria identified in a novel laboratory assay for biological control of fire blight of pear. *Biological Control* 22(1): 66-71.
- Messéan A., Angevin F., Gómez-Barbero M., Menrad K., Rodríguez-Cerezo E. (2006). New case studies on the coexistence of GM and non-GM crops in European agriculture, Technical Report EUR 22102 EN, Institute for Prospective Technological Studies, 112 p.
- Meyer-Aurich A. (2005). Economic and environmental analysis of sustainable farming practices - a Bavarian case study. *Agricultural Systems* 86(2): 190-206.
- Michaud J.P. (2002). Invasion of the Florida citrus ecosystem by *Harmonia axyridis* (Coleoptera: Coccinellidae) and asymmetric competition with a native species, *Cycloneda sanguinea*. *Environmental Entomology* 31(5): 827-835.
- Milai Canals L., Bauer C., Depestele J., Dubreuil A., Knuchel R.F., Gaillard G., Michelsen O., Muller-Wenk R., Rydgren B. (2007). Key elements in a framework for land use impact assessment within LCA. *International Journal of Life Cycle Assessment* 12(1): 5-15.
- Milchunas D.G., Sala O.E., Lauenroth W.K. (1988). A generalized model of the effects of grazing by large herbivores on grassland community structure. *American Naturalist* 132: 87-106.
- Mildén M., Münzbergová Z., Herben T., Ehrlén J. (2006). Metapopulation dynamics of a perennial plant, *Succisa pratensis*, in an agricultural landscape *Ecological Modelling* 199(4): 464-475.
- Miliczky E.R., Horton D.R. (2005). Densities of beneficial arthropods within pear and apple orchards affected by distance from adjacent native habitat and association of natural enemies with extra-orchard host plants. *Biological Control* 33(3): 249-259.
- Millan de la Pena N.M., Butet A., Delettre Y., Morant P., Burel F. (2003). Landscape context and carabid beetles (Coleoptera : Carabidae) communities of hedgerows in western France. *Agriculture, Ecosystems & Environment* 94(1): 59-72.
- Mineau P. (2002). Estimating the probability of bird mortality from pesticide sprays on the basis of the field study record. *Environmental Toxicology and Chemistry* 21: 1497-1506.
- Mineau P. (2005). Direct losses of birds to pesticides - Beginnings of a quantification, *Bird Conservation Implementation and Integration in the Americas*, 3. International Partners in Flight Conference 2002, Ralph C.J., Rich T.D., eds.: 1065-1070.
- Mineau P., Downes C.M., Kirk D.A., Bayne E., Csizy M. (2005). Patterns of bird species abundance in relation to granular insecticide use in the Canadian prairies *Ecoscience* 12(2): 267-278.
- Mineau P., McLaughlin A. (1996). Conservation of biodiversity within Canadian agricultural landscapes : integrating habitat for wildlife. *Journal of Agricultural and Environmental Ethics* 9(2): 93-113.
- Mineau P., Whiteside M. (2006). Lethal risk to birds from insecticide use in the United States: a spatial and temporal analysis. *Environmental Toxicology and Chemistry* 25(5): 1214-1222.
- Mitchell C.E., Tilman D., Groth J.V. (2002). Effects of grassland plant species diversity, abundance, and composition on foliar fungal disease. *Ecology* 83(6): 1713-1726.
- Mitchell G., May A., McDonald A. (1995). Picabue - a Methodological Framework for the Development of Indicators of Sustainable Development. *International Journal of Sustainable Development and World Ecology* 2(2): 104-123.
- Mitchell-Jones A.J. (1998). Landscapes for greater horseshoe bats, *Enact*, English Nature, 6: 11-14.
- Monetti L.N., Fernandez N.A. (1995). Seasonal population dynamics of the European red mite (*Panonychus ulmi*) and its predator *Neoseiulus californicus* in a sprayed apple orchard in Argentina (Acari: Tetranychidae, Phytoseiidae). *Acarologia* 36(4): 325-331.
- Moonen A.C., Marshall E.J.P. (2001). The influence of sown margin strips, management and boundary structure on herbaceous field margin vegetation in two neighbouring farms in southern England. *Agriculture, Ecosystems & Environment* 86(2): 187-202.

- Moore J.C., Berlow E.L., Coleman D.C., de Ruiter P.C., Dong Q., Hastings A., Johnson N.C., McCann K.S., Melville K., Morin P.J., Nadelhoffer K., Rosemond A.D., Post D.M., Sabo J.L., Scow K.M., Vanni M.J., Wall D.H. (2004). Detritus, trophic dynamics and biodiversity. *Ecology Letters* 7(7): 584-600.
- Moore J.L., Balmford A., Brooks T., Burgess N.D., Hansen L.A., Rahbek C., Williams P.H. (2003). Performance of sub-Saharan vertebrates as indicator groups for identifying priority areas for conservation. *Conservation Biology* 17(1): 207-218.
- Morales H., Perfecto I., Ferguson B. (2001). Traditional fertilization and its effect on corn insect populations in the Guatemalan highlands. *Agriculture, Ecosystems & Environment* 84(2): 145-155.
- Moreby S.J., Southway S.E. (1999). Influence of autumn applied herbicides on summer and autumn food available to birds in winter wheat fields in southern England. *Agriculture Ecosystems & Environment* 72(3): 285-297.
- Moreno C.E., Sanchez-Rojas G., Pineda E., Escobar F. (2007). Shortcuts for biodiversity evaluation: a review of terminology and recommendations for the use of target groups, bioindicators and surrogates. *International Journal of Environment and Health* 1(1): 71-86.
- Moser B.W., Witmer G.W. (2000). The effects of elk and cattle foraging on the vegetation, birds, and small mammals of the Bridge Creek Wildlife Area, Oregon. *International Biodeterioration & Biodegradation* 45(3-4): 151-157.
- Moss S.R., Clarke J. (1994). Guidelines for the prevention and control of herbicide-resistant blackgrass (*Alopecurus myosuroides*). *Crop Protection* 13(3): 230-234.
- Motavalli P.P., Kremer R.J., Fang M., Means N.E. (2004). Impact of genetically modified crops and their management on soil microbially mediated plant nutrient transformations. *Journal of Environmental Quality* 33(3): 816-824.
- Motte G., Libois R. (2002). Conservation of the lesser horseshoe bat (*Rhinolophus hipposideros* Bechstein, 1800) (Mammalia : Chiroptera) in Belgium. A case study of feeding habitat requirements. *Belgian Journal of Zoology* 132(1): 49-54.
- Mougel C., Offre P., Ranjard L., Corberand T., Gamalero E., Robin C., Lemanceau P. (2006). Dynamic of the genetic structure of bacterial and fungal communities at different developmental stages of Medicago truncatula Gaertn. cv. Jemalong line J5. *New Phytologist* 170(1): 165-175.
- Mountford J.O., Lakhani K.H., Holland R.J. (1996). Reversion of grassland vegetation following the cessation of fertilizer application. *Journal of Vegetation Science* 7(2): 219-228.
- Moutier J.N. (2006). Le suivi des dégâts de Chevreuil sur plantations en terre agricole. *Forêt Entreprise* 167: 50-52.
- Mulder C., de Zwart D., van Wijnen H.J., Schouten A.J., Breure A.M. (2003). Observational and simulated evidence of ecological shifts within the soil nematode community of agroecosystems under conventional and organic farming. *Functional Ecology* 17(4): 516-525.
- Mulder C., van Wijnen H.J., van Wezel A.P. (2005). Numerical abundance and biodiversity of below-ground taxocenes along a pH gradient across the Netherlands. *Journal of Biogeography* 32(10): 1775-1790.
- Müller-Schärer H., Scheepens P.C., Greaves M.P. (2000). Biological control of weeds in European crops: recent achievements and future work. *Weed Research* 40(1): 83-98.
- Mundt C.C. (2002). Use of multiline cultivars and cultivar mixtures for disease management. *Annual Review of Phytopathology* 40: 381-410.
- Mundt C.C., Cowger C., Garrett K.A. (2002). Relevance of integrated disease management to resistance durability. *Euphytica* 124(2): 245-252.
- Murray P.J., Cook R., Currie A.F., Dawson L.A., Gange A.C., Grayston S.J., Treonis A.M. (2006). Interactions between fertilizer addition, plants and the soil environment: Implications for soil faunal structure and diversity. *Applied Soil Ecology* 33(2): 199-207.
- Myers W., Patil G.P. (2006). Biodiversity in the age of ecological indicators. *Acta Biotheoretica* 54(2): 119-123.
- Myklestad A., Saetersdal M. (2004). The importance of traditional meadow management techniques for conservation of vascular plant species richness in Norway *Biological Conservation* 118(2): 133-139.
- Mysterud A., Hansen L.O., Peters C., Austrheim G. (2005). The short-term effect of sheep grazing on selected invertebrates (Diptera and Hemiptera) relative to other environmental factors in an alpine ecosystem. *Journal of Zoology* 266: 411-418.
- Nakamoto T., Tsukamoto M. (2006). Abundance and activity of soil organisms in fields of maize grown with a white clover living mulch. *Agriculture, Ecosystems & Environment* 115(1-4): 34-42.
- Nardo M., Saisana M., Saltelli A., Tarantola S. (2005). Tools for composite indicators building, Joint Research Center, European Commission, JRC EUR 21682 report 134 p. http://composite-indicators.jrc.ec.europa.eu/Document/EUR%2021682%20EN_Tools_for_Composite_Indicator_Building.pdf
- Navas M.-L., Ducout B., Roumet C., Richarte J., Garnier J., Garnier E. (2003). Leaf life span, dynamics and construction cost of species from Mediterranean old-fields differing in successional status. *New Phytologist* 159: 213-228.
- Neher D.A. (1995). Biological diversity in soils of agricultural and natural ecosystems. In *Exploring the role of diversity in sustainable agriculture*, (Olson R.K. et al., eds.), American Society of Agronomy, Madison, WI, USA: 55-72.
- Neher D.A., Barbercheck M.E. (1999). Diversity and function of soil mesofauna. In *Biodiversity in agroecosystems*, (Collins W.W., Qualset C.O., eds.), CRC Press, Boca Raton, London, New York, Washington D.C.: 27-48.
- Neve P. (2007). Challenges for herbicide resistance evolution and management: 50 years after Harper. *Weed Research* 47: 365-369.
- Nicolardot B., Bouziri L., Bastian F., Ranjard L. (2007). A microcosm experiment to evaluate the influence of location and quality of plant residues on residue decomposition and genetic structure of soil microbial communities. *Soil Biology & Biochemistry* 39(7): 1631-1644.
- Niehoff B., Poehling H.M. (1995). Population dynamics of aphids and syrphid larvae in winter wheat treated with different rates of pirimicarb. *Agriculture, Ecosystems & Environment* 52(1): 51-55.
- Niemela J., Baur B. (1998). Threatened species in a vanishing habitat: plants and invertebrates in calcareous grasslands in the Swiss Jura mountains. *Biodiversity and Conservation* 7(11): 1407-1416.
- Noe E., Halberg N., Reddersen J. (2005). Indicators of biodiversity and conservational wildlife quality on danish organic farms for use in farm management: A multidisciplinary approach to indicator development and testing. *Journal of Agricultural & Environmental Ethics* 18(4): 383-414.

- Norris K., Brindley E., Cook T., Babbs S., Brown C.F., Yaxley R. (1998). Is the density of redshank *Tringa totanus* nesting on saltmarshes in Great Britain declining due to changes in grazing management? *Journal of Applied Ecology* 35(5): 621-634.
- Norris K., Cook T., Odowd B., Durdin C. (1997). The density of redshank *Tringa totanus* breeding on the salt-marshes of the Wash in relation to habitat and its grazing management. *Journal of Applied Ecology* 34(4): 999-1013.
- Norris R.F., Kogan M. (2005). Ecology of interactions between weeds and arthropods. *Annual Review of Entomology* 50: 479-503.
- Noy-Meir I., Gutman M., Kaplan Y. (1989). Responses of Mediterranean grassland plants to grazing and protection. *Journal of Ecology* 77: 290-310.
- Oaks J.L., Gilbert M., Virani M.Z., Watson R.T., Meteyer C.U., Rideout B.A., Shivaprasad H.L., Ahmed S., Chaudhry M.J.I., Arshad M., Mahmood S., Ali A., Khan A.A. (2004). Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* 427(6975): 630-633.
- OECD (2001). Agriculture and Biodiversity : developing indicators for policy analysis Proceedings from OECD Expert Meeting on Agri-biodiversity Indicators, Zürich, Switzerland, 2001/11/05-08, 280 p.
- Oertli S., Muller A., Steiner D., Breitenstein A., Dorn S. (2005). Cross-taxon congruence of species diversity and community similarity among three insect taxa in a mosaic landscape. *Biological Conservation* 126(2): 195-205.
- Oindo B.O. (2002). Patterns of herbivore species richness in Kenya and current ecoclimatic stability. *Biodiversity and Conservation* 11(7): 1205-1221.
- Ojiambo P.S., Scherm H. (2006). Biological and application-oriented factors influencing plant disease suppression by biological control: A meta-analytical review. *Phytopathology* 96(11): 1168-1174.
- Oka Y., Shapira N., Fine P. (2007). Control of root-knot nematodes in organic farming systems by organic amendments and soil solarization. *Crop Protection* 26(10): 1556-1565.
- Oldroyd G.E.D. (2001). Dissecting symbiosis: Developments in Nod factor signal transduction. *Annals of Botany* 87(6): 709-718.
- Ollif H., Ritchie M.E. (1998). Effects of herbivores on grassland plant diversity. *Trends in Ecology & Evolution* 13(7): 261-265.
- Oppermann R. (2003). Nature balance scheme for farms - evaluation of the ecological situation. *Agriculture, Ecosystems & Environment* 98(1/3): 463-475
- Oppermann R., Braband D., Haack S. (2005). Nature indicators for agriculture. *Berichte über Landwirtschaft* 83(1): 76-102
- Orth D., Loiseau P., Loison A., Perrin O., Balay C. (2004). Un cas d'évaluation phytosociologique de la biodiversité: options et questions. *Fourrages* 179: 335-352.
- Osbornová J., Kovářová M., Lepš J., Prach K. (1990). Succession in abandoned fields. Studies in Central Bohemia, Czechoslovakia, Geobotany, (Werger M.J.A., ed.), 15, Kluwer Academic Publishers, Dordrecht, 168 p.
- Osem Y., Perevolotsky A., Kigel J. (2004). Site productivity and plant size explain the response of annual species to grazing exclusion in a Mediterranean semi-arid rangeland. *Journal of Ecology* 92: 297-309.
- Osterblad M., Norrdahl K., Korpimäki E., Huovinen P. (2001). Antibiotic resistance - How wild are wild mammals? *Nature* 409(6816): 37-38.
- Oxbrough A.G., Gittings T., O'Halloran J., Giller P.S., Kelly T.C. (2007). Biodiversity of the ground-dwelling spider fauna of afforestation habitats. *Agriculture Ecosystems & Environment* 120(2-4): 433-441.
- Pacha M.J., Petit S. (2008). The effect of landscape structure and habitat quality on the occurrence of *Geranium sylvaticum* in fragmented hay meadows. *Agriculture, Ecosystems & Environment* 123(1-3): 81-87.
- Paine L., Undersander D.J., Sample D.W., Bartelt G.A., Schatteman T.A. (1996). Cattle trampling of simulated ground nests in rotationally grazed pastures. *Journal of Range Management* 49(4): 294-300.
- Paine L., Undersander D.J., Sample D.W., Bartelt G.A., Schatteman T.A. (1997). Comparison of simulated ground nest types for grazing/trampling research. *Journal of Range Management* 50(3): 231-233.
- Pakeman R.J. (2004). Consistency of plant species and trait responses to grazing along a productivity gradient: a multi-site analysis. *Journal of Ecology* 92(5): 893-905.
- Palacio-Rabaud V. (2000). La conduite des prairies en France: pratiques intensives et rendements élevés dans le Nord-Ouest. *Agreste Cahiers* 4: 15-26.
- Palmer M.W. (1994). Variation in species richness: Towards a unification of hypotheses. *Folia Geobotanica et Phytotaxonomica* 29: 511-530.
- Paoletti M.G. (1999). The role of earthworms for assessment of sustainability and as bioindicators. *Agriculture, Ecosystems & Environment* 74(1-3): 137-155.
- Paoletti M.G., Bressan M. (1996). Soil invertebrates as bioindicators of human disturbance. *Critical Reviews in Plant Sciences* 15(1): 21-62.
- Paoletti M.G., Favretto M.R., Stinner B.R., Purrington F.F., Bajer J.E. (1991). Invertebrates as bioindicators of soil use. *Agriculture, Ecosystems & Environment* 34(1-4): 341-362.
- Parfitt R.L., Yeates G.W., Ross D.J., Mackay A.D., Budding P.J. (2005). Relationships between soil biota, nitrogen and phosphorus availability, and pasture growth under organic and conventional management. *Applied Soil Ecology* 28(1): 1-13.
- Pastor J., Muen R., Cohen Y. (1997). Spatial heterogeneity, carrying capacity and feedbacks in animal landscape interactions. *Journal of Mammalogy* 78: 1040-1052.
- Pastoret P.P., Thiry E., Brochier B., Schwers A., Thomas I., Dubuisson J. (1988). Maladies de la faune sauvage transmissibles aux animaux domestiques. *Revue scientifique et technique: office international des epizootics (O.I.E)* 7(4): 661-704.
- Patra A.K., Abbadie L., Clays A., Degrange V., Grayston S., Loiseau P., Louault F., Mahmood S., Nazaret S., Philippot L., Poly F., Prosser J.I., Richaume A., le Roux X. (2005). Effect of grazing on microbial groups involved in soil N dynamics. *Ecological Monographs* 75(1): 65-80.
- Patra A.K., Abbadie L., Clays-Josserand A., Degrange V., Grayston S.J., Guillaumaud N., Loiseau P., Louault F., Mahmood S., Nazaret S., Philippot L., Poly F., Prosser J.I., Le Roux X. (2006). Effects of management regime and plant species on the enzyme activity and genetic structure of N-fixing, denitrifying and nitrifying bacterial communities in grassland soils. *Environmental Microbiology* 8(6): 1005-1016.

- Patra A.K., Le Roux X., Abbadie L., Clays-Josserand A., Poly F., Loiseau P., Louault F. (2007). Effect of Microbial Activity and Nitrogen Mineralization on Free-living Nitrogen Fixation in Permanent Grassland Soils. *J. Agronomy & Crop Science* 193: 153-156.
- Patra A.K., Le Roux X., Grayston S.J., Loiseau P., Louault F. (2008). Unraveling the effects of management regime and plant species on soil organic carbon and microbial phospholipid fatty acid profiles in grassland soils. *Bioresource Technology* 99: 3545-3551.
- Pausas J.G., Austin M.P. (2001). Patterns of plant species richness in relation to different environments: an appraisal. *Journal of Vegetation Science* 12(2): 153-166.
- Pavlu V., Hejzman M., Pavlu L., Gaisler J. (2003). Effect of rotational and continuous grazing on vegetation of an upland grassland in the Jizerske Hory Mts., Czech Republic. *Folia Geobotanica* 38: 21-34.
- Payraudeau S., Van der Werf H.M.G. (2005). Environmental impact assessment for a farming region: a review of methods *Agriculture, Ecosystems & Environment* 107(1): 1-19.
- Pearson D.L. (1994). Selecting Indicator Taxa for the Quantitative Assessment of Biodiversity. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 345(1311): 75-79.
- Pearson D.L., Cassola F. (1992). World-wide species richness patterns of tiger beetles (Coleoptera: Cicindelidae): indicator taxon for biodiversity and conservation studies. *Conservation Biology* 6(3): 376-391.
- Peco B., de Pablos I., Traba J., Levassor C. (2005). The effect of grazing abandonment on species composition and functional traits: the case of dehesa grasslands. *Basic And Applied Ecology* 6(2): 175-183.
- Peeters A., Maljean J.F., Biala K., Brouckaert V. (2004). Indicators of biodiversity in grasslands: a tool for evaluating the sustainability of livestock-rearing systems. *Fourrages*(No.178): 217-232.
- Pekar S. (1999). Effect of IPM practices and conventional spraying on spider population dynamics in an apple orchard. *Agriculture, Ecosystems & Environment* 73(2): 155-166.
- Pépin D., Menaut P., Desneux L., Cargnelutti B. (1992). Seasonal changes in the use of space by isards (*Rupicapra pyrenaica*) in a protected area, *Ongulés / Ungulates 91*, Paris-Toulouse, Spitz F. *et al.*, eds., S.F.E.P.M.-I.R.G.M.: 327-330.
- Pérés G., Cluzeau D., Curmi P., Hallaire V. (1998). Earthworm activity and soil structure changes due to organic enrichments in vineyard systems. *Biology and Fertility of Soils* 27(4): 417-424.
- Perez-Piqueres A., Edel-Hermann W., Alabouvette C., Steinberg C. (2006). Response of soil microbial communities to compost amendments. *Soil Biology & Biochemistry* 38(3): 460-470.
- Perrow M.R., Jowitt A.J.D. (1995). What future for the Harvest Mouse ? *British Wildlife* 6: 356-365.
- Peters R.D., Sturz A.V., Carter M.R., Sanderson J.B. (2004). Influence of crop rotation and conservation tillage practices on the severity of soil-borne potato diseases in temperate humid agriculture. *Canadian Journal of Soil Science* 84(4): 397-402.
- Petersen H., Jucevica E., Gjelstrup P. (2004). Long-term changes in collembolan communities in grazed and non-grazed abandoned arable fields in Denmark. *Pedobiologia* 48(5-6): 559-573.
- Petersen S., Axelsen J.A., Tybirk K., Aude E., Vestergaard P. (2006). Effects of organic farming on field boundary vegetation in Denmark. *Agriculture, Ecosystems & Environment* 113(1-4): 302-306.
- Pfiffner L., Luka H., Jeanneret P., Schupbach B. (2000). Effects of ecological compensation areas on the carabid fauna. *Agrarforschung* 7(5): 212-217.
- Pearce T.G. (1984). Earthworm populations in soils disturbed by trampling. *Biological conservation* 29(3): 241-252.
- Pilar-Izquierdo M., Gibert P. (2004). Cohabitation entre les cheptels domestiques et la faune sauvage dans la Réserve Nationale des Bauges. *Faune Sauvage* 261: 48-53.
- Pilcher C.D., Obrycki J.J., Rice M.E., Lewis L.C. (1997). Preimaginal development, survival, and field abundance of insect predators on transgenic *Bacillus thuringiensis* corn. *Environmental Entomology* 26: 446-454.
- Piotrowski J.S., Denich T., Klironomos J.N., Graham J.M., Rillig M.C. (2004). The effects of arbuscular mycorrhizas on soil aggregation depend on the interaction between plant and fungal species. *New Phytologist* 164(2): 365-373.
- Pioz M., Loison A., Gibert P., Dubray D., Menaut P., Le Tallec B., Artois M., Gilot-Fromont E. (2007). Transmission of a pestivirus infection in a population of Pyrenean chamois. *Veterinary Microbiology* 119(1): 19-30.
- Piper J.K., Billings D.N., Leite V.J. (2005). Effects of nitrogen fertilizer on the composition of two prairie plant communities. *Community Ecology* 6(1): 93-100.
- Piutti S., Hallet S., Rousseaux S., Philippot L., Soulas G., Martin-Laurent F. (2002). Accelerated mineralisation of atrazine in maize rhizosphere soil. *Biology and Fertility of Soils* 36(6): 434-441.
- Plantureux S. (1996). Biodiversity, type of soil and management intensity of permanent pastures of Plateau Lorrain. *Acta Botanica Gallica* 143(4-5): 339-348.
- Plantureux S., Peeters A., McCracken D. (2005). Biodiversity in intensive grasslands: Effect of management, improvement and challenges. *Agronomy Research* 3(2): 153-164.
- Plassart P., Vincelas M.A., Gangneux C., Mercier A., Barray S., Laval K. (2008). Molecular and functional responses of soil microbial communities under grassland restoration. *Agriculture Ecosystems & Environment* 127(3-4): 286-293.
- Plumb R.T., Browilow R.H. (2005). Pesticides and Birds: A Report on the evidence for changes in farmland bird populations and the proposals for a pesticide tax, Home grown cereals authority, project n°2478, 46, 12: 23 p. http://www.pesticides.gov.uk/pesticides_forum.asp?id=540&link=%2Fuploadedfiles%2FWeb%5FAssets%2FPesticides%5FForum%2FPF115%2Epdf
- Poll C., Thiede A., Wermbter N., Sessitsch A., Kandeler E. (2003). Micro-scale distribution of microorganisms and microbial enzyme activities in a soil with long-term organic amendment. *European Journal of Soil Science* 54(4): 715-724.
- Poorter H., Garnier E. (2007). Ecological significance of inherent variation in relative growth rate and its components. In *Functional Plant Ecology* (2nd Edition), (Pugnaire F.I., Valladares F., eds.), CRC Press, Boca Raton: 67-100.
- Popay I., Field, R. (1996). Grazing animals as weed control agents. *Weed Technology* 10: 217-231.
- Poschlod P., Kiefer S., Tränkle U., Fischer S., Bonn S. (1998). Plant species richness in calcareous grasslands as affected by dispersability in space and time. *Applied Vegetation Science* 1(1): 75-90.
- Poschold P., Bakker J.P., Kahmen S. (2005). Changing land use and its impact on biodiversity. *Basic and Applied Ecology* 6(2): 93-98.

- Postma J., Willemsen-de Klein M., van Elsas J.D. (2000). Effect of the indigenous microflora on the development of root and crown rot caused by *Pythium aphanidermatum* in cucumber grown on rockwool. *Phytopathology* 90(2): 125-133.
- Powell D.F. (1988). The response of non-target predatory arthropods to deltamethrin applied to winter wheat on two alternative dates in autumn. *Aspects of Applied Biology* 17(2): 235-236.
- Powles S., B, Preston, C, Bryan, I, B, Jutsum, A, R, (1997). Herbicide resistance: impact and management *Advances in Agronomy* 58: 57-93.
- Poyry J., Lindgren S., Salminen J., Kuussaari M. (2004). Restoration of butterfly and moth communities in semi-natural grasslands by cattle grazing. *Ecological Applications* 14(6): 1656-1670.
- Poyry J., Lindgren S., Salminen J., Kuussaari M. (2005). Responses of butterfly and moth species to restored cattle grazing in semi-natural grasslands. *Biological Conservation* 122(3): 465-478.
- Poyry J., Luoto M., Paukkunen J., Pykala J., Raatikainen K., Kuussaari M. (2006). Different responses of plants and herbivore insects to a gradient of vegetation height: an indicator of the vertebrate grazing intensity and successional age. *Oikos* 115(3): 401-412.
- Prach K., Pysek P., Smlauer P. (1997). Changes in species traits during succession: a search for pattern. *Oikos* 79: 201-205.
- Prentice H.C., Lonn M., Lager H., Rosen E., Van Der Maarel E. (2000). Changes in allozyme frequencies in *Festuca ovina* populations after a 9-year nutrient/water experiment. *Journal of Ecology* 88(2): 331-347.
- Prentice H.C., Lonn M., Rosquist G., Ihse M., Kindstrom M. (2006). Gene diversity in a fragmented population of *Briza media*: grassland continuity in a landscape context. *Journal of Ecology* 94(1): 87-97.
- Prokopy R.J., Mason J.L., Christie M., Wright S.E. (1995). Arthropod pest and natural enemy abundance under second-level versus first-level integrated pest management practices in apple orchards: a 4-year study. *Agriculture, Ecosystems & Environment* 57(1): 35-47.
- Proulx M., Mazumder A. (1998). Reversal of grazing impact on plant species richness in nutrient-poor vs. nutrient-rich ecosystems. *Ecology* 79: 2581-2592.
- Puricelli E., Tuesca D. (2005). Weed density and diversity under glyphosate-resistant crop sequences. *Crop Protection* 24: 533-542.
- Purtauf T., Roschewitz I., Dauber J., Thies C., Tscharntke T., Wolters V. (2005). Landscape context of organic and conventional farms: Influences on carabid beetle diversity. *Agriculture, Ecosystems & Environment* 108(2): 165-174.
- Purvis G. (1992). A long-term study of the impact of methiocarb molluscicide on carabid populations and case history for interpretation of non-target pesticide effects in the field. *Aspects of Applied Biology*(31): 97-104.
- Purvis G., Abdoulla F. (2002). The influence of cropping rotations and soil cultivation practice on the population ecology of carabids (Coleoptera: Carabidae) in arable land. *Pedobiologia* 46(5): 452-474.
- Purvis G., Bannon J.W. (1992). Non-target effects of repeated methiocarb slug pellet application on carabid beetle (Coleoptera: Carabidae) activity in winter-sown cereals. *Annals of Applied Biology* 121(2): 401-422.
- Putman R.J., Fowler A.D., Tout S. (1991). Patterns of use of ancient grassland by cattle and horses and effects on vegetational composition and structure. *Biological Conservation* 56(3): 329-347.
- Putman R.J., Moore N.P. (1998). Impact of deer in lowland Britain on agriculture, forestry and conservation habitats. *Mammal review* 28(4): 141-164.
- Pykala J. (2000). Mitigating human effects on European biodiversity through traditional animal husbandry. *Conservation Biology* 14(3): 705-712.
- Pykälä J. (2004). Cattle grazing increases plant species richness of most species trait groups in mesic semi-natural grasslands. *Plant Ecology* 175: 217-226.
- Pykälä J., Luoto M., Heikkinen R.K., Kontula T. (2005). Plant species richness and persistence of rare plants in abandoned semi-natural grasslands in northern Europe. *Basic and Applied Ecology* 6: 25-33.
- Pysek P., Jarosik, V, Kropac, Z, Chytry, M, Wild, J, Tichy, L, (2005). Effects of abiotic factors on species richness and cover in Central European weed communities. *Agriculture Ecosystems & Environment* 109: 1-8.
- Pywell R.F., Bullock J.M., Tallowin J.B., Walker K.J., Warman E.A., Masters G. (2007). Enhancing diversity of species-poor grasslands: an experimental assessment of multiple constraints. *Journal of Applied Ecology* 44(1): 81-94.
- Quimby P.C., Bruckart W.L., Deloach C., J., Knutson L., Ralphs M.H. (1991). Biological control of rangeland weeds. In *Noxious Range Weeds*, Westview Press, Boulder: 83-102.
- Quinn M.S., Robinson G.R. (1987). The effects of experimental subdivision on flowering plant diversity in a California annual grassland. *Journal of Ecology* 75: 837-856.
- Raaijmakers J.M., Bonsall R.E., Weller D.M. (1999). Effect of population density of *Pseudomonas fluorescens* on production of 2,4-diacetylphloroglucinol in the rhizosphere of wheat. *Phytopathology* 89(6): 470-475.
- Raaijmakers J.M., Paulitz T.C., Alabouvette C., Steinberg C., Moëgne-Loccoz Y. (2008). The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant and Soil*. DOI 10.1007/s11104-008-9568-6.
- Rabalais N.N., Turner R.E., Scavia D. (2002). Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River. *BioScience* 52(2): 129-142.
- Rajaniemi T.K. (2002). Why does fertilization reduce plant species diversity? Testing three competition-based hypotheses. *Journal of Ecology* 90(2): 316-324.
- Ranjard L., Echairi A., Nowak V., Lejon D.P.H., Nouaim R., Chaussod R. (2006b). Field and microcosm experiments to evaluate the effects of agricultural Cu treatment on the density and genetic structure of microbial communities in two different soils. *Fems Microbiology Ecology* 58(2): 303-315.
- Ranjard L., Lignier L., Chaussod R. (2006a). Cumulative effects of short-term polymetal contamination on soil bacterial community structure. *Applied and Environmental Microbiology* 72(2): 1684-1687.
- Ransome R.D. (1996). The management of feeding areas for greater horseshoe bats ? . *Engl. Nat. Res. Rep.* 174: 1-74.
- Rantalainen M.L., Fritze H., Haimi J., Pennanen T., Setälä H. (2005). Species richness and food web structure of soil decomposer community as affected by the size of habitat fragment and habitat corridors. *Global Change Biology* 11(10): 1614-1627.
- Rantalainen M.L., Haimi J., Fritze H., Setälä H. (2006). Effects of small-scale habitat fragmentation, habitat corridors and mainland dispersal on soil decomposer organisms. *Applied Soil Ecology* 34(2-3): 152-159.

- Raps A., Kehr J., Gugerli P., Moar W.J., Bigler F., Hilbeck A. (2001). Immunological analysis of phloem sap of *Bacillus thuringiensis* corn and the nontarget herbivore *Rhopalosiphum padi* (Homoptera: Aphididae) for the presence of Cry1Ab. *Molecular Ecology* 10: 523-525.
- Rasmussen K.J. (1999). Impact of ploughless soil tillage on yield and soil quality: a Scandinavian review. *Soil & Tillage Research* 53(1): 3-14.
- Ratschker U.M., Roth M. (1997). The spider fauna of agroecosystems - effects of different forms of land use management and intensities on faunal parameters. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 11(1-6): 125-130.
- Raworth D.A., Robertson M.C., Bittman S. (2004). Effects of dairy slurry application on carabid beetles in tall fescue, British Columbia, Canada. *Agriculture, Ecosystems & Environment* 103(3): 527-534.
- Redzic S. (2007). Syntaxonomic diversity as an indicator of ecological diversity - case study Vranica Mts in the Central Bosnia. *Biologia* 62(2): 173-184.
- Reeleder R.D. (2003). Fungal plant pathogens and soil biodiversity. *Canadian Journal of Soil Science* 83(3): 331-336.
- Rehman S.U., Browning H.W., Nigg H.N., Harrison J.M. (2000). Increases in florida red scale populations through pesticidal elimination of *Aphytis holoxanthus* Debach in Florida citrus. *Biological Control* 18(2): 87-93.
- Reich P.B., Walters M.B., Ellsworth D.S. (1992). Leaf life-span in relation to leaf, plant, and stand characteristics among diverse ecosystems. *Ecological Monographs* 62: 365-392.
- Reidsma P., Tekelenburg T., van den Berg M., Alkemade J.R.M. (2006). Impacts of land-use change on biodiversity: an assessment of agricultural biodiversity in the European Union. *Agriculture, Ecosystems & Environment* 114(1): 86-102.
- Reus J., Leendertse P., Bockstaller C., Fomsgaard I., Gutsche V., Lewis K., Nilsson C., Pussemier L., Trevisan M., van der Werf H., Alfarroba F., Blumel S., Isart J., McGrath D., Seppala T. (2002). Comparison and evaluation of eight pesticide environmental risk indicators developed in Europe and recommendations for future use. *Agriculture Ecosystems & Environment* 90(2): 177-187.
- Rew L.J., Wilson P.J., Froud-Williams R.J., Boatman N.D. (1992). Changes in vegetation composition and distribution within set-aside land, *Set-aside*, Cambridge University, 15th-18th september 1992, 50, Clarke J., ed., BCPC: 79-84.
- Reyers B., Jaarsveld A.S.v., McGeoch M.A., James A.N. (1998). National biodiversity risk assessment: a composite multivariate and index approach. *Biodiversity and Conservation* 7(7): 945-965.
- Reynolds H.L., Mittelbach G.G., Darcy-Hall T.L., Houseman G.R., Gross K.L. (2007). No effect of varying soil resource heterogeneity on plant species richness in a low fertility grassland. *Journal of Ecology* 95(4): 723-733.
- Reznik S.Y. (1991). The effects of feeding damage in ragweed *Ambrosia artemisiifolia* (Asteraceae) on populations of *Zygogramma suturalis* (Coleoptera, Chrysomelidae). *Oecologia* 88(2): 204-210.
- Rhett R.G., Simmers J.S., Lee C.R. (1988). *Eisenia fetida* used as a biomonitoring tool to predict the potential bioaccumulation of contaminants from contaminated dredged material. In *Earthworms in waste and environmental management*, (Edwards C.A., Neuhauser E.F., eds.), SPB Academic Publishing, The Hague, The Netherlands: 321-328.
- Rhine E.D., Fuhrmann J.J., Radosevich M. (2003). Microbial community responses to atrazine exposure and nutrient availability: Linking degradation capacity to community structure. *Microbial Ecology* 46(2): 145-160.
- Ricketts T.H., Regetz J., Steffan-Dewenter I., Cunningham S.A., Kremen C., Bogdanski A., Gemmill-Herren B., Greenleaf S.S., Klein A.M., Mayfield M.M., Morandin L.A., Ochieng A., Viana B.F. (2008). Landscape effects on crop pollination services: are there general patterns? *Ecology Letters* 11(5): 499-515.
- Ricklefs R.E., Miller T.E. (2005). *Ecology*, De Boeck University, Paris, 821 p.
- Ricklefs R.E., Schlüter D. (1993). *Species diversity in ecological communities: historical and geographical perspectives*, University of Chicago Press, Chicago, 414 p.
- Riddick E.W., Dively G., Barbosa P. (2000). Season-long abundance of generalist predators in transgenic versus nontransgenic potato fields. *Journal of Entomological Science* 35(4): 349-359.
- Rigamonti I.E., Lozzia G.C. (2002). Phytoseiid mites (Acari: Phytoseiidae) on apple tree and spontaneous flora under different environmental and cultural conditions in Valtellina (Lombardy, Northern Italy). *Bollettino di Zoologia Agraria e di Bachicoltura* 34(1): 53-70.
- Rigby D., Woodhouse P., Young T., Burton M. (2001). Constructing a farm level indicator of sustainable agricultural practice. *Ecological Economics* 39(3): 463-478.
- Riley J. (2001). The indicator explosion: Local needs and international challenges - Preface. *Agriculture Ecosystems & Environment* 87(2): 119-120.
- Ritchie M.E., Tilman D., Knops J.H.M. (1998). Herbivore Effects on Plant and Nitrogen Dynamics in Oak Savanna *Ecology* 79: 165-177.
- Rodriguez A., Mamarot J. (1992). La jachère dans les rotations : étude des risques de salissement des terres, ACTA Point, 2, 39 p.
- Roem W.J., Berendse F. (2000). Soil acidity and nutrient supply ratio as possible factors determining changes in plant species diversity in grassland and heathland communities. *Biological Conservation* 92(2): 151-161.
- Romeis J., Babendreier D., Wackers F.L. (2003). Consumption of snowdrop lectin (*Galanthus nivalis* agglutinin) causes direct effects on adult parasitic wasps. *Oecologia* 134(4): 528-536.
- Romijn C., Luttik R., Canton J.H. (1994). Presentation of a General Algorithm to Include Effect Assessment on Secondary Poisoning in the Derivation of Environmental-Quality Criteria .2. Terrestrial Food-Chains. *Ecotoxicology and Environmental Safety* 27(2): 107-127.
- Rook A.J., Dumont B., Isselein J., Osoro K., WallisDeVries M.F., Parente G., Mills J. (2004). Matching type of livestock to desired biodiversity outcomes in pastures - a review. *Biological Conservation* 119(2): 137-150.
- Rook A.J., Tallowin J.R.B. (2003). Grazing and pasture management for biodiversity benefit. *Animal Research* 52(2): 181-189.
- Roper M.M., Gupta V. (1995). Management-Practices and Soil Biota. *Australian Journal of Soil Research* 33(2): 321-339.
- Ros M., Klammer S., Knapp B., Aichberger K., Insam H. (2006). Long-term effects of compost amendment of soil on functional and structural diversity and microbial activity. *Soil Use and Management* 22(2): 209-218.
- Roschewitz I., Gabriel D., Tschamtké T., Thies C. (2005). The effects of landscape complexity on arable weed species diversity in organic and conventional farming. *Journal of Applied Ecology* 42(5): 873-882.
- Roschewitz I., Hucker M., Tschamtké T., Thies C. (2005). The influence of landscape context and farming practices on parasitism of cereal aphids. *Agriculture, Ecosystems & Environment* 108(3): 218-227.

- Rosi-Marshall E.J., Tank J.L., Royer T.V., Whiles M.R., Evans-White M., Chambers C., Griffiths N.A., Pokelsek J., Stephen M.L. (2007). Toxins in transgenic crop byproducts may affect headwater stream ecosystems. *PNAS* 104(41): 16204-16208.
- Rosnoblet J., Girardin P., Weinzaepflen E., Bockstaller C. (2006). Analysis of 15 years of agriculture sustainability evaluation methods, 9. *ESA Congress*, Warsaw, Poland, 2006/09/04-06, Fotyma M., Kaminska B., eds.: 707-708.
- Rossi S., Hars J., Louquet Y., Masse-Provin N., Pol F., Le Potier M.F. (2006). Gestion d'un réservoir sauvage : la peste porcine du Sanglier (*Sus scrofa*). *Bulletin de l'Académie vétérinaire de France* 159(5): 389-392.
- Rossignol N., Bonis A., Bouzii J.B. (2006). Consequence of grazing pattern and vegetation structure on the spatial variations of net N mineralisation in a wet grassland. *Applied Soil Ecology* 31(1-2): 62-72.
- Roulois V. (2006). Evaluation de l'impact de mesures environnementales sur un territoire ordinaire : méthodologie et premiers résultats, Université de Poitiers: 1-63.
- Roux-Michollet D. (2007). Effet de la désinfection à la vapeur des sols agricoles sur les communautés bactériennes fonctionnelles et leur environnement: résistance, résilience et dynamique de colonisation. PhD thesis, Soil microbiology, University Claude Bernard-Lyon I, Lyon, 125 p.
- Rukayadi Y., Suwanto A., Tjahjono B., Harling R. (2000). Survival and epiphytic fitness of a nonpathogenic mutant of *Xanthomonas campestris* pv. *glycines*. *Applied and Environmental Microbiology* 66(3): 1183-1189.
- Russ J.M., Briffa M., Montgomery W.I. (2003). Seasonal patterns in activity and habitat use by bats (*Pipistrellus* spp. and *Nyctalus leisleri*) in Northern Ireland, determined using a driven transect. *Journal of Zoology* 259(3): 289-299.
- Russell E.P. (1989). Enemies hypothesis: a review of the effect of vegetational diversity on predatory insects and parasitoids. *Environmental Entomology* 18(4): 590-599.
- Russo D., Jones G. (2003). Use of foraging habitats by bats in a Mediterranean area determined by acoustic surveys: conservation implications. *Ecography* 26(2): 197-209.
- Russo D., Jones G., Migliozzi A. (2002). Habitat selection by the Mediterranean horseshoe bat, *Rhinolophus euryale* (Chiroptera : Rhinolophidae) in a rural area of southern Italy and implications for conservation. *Biological Conservation* 107(1): 71-81.
- Rydberg N.T., Milberg P. (2000). A survey of weeds in organic farming in Sweden. *Biological Agriculture & Horticulture* 18(2): 175-185.
- Ryser P., Langenauer R., Gigon A. (1995). Species richness and vegetation structure in a limestone grassland after 15 years management with six biomass removal regimes. *Folia Geobotanica et Phytotaxonomica* 30(2): 157-167.
- Saarinen K., Jantunen J. (2005). Grassland butterfly fauna under traditional animal husbandry: Contrasts in diversity in mown meadows and grazed pastures. *Biodiversity and Conservation* 14(13): 3201-3213.
- Salamolard M., Butet A., Leroux A., Bretagnolle V. (2000). Responses of an avian predator to variations in prey density at a temperate latitude. *Ecology* 81(9): 2428-2441.
- Sall S.N., Masse D., Ndour N.Y.B., Chotte J.L. (2006). Does cropping modify the decomposition function and the diversity of the soil microbial community of tropical fallow soil? *Applied Soil Ecology* 31(3): 211-219.
- Sammul M., Kull K., Tamm A. (2003). Clonal growth in a species-rich grassland: result of a 20-year fertilization experiment. *Folia Geobotanica* 38(1): 1-20.
- Sanvido O., Widmer F., Winzeler M., Streit B., Szerencsi E., Bigler F. (2005). Coexistence de cultures avec ou sans OGM en Suisse. *Revue Suisse d'Agriculture* 37(5): 221-226.
- Sauberer N., Zulka K.P., Abensperg-Traun M., Berg H.M., Bieringer G., Milasowszky N., Moser D., Plutzer C., Pollheimer M., Storch C., Trostl R., Zechmeister H., Grabherr G. (2004). Surrogate taxa for biodiversity in agricultural landscapes of eastern Austria. *Biological Conservation* 117(2): 181-190.
- Sauphanor B., Bouvier J.C., Boisneau C., Rieux R., Simon S., Capowiez Y., Toubon J.F. (2005). Impacts biologiques des systèmes de protection en vergers de pommiers. *Phytoma, la défense des végétaux* 581: 32-36.
- Saxena D., Flores S., Stotzky G. (1999). Transgenic plants - Insecticidal toxin in root exudates from Bt corn. *Nature* 402(6761): 480-480.
- Saxena D., Stotzky G. (2001). *Bacillus thuringiensis* (Bt) toxin released from root exudates and biomass of Bt corn has no apparent effect on earthworms, nematodes, protozoa, bacteria, and fungi in soil. *Soil Biology & Biochemistry* 33(9): 1225-1230.
- Schechtner G. (1993). Efficiency of liming on grasslands. *Bodenkultur* 44(2): 135-152.
- Scheepens P.C., Müller-Schärer H., Kempenaar C. (2001). Opportunities for biological weed control in Europe. *BioControl* 46: 127-138.
- Schellberg J., Mösel B.M., Kühbauch W., Rademacher I.F. (1999). Long-term effects of fertilizer on soil nutrient concentration, yield, forage quality and floristic composition of a hay meadow in the Eifel mountains, Germany. *Grass & Forage Science* 54(3): 195-207.
- Scheu S., Schulz E. (1996). Secondary succession, soil formation and development of a diverse community of oribatids and saprophagous soil macro-invertebrates. *Biodiversity and Conservation* 5(2): 235-250.
- Schlöter M., Bach H.J., Metz S., Sehy U., Munch J.C. (2003). Influence of precision farming on the microbial community structure and functions in nitrogen turnover. *Agriculture Ecosystems & Environment* 98(1-3): 295-304.
- Schmidt M.H., Roschewitz I., Thies C., Tschamtké T. (2005). Differential effects of landscape and management on diversity and density of ground-dwelling farmland spiders. *Journal of Applied Ecology* 42(2): 281-287.
- Schmidt M.H., Tschamtké T. (2005). Landscape context of sheetweb spider (Araneae : Linyphiidae) abundance in cereal fields. *Journal of Biogeography* 32(3): 467-473.
- Schmidt N.M., Olsen H. (2003). The response of small mammal communities to cattle grazing on a coastal meadow. *Polish Journal of Ecology* 51(1): 79-84.
- Schmidt N.M., Olsen H., Bildsoe M., Sluydts V., Leirs H. (2005). Effects of grazing intensity on small mammal population ecology in wet meadows. *Basic and Applied Ecology* 6(1): 57-66.
- Schmutterer H. (1999). Side effects of neem products on insect pathogens and natural enemies of spider mites and insects. In *Azadirachta indica* A. Juss., (Singh R.P., Saxena R.C., eds.), Enfield USA: 147-162.
- Schneider O., Aubertot J.-N., Roger-Estrade J., Doré T. (2003). Analysis and modelling of the amount of oilseed rape residues left at the soil surface after different soil tillage operations, 7. *International Conference on Plant Pathology*, Tours, France, 3-5/12/2003.
- Schnepf E., Crickmore N., Van Rie J., Lereclus D., Baum J., Feitelson J., Zeigler D.R., Dean D.H. (1998). *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiology and Molecular Biology Reviews* 62(3): 775-806.

- Schnitzler A., Muller S. (1998). Towards an ecological basis for the conservation of subalpine heath-grassland on the upper ridges of the Vosges. *Journal of Vegetation Science* 9(3): 317-326.
- Schuld M., Schmuck R. (2000). Effects of thiacloprid, a new chloronicotinyl insecticide, on the egg parasitoid *Trichogramma cacoeciae*. *Ecotoxicology* 9(3): 197-205.
- Schuster D.J., Calderon M. (1986). Interactions of host plant resistant genotypes and beneficial insects in cotton ecosystems. In *Interactions of Plant Resistance and Parasitoids and Predators of Insects* (Boethel D.J., Eikenberry R.D., eds.), Ellis Horwood Limited, Chichester: 84-97.
- Schwab A., Dubois D., Fried P.M., Edwards P.J. (2002). Estimating the biodiversity of hay meadows in north-eastern Switzerland on the basis of vegetation structure. *Agriculture, Ecosystems & Environment* 93(1/3): 197-209.
- Schweiger O., Maelfait J.P., van Wingerden W., Hendrickx F., Billeter R., Speelmans M., Augenstein I., Aukema B., Aviron S., Bailey D., Bukacek R., Burel F., Diekötter T., Dirkens J., Frenzel M., Herzog F., Liira J., Roubalova M., Bugter R. (2005). Quantifying the impact of environmental factors on arthropod communities in agricultural landscapes across organizational levels and spatial scales. *Journal of Applied Ecology* 42(6): 1129-1139.
- Scimone M., Rook A.J., Garel J.P., Sahin N. (2007). Effects of livestock breed and stocking rate on sustainable grazing systems: 3. Effects on vegetation diversity. *Grass & Forage Science* 62(2): 172-184.
- Scopa A., Dumontet S. (2007). Soil solarization: Effects on soil microbiological parameters. *Journal of Plant Nutrition* 30(4-6): 537-547.
- Scursoni J., Forcella, F., Gunsolus, J., Owen, M., Oliver, R., Smeda, R., Vidrine, R. (2006). Weed diversity and soybean yield with glyphosate management along a north-south transect in the United States. *Weed Science* 54: 713-719.
- Sébillotte M., Allain S., Doré T., Meynard J.-M. (1993). La jachère et ses fonctions agronomiques, économiques et environnementales. Diagnostic actuel. *Comptes-Rendus de l'Académie d'Agriculture de France* 79: 105-118.
- Seeber J., Seeber G.U.H., Kossler W., Langel R., Scheu S., Meyer E. (2005). Abundance and trophic structure of macro-decomposers on alpine pastureland (Central Alps, Tyrol): effects of abandonment of pasturing. *Pedobiologia* 49(3): 221-228.
- Seghers D., Top E.M., Reheul D., Bulcke R., Boeckx P., Verstraete W., Siciliano S.D. (2003). Long-term effects of mineral versus organic fertilizers on activity and structure of the methanotrophic community in agricultural soils. *Environmental Microbiology* 5(10): 867-877.
- Seghers D., Verthe K., Reheul D., Bulcke R., Siciliano S.D., Verstraete W., Top E.M. (2003). Effect of long-term herbicide applications on the bacterial community structure and function in an agricultural soil. *Fems Microbiology Ecology* 46(2): 139-146.
- Selinger-Looten R., Muller S. (2001). Restauration d'un couvert prairial sur ancienne culture de maïs : Impact de la fauche et rôle de la banque de graines du sol. *Revue d'Ecologie* 56(1): 3-19.
- Senapati B.K., Lavelle P., Giri S., Pashanasi B., Alegre J., Decaëns T., Jiménez J.J., Albrecht A., Blanchart E., Mahieux M., Rousseaux L., Thomas R.J., Panigrahi P.K., Venkatachalam M. (1999). Soil earthworm technologies for tropical agroecosystems. In *Earthworm management in tropical agroecosystems*, (Lavelle P. et al., eds.), CAB International, Oxford: 199-237.
- Sepp K., Ivask M., Kaasik A., Mikk M., Peepson A. (2005). Soil biota indicators for monitoring the Estonian agri-environmental programme. *Agriculture, Ecosystems & Environment* 108(3): 264-273
- Serra-Cobo J., López-Roig M., Marquès-Bonet T., Lahuerta E. (2000). Rivers as possible landmarks in the orientation flight of *Miniopterus schreibersii*. *Acta Theriologica* 45(3): 347-352.
- Sessitsch A., Howieson J.G., Perret X., Antoun H., Martínez-Romero E. (2002). Advances in Rhizobium research. *Critical Reviews in Plant Sciences* 21(4): 323-378.
- Setälä H., McLean M.A. (2004). Decomposition rate of organic substrates in relation to the species diversity of soil saprophytic fungi. *Oecologia* 139(1): 98-107.
- Shelton M.D., Edwards C.R. (1983). Effects of weeds on the diversity and abundance of insects in soybeans. *Environmental Entomology* 12: 296-298.
- Shi S.J., Bending G.D. (2007). Changes to the structure of *Sphingomonas* spp. communities associated with biodegradation of the herbicide isoproturon in soil. *Fems Microbiology Letters* 269(1): 110-116.
- Shipley B., Vile D., Garnier E. (2006). From plant traits to plant communities: a statistical mechanistic approach to biodiversity. *Science* 314: 812-814.
- Shirai Y. (2006). Laboratory evaluation of effects of transgenic Bt corn pollen on two non-target herbivorous beetles, *Epilachna vigintioctopunctata* (Coccinellidae) and *Galerucella vittaticollis* (Chrysomelidae). *Applied Entomology and Zoology* 41(4): 607-611.
- Shiskoff N., Bruckart W.L. (1993). Evaluation of infection of target and non-target hosts by isolates of the potential biocontrol agent *Puccinia jaceae* that infect *Centaurea* spp. *Phytopathology* 83: 894-898.
- Shrewsbury P.M., Raupp M.J. (2006). Do top-down or bottom-up forces determine *Stephanitis pyrioides* abundance in urban landscapes? *Ecological Applications* 16(1): 262-272.
- Shuster W.D., Edwards C.A. (2003). Interactions between tillage and earthworms in Agroecosystems. In *Soil tillage in Agroecosystems*, (El-Titi A., ed.), Press, CRC, London: 229-260.
- Siepel H. (1996). Biodiversity of soil microarthropods: the filtering of species. *Biodiversity and Conservation* 5(2): 251-260
- Sih A., Kerby J., Bell A., Relyea R. (2004). Response to Schmidt. Pesticides, mortality and population growth rate. *Trends in Ecology & Evolution* 19(9): 460-461.
- Silvertown J., Poulton P., Johnston E., Edwards G., Heard M., Biss P.M. (2006). The Park Grass Experiment 1856-2006: its contribution to ecology. *Journal of Ecology* 94: 801-814.
- Simberloff D. (1998). Flagships, umbrellas, and keystones: Is single-species management passé in the landscape era? *Biological Conservation* 83(3): 247-257.
- Simberloff D., Stiling P. (1996). Risks of species introduced for biological control. *Biological Conservation* 78: 185-192.
- Simon S., Defrance H., Sauphanor B. (2007). Effect of codling moth management on orchard arthropods. *Agriculture, Ecosystems & Environment* 122(3): 340-348.
- Simon S., Sauphanor B., Lauri P.E. (2007). Control of fruit tree pests through manipulation of tree architecture. *Pest Technology* 1(1): 33-37.

- Sims S.R., Holden L.R. (1996). Insect bioassay for determining soil degradation of *Bacillus thuringiensis* subsp *kurstaki* CryIA(b) protein in corn tissue. *Environmental Entomology* 25(3): 659-664.
- Sims S.R., Martin J.W. (1997). Effect of the *Bacillus thuringiensis* insecticidal proteins CryIA(b), CryIA(c), CryIIA, and CryIIIA on *Folsomia candida* and *Xenylla grisea* (Insecta: Collembola). *Pedobiologia* 41(5): 412-416.
- Six J., Elliot E.T., Paustian K. (2000). Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology & Biochemistry* 32: 2099-2103.
- Six J., Frey S.D., Thiet R.K., Batten K.M. (2006). Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Science Society of America Journal* 70(2): 555-569.
- Slotow R., Hamer M. (2000). Biodiversity research in South Africa: comments on current trends and methods. *South African Journal of Science* 96(5): 222-224.
- Smart S.M., Bunce R.G.H., Firbank L.G., Coward P. (2002). Do field boundaries act as refugia for grassland plant species diversity in intensively managed agricultural landscapes in Britain? *Agriculture Ecosystems & Environment* 91(1-3): 73-87.
- Smart S.M., Marrs R.H., Le Duc M.G., Thompson K., Bunce R.G.H., Firbank L.G., Rossall M.J. (2006). Spatial relationships between intensive land cover and residual plant species diversity in temperate farmed landscapes. *Journal of Applied Ecology* 43(6): 1128-1137.
- Smith L.M., Haukos D.A. (2002). Floral diversity in relation to playa wetland area and watershed disturbance. *Conservation Biology* 16(4): 964-974.
- Smith R.S., Shiel R.S., Bardgett R.D., Millward D., Corkhill P., Evans P., Quirk H., Hobbs P.J., Kometa S.T. (2008). Long-term change in vegetation and soil microbial communities during the phased restoration of traditional meadow grassland. *Journal of Applied Ecology* 45: 670-679.
- Smith R.S., Shiel R.S., Bardgett R.D., Millward D., Corkhill P., Rolph G., Hobbs P.J., Peacock S. (2003). Soil microbial community, fertility, vegetation and diversity as targets in the restoration management of a meadow grassland. *Journal of Applied Ecology* 40(1): 51-64.
- Snyder W.E., Clevenger G.M., Eigenbrode S.D. (2004). Intraguild predation and successful invasion by introduced ladybird beetles. *Oecologia* 140(4): 559-565.
- Soberon J., Rodriguez P., Vazquez-Dominguez E. (2000). Implications of the hierarchical structure of biodiversity for the development of ecological indicators of sustainable use. *Ambio* 29(3): 136-142.
- Sosnoskie L., M, Herms, N.P, Cardina, J, (2006). Weed seedbank community composition in a 35-yr-old tillage and rotation experiment. *Weed Science* 54: 263-273.
- Southwood T.R.E. (1988). Tactics, strategies and templets. *Oikos* 52: 3-18.
- Spahillari M., Hammer K., Gladis T., Diederichsen A. (1999). Weeds as part of agrobiodiversity. *Outlook on Agriculture* 28(4): 227-232.
- Spurgeon D.J., Weeks J.M., Van Gestel C.A.M. (2004). A summary of eleven years progress in earthworm ecotoxicology. *Pedobiologia* 47(5-6): 588-606.
- Stanley-Horn D.E., Dively G.P., Hellmich R.L., Mattila H.R., Sears M.K., Rose R., Jesse L.C.H., Losey J.E., Obrycki J.J., Lewis L. (2001). Assessing the impact of Cry1Ab-expressing corn pollen on monarch butterfly larvae in field studies, *Proc. Natl Acad. Sci. USA*, 98: 11931±11936.
- Stark J.D., Banks J.E., Acheampong S. (2004). Estimating susceptibility of biological control agents to pesticides: influence of life history strategies and population structure. *Biological Control* 29(3): 392-398.
- Steffey K.L., Venditti M., Barrido B.R., Felsot A.S. (2004). Effect of *Bacillus thuringiensis* corn on natural enemies of the European corn borer. In *Agricultural Biotechnology: Challenges and Prospects*, Acs Symposium Series, 866, AMER CHEMICAL SOC, Washington: 139-150.
- Steinbauer M.J., Short M.W., Schmidt S. (2006). The influence of architectural and vegetational complexity in eucalypt plantations on communities of native wasp parasitoids: towards silviculture for sustainable pest management. *Forest Ecology and Management* 233(1): 153-164.
- Steinberg C., Edel-Hermann V., Alabouvette C., Lemanceau P. (2007). Soil suppressiveness to plant diseases. In *Modern Soil Microbiology*, (van Elsas J.D. et al., eds.), CRC Press: 455-478.
- Steinmann H.H., Gerowitt B. (1993). Mechanical control of *Galium aparine* in winter wheat. , IV. *IFOAM-conference "Non-chemical Weed Control"*, Dijon (France), Thomas J.-M., ed., IFOAM: 257-261.
- Stotzky G. (2004). Persistence and biological activity in soil of the insecticidal proteins from *Bacillus thuringiensis*, especially from transgenic plants. *Plant and Soil* 266(1-2): 77-89.
- Strijker D. (2005). Marginal land in Europe, causes of decline. *Basic and applied ecology* 6: 99-106.
- Stromberg K.D., Kinkel L.L., Leonard K.J. (2004). Quantifying the effect of bacterial antagonists on the relationship between phyllosphere population sizes of *Xanthomonas translucens* pv. *translucens* and subsequent bacterial leaf streak severity on wheat seedlings. *Biological Control* 29(1): 58-65.
- Strong D.R. (1997). Fear no weevil? Perspectives. *Science* 277: 1058-1059.
- Sturz A.V., Christie B.R. (2003). Beneficial microbial allelopathies in the root zone: the management of soil quality and plant disease with rhizobacteria. *Soil & Tillage Research* 72(2): 107-123.
- Sturz A.V., Ryan D.A.J., Coffin A.D., Matheson B.G., Arsenault W.J., Kimpinski J., Christie B.R. (2004). Stimulating disease suppression in soils: sulphate fertilizers can increase biodiversity and antibiosis ability of root zone bacteria against *Streptomyces scabies*. *Soil Biology & Biochemistry* 36(2): 343-352.
- Suckling D.M., Walker J.T.S., Wearing C.H. (1999). Ecological impact of three pest management systems in New Zealand apple orchards. *Agriculture, Ecosystems & Environment* 73(2): 129-140.
- Sullivan T.P., Sullivan D.S. (2006). Plant and small mammal diversity in orchard versus non-crop habitats. *Agriculture, Ecosystems & Environment* 116(3-4): 235-243.
- Suominen O., Danell K. (2006). Effects of large herbivores on other fauna. In *Large Herbivore Ecology, Ecosystem Dynamics and Conservation* (Danell K. et al., eds.), Cambridge University Press: 383-412.

- Svendsen T.S., Baker G.H. (2002). Survival and growth of *Aporrectodea longa* (Lumbricidae) fed on sheep and cow dung with and without moxidectin residues. *Australian Journal of Agricultural Research* 53(4): 447-451.
- Svendsen T.S., Gronvold J., Holter P., Sommer C. (2003). Field effects of ivermectin and fenbendazole on earthworm populations and the disappearance of dung pats from bolus-treated cattle. *Applied Soil Ecology* 24(3): 207-218.
- Svendsen T.S., Sommer C., Holter P., Gronvold J. (2002). Survival and growth of *Lumbricus terrestris* (Lumbricidae) fed on dung from cattle given sustained-release boluses of ivermectin or fenbendazole. *European Journal of Soil Biology* 38(3-4): 319-322.
- Swanton C.J., Booth B.D., Chandler K., Clements D.R., Shrestha A. (2006). Management in a modified no-tillage corn-soybean-wheat rotation influences weed population and community dynamics. *Weed Science* 54(1): 47-58.
- Swengel A.B. (2001). A literature review of insect responses to fire, compared to other conservation managements of open habitat. *Biodiversity and Conservation* 10(7): 1141-1169.
- Swift M.J., Heal O.W., J.M. A. (1979). Decomposition in terrestrial ecosystems, Studies in Ecology, Vol. 5. , Blackwell Scientific Publications, Oxford, U.K., 384 p.
- Swift M.J., Izac A.M.N., van Noordwijk M. (2004). Biodiversity and ecosystem services in agricultural landscapes - are we asking the right questions? *Agriculture, Ecosystems & Environment* 104(1): 113-134.
- Tallowin J.R.B. (1996). Effects of inorganic fertilizers on flower-rich hay meadows: a review using a case study on the Somerset Levels, UK. *Grasslands and Forage Abstracts* 66(4): 147-152.
- Tallowin J.R.B., Kirkham F.W., Wilkins R.J., Smith R.E.N., Thomas G.H., Mountford O., Lakhani K.H. (1994). The effects of inorganic fertilizers in flower-rich hay meadows on the Somerset Levels, MAFF/DoE/English Nature Tadham Moor Project 1986-93, Institute of Grassland and Environmental Research, North Wyke, Devon.
- Tallowin J.R.B., Rook A.J., Rutter S.M. (2005). Impact of grazing management on biodiversity of grasslands. *Animal Science* 81: 193-198.
- Tallowin J.R.B., Smith R.E.N., Goodyear J., Vickery J.A. (2005). Spatial and structural uniformity of lowland agricultural grassland in England: a context for low biodiversity. *Grass & Forage Science* 60(3): 225-236.
- Tarmi S., Tuuri H., Helenius J. (2002). Plant communities of field boundaries in Finnish farmland. *Agricultural and Food Science in Finland* 11: 121-135.
- Tatoni T., Magnin F., Bonin G., Vaudour J. (1994). Secondary successions on abandoned cultivation terraces in calcareous Provence. I - Vegetation and soil. *Acta Oecologica* 15: 431-447.
- ten Berge H.F.M., Burgers S.L.G.E., Schröder E.J., Hofstad E.J. (2002). 'Partial balance'-regression models for $N_{min,H}$. In A review of potential indicators for nitrate loss from cropping and farming systems in the Netherlands, (ten Berge H.F.M., ed.), Plant Research International B.V., Wageningen 25-60.
- ter Braak C., J.F., Wiertz, J. (1994). On the statistical analysis of vegetation change : a wetland affected by water extraction and soil acidification. *Journal of vegetation Soil Science* 5: 361-374.
- Tew T.E., Macdonald D.W., Rands M.R.W. (1992). Herbicide Application Affects Microhabitat Use by Arable Wood Mice (*Apodemus sylvaticus*). *Journal of Applied Ecology* 29(2): 532-539.
- Thies C., Roschewitz I., Tschamtkke T. (2005). The landscape context of cereal aphid-parasitoid interactions. *Proceedings of the Royal Society B-Biological Sciences* 272(1559): 203-210.
- Thirumurthi S. (1995). Insecticidal effects on predator populations in tobacco fields. *Insect Environment* 1(2): 10-11.
- Thomas J.A. (2005). Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Philosophical Transactions of the Royal Society B-Biological Sciences* 360(1454): 339-357.
- Thomas M.B., Wratten S.D., Sotherton N.W. (1992). Creation of 'island' habitats in farmland to manipulate populations of beneficial arthropods: predator densities and species composition. *Journal of Applied Ecology* 29(2): 524-531.
- Thompson I.P., Bailey M.J., Fenlon J.S., Fermor T.R., Lilley A.K., Lynch J.M., McCormack P.J., McQuilken M.P., Purdy K.J., Rainey P.B., Whipps J.M. (1993). Quantitative and qualitative seasonal changes in the microbial community from the phyllosphere of sugar beet (*Beta vulgaris*). *Plant and Soil* 150: 177-191.
- Thomson J.R., Fleishman E., Mac Nally R., Dobkin D.S. (2007). Comparison of predictor sets for species richness and the number of rare species of butterflies and birds. *Journal of Biogeography* 34(1): 90-101.
- Thomson L.J., Hoffmann A.A. (2006). Field validation of laboratory-derived IOBC toxicity ratings for natural enemies in commercial vineyards. *Biological Control* 39(3): 507-515.
- Tichit M., Doyen L., Lemel J.Y., Renault O., Durant D. (2007). A co-viability model of grazing and bird community management in farmland. *Ecological Modelling* 206(3-4): 277-293.
- Tichit M., Durant D., Kernéis E. (2005). The role of grazing in creating suitable sward structures for breeding waders in agricultural landscapes. *Livestock Production Science* 96(1): 119-128.
- Tichit M., Renault O., Potter T. (2005). Grazing regime as a tool to assess positive side effects of livestock farming systems on wading birds. *Livestock Production Science* 96(1): 109-117.
- Tietjen W.J., Cady A.B. (2007). Sublethal exposure to a neurotoxic pesticide affects activity rhythms and patterns of four spider species. *Journal of Arachnology* 35(2): 396-406.
- Tilman D. (1990). Constraints and tradeoffs: toward a predictive theory of competition and succession. *Oikos* 58: 3-15.
- Tilman D., Fargione J., Wolff B., D'Antonio C., Dobson A., Howarth R., Schindler D., Schlesinger W.H., Simberloff D., Swackhamer D. (2001). Forecasting agriculturally driven global environmental change. *Science* 292: 281-284.
- Tilman D., Reich P.B., Knops J., Wedin D., Mielke T., Lehman C. (2001). Diversity and productivity in a long-term grassland experiment. *Science* 294:843-845. *Science* 294(5543): 843-845.
- Timms-Wilson T.M., Kilshaw K., Bailey M.J. (2004). Risk assessment for engineered bacteria used in biocontrol of fungal disease in agricultural crops. *Plant and Soil* 266(1-2): 57-67.
- Tobor-Kaplon M.A., Bloem J., Romkens P., de Ruiter P.C. (2005). Functional stability of microbial communities in contaminated soils. *Oikos* 111(1): 119-129.
- Tonhasca A. (1993). Effects of agroecosystem diversification on natural enemies of soybean herbivores. *Entomologia Experimentalis et Applicata* 69(1): 83-90.

- Torres J.B., Ruberson J.R. (2007). Abundance and diversity of ground-dwelling arthropods of pest management importance in commercial Bt and non-Bt cotton fields. *Annals of Applied Biology* 150(1): 27-39.
- Tratalos J., Fuller R.A., Evans K.L., Davies R.G., Newson S.E., Greenwood J.J.D., Gaston K.J. (2007). Bird densities are associated with household densities. *Global Change Biology* 13(8): 1685-1695.
- Triplet P., Durant J., Bacquet S. (1997). Reproduction du vanneau huppé *Vanellus vanellus* et pratiques agricoles : caractéristiques des sites utilisés en plaine maritime picarde. *Alauda* 65: 121-130.
- Tscharntke T., Brandl R. (2004). Plant-insect interactions in fragmented landscapes. *Annual Review of Entomology* 49(1): 405-430.
- Tscharntke T., Greiler H.J. (1995). Insect Communities, Grasses, and Grasslands. *Annual Review of Entomology* 40: 535-558.
- Tscharntke T., Klein A.M., Kruess A., Steffan-Dewenter I., Thies C. (2005). Landscape perspectives on agricultural intensification and biodiversity - ecosystem service management. *Ecology Letters* 8(8): 857-874.
- Tscharntke T., Rand T.A., Bianchi F. (2005). The landscape context of trophic interactions: insect spillover across the crop-noncrop interface. *Annales Zoologici Fennici* 42(4): 421-432.
- Tscharntke T., Steffan Dewenter I., Kruess A., Thies C. (2002). Contribution of small habitat fragments to conservation of insect communities of grassland-cropland landscapes. *Ecological Applications* 12(2): 354-363.
- Tucker G.M. (1997). Priorities for bird conservation in Europe : the importance of farmed landscape. In *Farming and birds in Europe : The Common Agricultural Policy and its implications for birds conservation.*, (Pain D., Pienkowski M.W., eds.), Academic Press, San Diego, CA: 79-116.
- Tuovinen T. (1992). Predatory mites in Finnish apple orchards. *Acta Phytopathologica et Entomologica Hungarica* 27(1-4): 609-613.
- Turner M.G. (2005). Landscape ecology in North America: Past, present, and future. *Ecology* 86(8): 1967-1974.
- Tuttle M.D. (1976). Population Ecology of the Gray Bat (*Myotis grisescens*): Factors Influencing Growth and Survival of Newly Volant Young. *Ecology* 57(3): 587-595.
- Unal G., Jepson P.C. (1991). The toxicity of aphicide residues to beneficial invertebrates in cereal crops. *Annals of Applied Biology* 118(3): 493-502.
- UNEP (2003). Monitoring and indicators : designing national-level monitoring programmes and indicators, Conventional Biological Diversity, UNEP/CBD/SBSTTA/9/10: 45 p. <http://www.cbd.int/2010-target/documents.shtml>
- van Bruggen A.H.C., Semenov A.M. (2000). In search of biological indicators for soil health and disease suppression. *Applied Soil Ecology* 15(1): 13-24.
- van Bruggen A.H.C., Semenov A.M., van Diepeningen A.D., de Vos O.J., Blok W.J. (2006). Relation between soil health, wave-like fluctuations in microbial populations, and soil-borne plant disease management. *European Journal of Plant Pathology* 115(1): 105-122.
- van Buskirk J., Willi Y. (2004). Enhancement of farmland biodiversity within set-aside land. *Conservation Biology* 18(4): 987-994.
- van Buskirk J., Willi Y. (2005). Meta-analysis of farmland biodiversity within set-aside land: reply to Kleijn and Baldi. *Conservation Biology* 19(3): 967-968
- van der Heijden M.G.A., Scheublin T.R., Brader A. (2004). Taxonomic and functional diversity in arbuscular mycorrhizal fungi - is there any relationship? *New Phytologist* 164(2): 201-204.
- van der Putten W.H., Mortimer S.R., Hedlund K., Van Dijk C., Brown V.K., Leps J., Rodriguez-Barrueco C., Roy J., FDiaz Len T.A., Gormsen D., Korthals G.W., Lavorel S., Santa Regina I., Smilauer P. (2000). Plant species diversity as a driver of early succession in abandoned fields: a multi-site approach. *Oecologia* 124(1): 91-99.
- van der Werf H.M.G., Petit J. (2002). Evaluation of the environmental impact of agriculture at the farm level: a comparison and analysis of 12 indicator-based methods. *Agriculture Ecosystems & Environment* 93(1-3): 131-145.
- van Elsas J.D., Costa R. (2007). Molecular assessment of soil microbial communities with potential for plant disease suppression. In *Biotechnology and plant disease management*, (Punja Z.K. et al., eds.), CAB International, Wallingford UK: 498-517.
- van Elsas J.D., Duarte G.F., Rosado A.S., Smalla K. (1998). Microbiological and molecular biological methods for monitoring microbial inoculants and their effects in the soil environment. *Journal of Microbiological Methods* 32(2): 133-154.
- van Elsas J.D., Garbeva P., Salles J. (2002). Effects of agronomical measures on the microbial diversity of soils as related to the suppression of soil-borne plant pathogens. *Biodegradation* 13(1): 29-40.
- van Elsen T. (2000). Species diversity as a task for organic agriculture in Europe. *Agriculture, Ecosystems & Environment* 77(1/2): 101-109.
- Van Lenteren J.C. (1993). Parasites and predators play a paramount role in insect pest management. In *Pest Management: Biologically Based Technologies* (Lumsden R.D., Vaughn J.L., eds.), Beltsville Agricultural Research Center, Beltsville, Maryland: 68-81.
- van Loon L.C., Rep M., Pieterse C.M.J. (2006). Significance of Inducible Defense-related Proteins in Infected Plants. *Annual Review of Phytopathology* 44(1): 135-162.
- Van Wieren S.E. (1998). Effects of large herbivores upon the animal community. In *Grazing and conservation management*, (Wallis De Vries M.F. et al., eds.), Kluwer Academic Publisher, Dordrecht: 185-214.
- Vellend M. (2005). Species diversity and genetic diversity: parallel processes and correlated patterns. *American Naturalist* 166(2): 199-215.
- Vellend M., Geber M.A. (2005). Connections between species diversity and genetic diversity. *Ecology Letters* 8: 767-781.
- Verboom B., Huitema H. (1997). The importance of linear landscape elements for the pipistrelle *Pipistrellus pipistrellus* and the serotine bat *Eptesicus serotinus*. *Landscape Ecology* 12(2): 117-125.
- Verboom B., Spoelstra K. (1999). Effects of food abundance and wind on the use of tree lines by an insectivorous bat, *Pipistrellus pipistrellus*. *Canadian Journal of Zoology* 77(9): 1393-1401.
- Vercesi M.L., Krogh P.H., Holmstrup M. (2006). Can *Bacillus thuringiensis* (Bt) corn residues and Bt-corn plants affect life-history traits in the earthworm *Aporrectodea caliginosa*? *Applied Soil Ecology* 32(2): 180-187.
- Verheyden H. (2005). Relation du Cerf et du milieu en Pyrénées Centrales. Rapport final, INRA-CEFS - FDC 31 - ONCFS-CNRA Faune de Montagne - CRPF, Toulouse.
- Verma M., Brar S.K., Tyagi R.D., Surampalli R.Y., Valero J.R. (2007). Antagonistic fungi, *Trichoderma* spp.: Panoply of biological control. *Biochemical Engineering Journal* 37(1): 1-20.

- Vesk P.A., Leishman M.R., Westoby M. (2004). Simple traits do not predict grazing response in Australian dry shrublands and woodlands. *Journal of Applied Ecology* 41(1): 22-31.
- Vessey J.K., Pawlowski K., Bergman B. (2005). Root-based N₂-fixing symbioses: Legumes, actinorhizal plants, Parasponia sp and cycads. *Plant and Soil* 274(1-2): 51-78.
- Vickery J.A., Tallwin J.R., Feber R.E., Asteraki E.J., Atkinson P.W., Fuller R.J., Brown V.K. (2001). The management of lowland neutral grasslands in Britain: effects of agricultural practices on birds and their food resources. *Journal of Applied Ecology* 38(3): 647-664.
- Viebahn M., Glandorf D.C.M., Ouwens T.W.M., Smit E., Leeflang P., Wernars K., Thomashow L.S., van Loon L.C., Bakker P. (2003). Repeated introduction of genetically modified *Pseudomonas putida* WCS358r without intensified effects on the indigenous microflora of field-grown wheat. *Applied and Environmental Microbiology* 69(6): 3110-3118.
- Vigl J., Boscheri S., Mantinger H. (1985). Effect of various insecticides and acaricides on predatory mites. *Obstbau Weinbau* 22(4): 108-112.
- Vile D., Shipley B., Garnier E. (2006). Ecosystem productivity relates to species' potential relative growth rate: a field test and a conceptual framework. *Ecology Letters* 9: 1061-1067.
- Violle C., Navas M.L., Vile D., Kazakou E., Fortunel C., Hummel I., Garnier E. (2007). Let the concept of trait be functional! *Oikos* 116: 882-892.
- Vitousek P.M., D'Antonio C.M., Loope L., Rejmanek M., Westbrooks R. (1997). Introduced species: a significant component of human-caused global change. *New Zealand Journal of Ecology* 21: 1-16.
- Vlasenko N.G., Shtundyuk D.A. (1994). The effect of pesticides on the community of carabids in spring rape fields. *Agrokhimiya*(2): 89-94.
- Vogt H. (1992). Acaricide tests in apple orchards with special regard to their effects on beneficials and on the apple rust mite. *Acta Phytopathologica et Entomologica Hungarica* 27(1-4): 659-667.
- Vogt H., Weigel A., Wyss E. (1998). Aspects of indirect plant protection strategies in orchards: are flowering strips an adequate measure to control apple aphids?, 6. *European Congress of Entomology*, České Budejovice, Rép. Tchèque, August 23-29 1998, Brunnhofer V., Soldan T., eds., Institute of Entomology, Academy of the Czech republic and University of South Bohemia: 625-626.
- Volkl W., Stechmann D.H. (1998). Parasitism of the black bean aphid (*Aphis fabae*) by *Lysiphlebus fabarum* (Hym., Aphidiidae): the influence of host plant and habitat. *Journal of Applied Entomology* 122(5): 201-206.
- Volkmar C., Kreuter T. (2006). Biodiversity of spiders (Araneae) and carabid beetles (Carabidae) on fields in Saxony. *Mitteilungen der Deutschen Gesellschaft für allgemeine und angewandte Entomologie* 15: 97-102.
- Volkmar C., Lubke-Al Hussein M., Jany D., Hunold I., Richter L., Kreuter T., Wetzel T. (2003). Ecological studies on epigeous arthropod populations of transgenic sugar beet at Friemar (Thuringia, Germany). *Agriculture, Ecosystems & Environment* 95(1): 37-47.
- von Tiedemann A. (2002). The suppressiveness potential of plants - An approach for "increasing the richness of the soil". *Berichte Über Landwirtschaft*: 57-66.
- Wagner H.H., Wildi O., Ewald K.C. (2000). Additive partitioning of plant species diversity in an agricultural mosaic landscape. *Landscape Ecology* 15: 219-227.
- Waldhardt R. (2003). Biodiversity and landscape - summary, conclusions and perspectives. *Agriculture, Ecosystems & Environment* 98(1/3): 305-309.
- Walker K.J., Stevens P.A., Stevens D.P., Mountford J.O., Manchester S.J., Pywell R.F. (2004). The restoration and re-creation of species-rich lowland grassland on land formerly managed for intensive agriculture in the UK. *Biological Conservation* 119(1): 1-19.
- Walker S., Wilson J.B., Lee W.G. (2005). Does fluctuating resource availability increase invasibility? Evidence from field experiments in New Zealand short tussock grassland. *Biological Invasions* 7(2): 195-211.
- Wall D.H., Adams G.A., Parsons A.N. (2001). Soil Biodiversity. In *Global biodiversity in a changing environment. Scenarios for the 21st Century*, (Chapin F.S. et al., eds.), Springer Verlag, New York: 47-82.
- Wall D.H., Moore J.C. (1999). Interactions underground - Soil biodiversity, mutualism, and ecosystem processes. *BioScience* 49(2): 109-117.
- Wallis de Vries M.F. (1998). Large herbivores as key factors for nature conservation. In *Grazing and conservation management*, (Wallis de Vries M.F. et al., eds.), Kluwer Academic Publishers, Dordrecht: 1-20.
- Wallis de Vries M.F., Raemakers I. (2001). Does extensive grazing benefit butterflies in coastal dunes? *Restoration Ecology* 9(2): 179-188.
- Walsh A.L., Harris S. (1996). Foraging habitat preferences of vespertilionid bats in Britain. *Journal of Applied Ecology* 33(3): 508-518.
- Wamelink G.W.W., de Jong J.J., van Dobben H.F., van Wijk M.N. (2005). Additional costs of nature management caused by deposition. *Ecological Economics* 52(4): 437-451.
- Wang B., Qiu Y.L. (2006). Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 16(5): 299-363.
- Wang Q.S., Shen D.A., Ma Z.Q. (1993). Effects of different pesticides on populations of main pest insects and natural enemies at the seedling stage in soyabean fields. *Entomological Knowledge* 30(6): 333-335.
- Wardhaugh K.G., Mahon R.J. (1991). Avermectin residues in sheep and cattle dung and their effects on dung-beetle (Coleoptera: Scarabaeidae) colonization and dung burial. *Bulletin of Entomological Research* 81(3): 333-339.
- Wardhaugh K.G., Ridsdill-Smith T.J. (1998). Antiparasitic drugs, the livestock industry and dung beetles - cause for concern? *Australian Veterinary Journal* 76(4): 259-261.
- Wardle D.A. (2006). The influence of biotic interactions on soil biodiversity. *Ecology Letters* 9(7): 870-886.
- Wardle D.A., Bardgett R.D., Klironomos J.N., Setälä H., Van der Putten W.H., Wall D.H. (2004). Ecological linkage between aboveground and belowground biota. *Science* 304: 1634-1637.
- Warren J., Paul A.G. (2006). A comparison of invertebrate populations and their relationship with rates of faecal decomposition in organic and conventionally managed pastures. *Biological Agriculture & Horticulture* 24(1): 61-71.
- Wartelle R. (2002). Impact de la Jachère Environnement Faune Sauvage sur la biodiversité : le volet arthropodes. *Faune Sauvage* 256: 74-77.
- Warwick S., I, Légère A., Simard M.J., James T. (2007). Do escaped transgenes persist in nature? The case of an herbicide resistance transgene in a weedy *Brassica rapa* population. *Molecular Ecology* 17(5): 1387-1395.
- Watson A.K., Wymore L.A. (1992). L'approche classique dans la lutte biologique dirigée contre les mauvaises herbes. TEC & DOC ed. In *La lutte biologique*, (Morin G., ed.), 17, Lavoisier: 343-360.

- Way M.J., Cammell M.E. (1981). Effects of weeds and weed control on invertebrate pest ecology. *In Proc. Univ. York and Br. Ecol. Soc. and Feder. Br. Plant Pathol.*, Pitman, London: 443-458.
- Weber D., Hintermann U., Zangger A. (2004). Scale and trends in species richness: considerations for monitoring biological diversity for political purposes. *Global Ecology and Biogeography* 13(2): 97-104.
- Weibull A.C., Ostman O. (2003). Species composition in agroecosystems: The effect of landscape, habitat, and farm management. *Basic and Applied Ecology* 4(4): 349-361.
- Weibull A.-C., Bengtsson J., Nohlgren E. (2000). Diversity of butterflies in the agricultural landscape: the role of farming system and landscape heterogeneity. *Ecography* 23(6): 743-750.
- Weibull A.-C., Östman Ö., Granqvist Å. (2003). Species richness in agroecosystems: the effect of landscape, habitat and farm management. *Biodiversity and Conservation* 12(7): 1335-1355.
- Weier E., Keddy P. (1999). Ecological assembly rules. Perspectives, advances, retreats, Cambridge University Press, Cambridge, 418 p.
- Weller D.M., Raaijmakers J.M., Gardener B.B.M., Thomashow L.S. (2002). Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annual Review of Phytopathology* 40: 309-348.
- Wellstein C., Otte A., Waldhardt R. (2007). Impact of site and management on the diversity of central European mesic grassland. *Agriculture, Ecosystems & Environment* 122(2): 203-210
- Westoby M., Falster D.S., Moles A.T., Vesk P.A., Wright I.J. (2002). Plant ecological strategies: Some leading dimensions of variation between species. *Annual Review of Ecology and Systematics* 33: 125-159.
- Whalen J.K., Hamel C. (2004). Effects of key soil organisms on nutrients dynamics in temperate agroecosystems. *Journal of Crop Improvement* 11(1/2): 175-207
- Whalen J.K., Parmelee R.W., Edwards C.A. (1998). Population dynamics of earthworm communities in corn agroecosystems receiving organic or inorganic fertilizer amendments. *Biology and Fertility of Soils* 27(4): 400-407.
- Whalon M.E., Croft B.A. (1986). Immigration and Colonization of Portable Apple-Trees by Arthropod Pests and Their Natural Enemies. *Crop Protection* 5(6): 376-384.
- Wheatley G.A., Hardman J.A. (1964). Insecticides and chlorinated hydrocarbons and organic phosphorus compounds and residues in soil and water, carrots and earthworms. *Rept. Natl. Veg. Res. Sta.* A5: 63-65.
- Wheeler B.D., Shaw S.C. (1991). Above-ground crop mass and species richness of the principal types of herbaceous rich-fen vegetation of lowland England and Wales. *Journal of Ecology* 79(2): 285-301.
- White T.A., Barker D.J., Kenneth J.M. (2004). Vegetation diversity, growth, quality and decomposition in managed grasslands. *Agriculture, Ecosystems & Environment* 101(1): 73-84.
- Whittaker R.H. (1972). Evolution and measurement of species diversity. *Taxon* 21: 213-251.
- Willems J.H., van Nieuwstadt M.G.L. (1996). Long-term after effects of fertilization on above-ground phytomass and species diversity in calcareous grassland. *Journal of Vegetation Science* 7(2): 177-184.
- Williams B.L., Grayston S.J., Reid E.J. (2000). Influence of synthetic sheep urine on the microbial biomass, activity and community structure in two pastures in the Scottish uplands. *Plant and Soil* 225(1-2): 175-185.
- Willis K.J., Whittaker R.J. (2002). Species diversity - Scale matters. *Science* 295: 1245-1246.
- Wilson E.O. (1992). *The Diversity of Life*, Harvard University Press, Harvard, 432 p.
- Wilson S.D., Shay J.M. (1990). Competition, fire, and nutrients in a mixed-grass prairie. *Ecology* 71(5): 1959-1967.
- Winstone L., Iles D.R., Kendall D.A. (1996). Effects of rotation and cultivation on polyphagous predators in conventional and integrated farming systems. *Aspects of Applied Biology* 47: 111-118.
- With K.A., Pavuk D.M., Worchuck J.L., Oates R.K., Fisher J.L. (2002). Threshold effects of landscape structure on biological control in agroecosystems. *Ecological Applications* 12(1): 52-65.
- Witsack W., Engler I., Schneider K., Schnitter P.H. (1997). Development of zoocenotic structures in nine-year-old agricultural and pastoral fallows. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 11(1/6): 95-98.
- Wolfe M.S. (1985). The Current Status and Prospects of Multiline Cultivars and Variety Mixtures for Disease Resistance. *Annual Review of Phytopathology* 23: 251-273.
- Wolfe M.S. (2000). Crop strength through diversity. *Nature* 406(6797): 681-682.
- Wolsing M., Prieme A. (2004). Observation of high seasonal variation in community structure of denitrifying bacteria in arable soil receiving artificial fertilizer and cattle manure by determining T-RFLP of nir gene fragments. *FEMS Microbiology Ecology* 48(2): 261-271.
- Wolters V. (2001). Biodiversity of soil animals and its function. *European Journal of Soil Biology* 37(4): 221-227.
- Wolters V., Ekschmitt K. (1997). Gasteropods, isopods, diplopods and chilopods: neglected groups of the decomposer food web. *In Fauna in soil ecosystems: recycling, processes, nutrient fluxes and agricultural production*, (Benckiser G., ed.), Marcel Dekker, NewYork - Base - Hong Kong: 265-306.
- Woodcock B.A., Pywell R.F., Roy D.B., Rose R.J., Bell D. (2005). Grazing management of calcareous grasslands and its implications for the conservation of beetle communities. *Biological Conservation* 125(2): 193-202.
- Woodhouse S.P., Good J.E.G., Lovett A.A., Fuller R.J., Dolman P.M. (2005). Effects of land-use and agricultural management on birds of marginal farmland: a case study in the Llyn peninsula, Wales. *Agriculture, Ecosystems & Environment* 107(4): 331-340.
- Woodward F.I., Diament A.D. (1991). Functional approaches to predicting the ecological effects of global change. *Functional Ecology* 5: 202-212.
- Wright I.J., Reich P.B., Westoby M., Ackerly D.D., Baruch Z., Bongers F., Cavender-Bares J., Chapin F.S.I., Cornelissen J.H.C., Diemer M., Flexas J., Garnier E., Groom P.K., Gulias J., Hikosaka K., Lamont B.B., Lee T., Lee W., Lusk C., Midgley J.J., Navas M.-L., Niinemets Ü., Oleksyn J., Osada N., Poorter H., Prior L., Pyankov V.I., Roumet C., Thomas S.C., Tjoelker M.G., Veneklaas E., Villar R. (2004). The worldwide leaf economics spectrum. *Nature* 428: 821-827.
- Wyss E. (1995). The effects of weed strips on aphids and aphidophagous predators in an apple orchard. *Entomologia Experimentalis et Applicata* 75(1): 43-49.
- Yamazaki K., Sugiura S., Kawamura K. (2003). Ground beetles (Coleoptera : Carabidae) and other insect predators overwintering in arable and fallow fields in central Japan. *Applied Entomology and Zoology* 38(4): 449-459.

- Yardim E.N., Edwards C.A. (1998). The influence of chemical management of pests, diseases and weeds on pest and predatory arthropods associated with tomatoes. *Agriculture, Ecosystems & Environment* 70(1): 31-48.
- Yeates G.W., Bongers T. (1999). Nematode diversity in agroecosystems. *Agriculture, Ecosystems & Environment* 74(1-3): 113-135.
- Yeates G.W., Skipp R.A., Gray R.A.J., Chen L.Y., Waghorn T.S. (2007). Impact on soil fauna of sheep faeces containing a range of parasite control agents. *Applied Soil Ecology* 35(2): 380-389.
- Zablotowicz R.M., Krutz L.J., Reddy K.N., Weaver M.A., Koger C.H., Locke M.A. (2007). Rapid development of enhanced atrazine degradation in a dundee silt loam soil under continuous corn and in rotation with cotton. *Journal of Agricultural and Food Chemistry* 55(3): 852-859.
- Zangerl A.R., McKenna D., Wraight C.L., Carroll M., Ficarello P., Warner R., Berenbaum M.R. (2001). Effects of exposure to event 176 *Bacillus thuringiensis* corn pollen on monarch and black swallowtail caterpillars under field conditions, *Proc. Natl Acad. Sci. USA*, 98: 11908-11912.
- Zechmeister H.G., Schmitzberger I., Peterseil J., Wrba T. (2003). The influence of land-use practices and economics on plant species richness in meadows. *Biological Conservation* 114(2): 165-177.
- Zhang N.X., Deng X., Chen J.F. (1986). Introduction and application of *Typhlodromus occidentalis* Nesbitt. *Plant Protection* 12(2): 17-18.
- Zhang Q.C., Wang G.H., Yao H.Y. (2007). Phospholipid fatty acid patterns of microbial communities in paddy soil under different fertilizer treatments. *Journal of Environmental Sciences-China* 19(1): 55-59.
- Zielezny Y., Groeneweg J., Vereecken H., Tappe W. (2006). Impact of sulfadiazine and chlorotetracycline on soil bacterial community structure and respiratory activity. *Soil Biology & Biochemistry* 38(8): 2372-2380.
- Zobel M. (1997). The relative role of species pools in determining plant species richness: an alternative explanation of species coexistence? *Trends in Ecology & Evolution* 12: 266-269.
- Zoebelein G. (1988). Long-term field studies about pesticide effects on ladybird beetles (Coleoptera: Coccinellidae). *Entomologia Generalis* 13(3-4): 175-187.
- Zwahlen C., Hilbeck A., Howald R., Nentwig W. (2003). Effects of transgenic Bt corn litter on the earthworm *Lumbricus terrestris*. *Molecular Ecology* 12(4): 1077-1086.