

Références bibliographiques citées dans le chapitre 2

- Abawi G.S., Widmer T.L. (2000). Impact of soil health management practices on soilborne pathogens, nematodes and root diseases of vegetable crops. *Applied Soil Ecology* 15: 37-47.
- Abrams P.A. (1993). Effect of increased productivity on the abundances of trophic levels. *American Naturalist* 141(3): 351-371.
- ACTA (2008). Index Phytosanitaire 44, ACTA, Paris, 844 p.
- Adnoy T., Haug A., Sorheim O., Thomassen M.S., Varszegi Z., Eik L.O. (2005). Grazing on mountain pastures - does it affect meat quality in lambs? *Livestock Production Science* 94: 25-31.
- Aebi A., Schonrogge K., Melika G., Quacchia A., Alma A., Stone G.N. (2007). Native and introduced parasitoids attacking the invasive chestnut gall wasp *Dryocosmus kuriphilus*. *Bulletin OEPP/EPPO Bulletin* 37(1): 166-171.
- Afun J.V.K., Johnson D.E., Russell-Smith A. (1999). Weeds and natural enemy regulation of insect pests in upland rice; a case study from West Africa. *Bulletin of Entomological Research* 89(5): 391-402.
- Agrama H.A. (2005). Application of molecular markers in breeding for nitrogen use efficiency. *Journal of Crop Improvement* 15(2): 175-211.
- Agreil C., Fritz H., Meuret M. (2005). Maintenance of daily intake through bite mass diversity adjustment in sheep grazing on heterogeneous and variable vegetation. *Applied Animal Behaviour Science* 91(1/2): 35-56.
- Agrios G.N. (1997). Plant Pathology, Fourth Edition ed., Academic Press, San Diego, 635 p.
- Aguhlon R. (1996). Les nouvelles techniques d'entretien du sol - Intérêt pour la viticulture, l'œnologie et l'environnement. *La Défense des Végétaux* 488: 41-46.
- Ahmed M.T., Ismail S.M.M., Mabrouk S.S. (1998). Residues of some chlorinated hydrocarbon pesticides in rain water, soil and ground water, and their influence on some soil microorganisms. *Environment International* 24(5-6): 665-670.
- Aitchinson J. (1995). Cultural landscapes in Europe: a geographical perspective. In Cultural landscapes of universal value, (von Droste B. P.H.R.M., ed.), Gustav Fischer Verlag, Jena, Germany: 272-288.
- Akbar K.F., Hale W.H.G., Headley A.D. (2003). Assessment of scenic beauty of the roadside vegetation in northern England. *Landscape and Urban Planning* 63: 139-144.
- Al Haj Khaled R., Duru M., Theau J.P., Plantureux S., Cruz P. (2005). Variation in leaf traits through seasons and N-availability levels and its consequences for ranking grassland species. *Journal of Vegetation Science* 16(4): 391-398.
- Alabouvette C., Backhouse D., Steinberg C., Donovan N.J., Edel-Hermann V., Burgess L.W. (2004). Microbial diversity in soil - Effects on crop health. In Managing Soil Quality: Challenges in Modern Agriculture, (Emholt S., Christensen B.T., eds.), CABI Publishing, New York: 121-138.
- Alabouvette C., Olivain C., Steinberg C. (2006). Biological control of plant diseases: the European situation. *European Journal of Plant Pathology* 114(3): 329-341.
- Alabouvette C., Raaijmakers J., De Boer W., Notz R., Défago G., Steinberg C., Lemanceau P. (2006). Concepts and Methods to Assess the Phytosanitary Quality of Soils. In Plant-microbe interactions and soil quality handbook, (Bloem J. et al., eds.), CABI Publishing, Wallingford, UK: 257-270.
- Alard V., Béranger C., Journet M. (2002). A la recherche d'une agriculture durable. Etude de systèmes herbagers économes en Bretagne, Collection Espace Ruraux, INRA, Paris, 340 p.
- Albrecht M., Duelli P., Müller C., Kleijn D., Schmid B. (2007). The Swiss agri-environment scheme enhances pollinator diversity and plant reproductive success in nearby intensively managed farmland. *Journal of Applied Ecology* 44(4): 813-822.
- Albrecht M., Duelli P., Schmid B., Müller C.B. (2007). Interaction diversity within quantified insect food webs in restored and adjacent intensively managed meadows. *Journal of Animal Ecology* 76(5): 1015-1025.
- Alix A., Cortesero A.M., Nènon J.P., Anger J.P. (2001). Selectivity assessment of chlorfenvinphos reevaluated by including physiological and behavioral effects on an important beneficial insect. *Environmental Toxicology and Chemistry* 20(11): 2530-2536.
- Allen M.F. (2007). Mycorrhizal fungi: Highways for water and nutrients in arid soils. *Vadose Zone Journal* 6(2): 291-297.
- Allen S.C., Jose S., Nair P.K.R., Brecke B.J., Nkedi-Kizza P., Ramsey C.L. (2004). Safety-net role of tree roots: evidence from a pecan (*Carya illinoensis* K. Koch)-cotton (*Gossypium hirsutum* L.) alley cropping system in the southern United States. *Forest Ecology and Management* 192(2-3): 395-407.
- Allen-Wardell G., Bernhardt P., Bitner R., Burquez A., Buchmann S., Cane J., Cox P.A., Dalton V., Feinsinger P., Ingram M., Inouye D., Jones C.E., Kennedy K., Kevan P., Koopowitz H., Medellín R., Medellín-Morales S., Nabhan G.P., Pavlik B., Tepedino V., Torchio P., Walker S. (1998). The potential consequences of pollinator declines on the conservation of biodiversity and stability of food crop yields. *Conservation Biology* 12(1): 8-17.
- Al-Mufti M.M., Sydes C.L., Furness S.B., Grime J.P., Band S.R. (1977). A quantitative analysis of shoot phenology and dominance in herbaceous vegetation. *Journal of Ecology* 65: 759-791.
- Alpert P., Bone E., Holzapfel C. (2000). Invasiveness, invasibility and the role of environmental stress in the spread of non-native plants. *Perspectives in Plant Ecology, Evolution and Systematics* 3(1): 52-66.
- Alpehi J., Bonkowski M., Scheu S. (1996). Protozoa, Nematoda and Lumbricidae in the rhizosphere of *Hordelymus europaeus* (Poaceae): Faunal interactions, response of microorganisms and effects on plant growth. *Oecologia* 106(1): 111-126.
- Alston D.G. (1994). Effect of apple orchard floor vegetation on density and dispersal of phytophagous and predaceous mites in Utah. *Agriculture, Ecosystems & Environment* 50(1): 73-84.
- Altieri M.A. (1986). L'Agroécologie - Bases scientifiques d'une agriculture alternative, Debard, 237 p.
- Altieri M.A., Letourneau D.K. (1982). Vegetation management and biological control in agroecosystems. *Crop Protection* 1: 405-430.
- Altieri M.A., Nicholls C.I. (2004). Effects of agroforestry systems on the ecology and management of insect pest populations. In Ecological engineering for pest management - Advances in habitat manipulation for Arthropods, (Gurr G.M. et al., eds.), CSIRO CABI, Collingwood Australia Oxon UK: 143-154.
- Altieri M.A., Nicholls C.I. (2004). Plant diversity and insect stability in agroecosystems. In Biodiversity and pest management in agroecosystems, (Altieri M.A., Nicholls C.I., eds.), Food Products Press, New-York: 29-45.

- Altieri M.A., Nicholss C.I. (2003). Soil fertility management and insect pests: harmonizing soil and plant health in agroecosystems. *Soil & Tillage Research* 72: 203-211.
- Altieri M.A., Schmidt L.L. (1986). The dynamics of colonizing arthropod communities at the interface of abandoned, organic and commercial apple orchards and adjacent woodland habitats. *Agriculture, Ecosystems & Environment* 16(1): 29-43.
- Altieri M.A., Schmidt L.L. (1986). Population trends and feeding preferences of flea beetles (*Phyllotreta cruciferae* Goeze) in collard-wild mustard mixtures. *Crop Protection* 5(3): 170-175.
- Altieri M.A., Schmidt L.L. (1987). Mixing cultivars of broccoli reduces cabbage aphid populations. *California Agriculture* 41(11-12): 24-26.
- Altieri M.A., Wilson R.C., Schmidt L.L. (1985). The effects of living mulches and weed cover on the dynamics of foliage- and soil-arthropod communities in three crop systems. *Crop Protection* 4(2): 201-213.
- Ambrosoli R., Petruzzelli L., Minati J.L., Marsan F.A. (2005). Anaerobic PAH degradation in soil by a mixed bacterial consortium under denitrifying conditions. *Chemosphere* 60(9): 1231-1236.
- Amiaud B., Touzard B. (2004). The relationships between soil seed bank, aboveground vegetation and disturbances in old embanked marshlands of Western France. *Flora* 199(1): 25-35.
- Anasiewicz A., Warakomska Z., Kozłowska A. (1989). Pollen of Umbelliferae family vegetables as a food of source for Syrphidae, Diptera. *Annales Universitatis Mariae Curie-Skłodowska Lublin-Polinia* sectio C, XLIV(18): 297-312.
- Ançay A., Delabays N., Mermillod G., De Joffrey J.P., Bohren C. (2006). Quel enherbement pour les cultures de framboisiers ? . *Revue Suisse de Viticulture, Arboriculture et d'Horticulture* 38(6): 363-369.
- Andow D.A. (1986). Plant diversification and insect population control in agroecosystems. In *Some aspects of Integrated Pest Management*, (Pimentel D.E., ed.), Cornell University Press, Ithaca, NY: 277-368.
- Andow D.A. (1988). Management of weeds for insect manipulation in agroecosystems. In *Weed management in agroecosystems: ecological approaches*, (Altieri M.A., Liebman M., eds.), CRC Press: 265-301.
- Andow D.A. (1990). Population dynamics of an insect herbivore in simple and diverse habitats. *Ecology* 72(3): 1006-1017.
- Andow D.A. (1991). Vegetational diversity and arthropod population response. *Annual Review of Entomology* 36: 561-586.
- Andow D.A., Risch S.J. (1985). Predation in diversified agroecosystems: relations between a coccinellid predator *Coleomegilla maculata* and its food. *Journal of Applied Ecology* 22(2): 357-372.
- Andrade D.D., Colozzi A., Balota E.L., Hungria M. (2003). Long-term effects of agricultural practices on microbial community. In *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-Economy, Policy*, KLUWER ACADEMIC PUBL, Dordrecht: 301-306.
- Andrade G., Mihara K.L., Linderman R.G., Bethlenfalvay G.J. (1997). Bacteria from the rhizosphere and hyphosphere of different arbuscular-mycorrhizal fungi. *Plant and Soil* 192: 71-79.
- Andreasen C., Stryhn H. (2008). Increasing weed flora in Danish arable fields and its importance for biodiversity. *Weed Research* 48: 1-9.
- Andrieu N., Coleno F.C., Duru M. (2008). L'organisation du système fourrager source de flexibilité face aux variations climatiques. In *Les exploitations d'élevage en mouvement. Flexibilités et dynamiques face aux aléas et aux nouveaux enjeux des filières et territoires*, (Dedieu B. et al., eds.), Editions Quae, Paris: 95-110.
- Andrieu N., Josien E., Duru M. (2007). Relationships between diversity of grassland vegetation, field characteristics and land use management practices assessed at the farm level. *Agriculture, Ecosystems & Environment* 120(2-4): 359-369.
- Andrieu N., Poix C., Josien E., Duru M. (2007). Simulation of forage management strategies considering farm-level land diversity: Example of dairy farms in the Auvergne. *Computers and Electronics in Agriculture* 55(1): 36-48.
- Anil L., Park J., Phipps R.H., Miller F.A. (1998). Temperate intercropping of cereals for forage: a review of the potential for growth and utilization with particular reference to the UK. *Grass & Forage Science* 53(4): 301-317.
- Ansquer P. (2006). Caractérisation agroécologique des végétations prairiales naturelles en réponse aux pratiques agricoles. Apports pour la construction d'outils de diagnostic. Thèse, Institut National Polytechnique de Toulouse, 272 p.
- Ansquer P., Theau J.P., Cruz P., Viegas J., Al Haj Khaled R., Duru M. (2004). Caractérisation de la diversité fonctionnelle des prairies naturelles. Une étape vers la construction d'outils pour gérer les milieux à flore complexe. *Fourrages* 179: 353-368.
- Antrop M. (2002). Rural-urban conflicts and opportunities, *The New Dimensions of the European Landscape*, Wageningen, NL, R. J., ed., Frontis workshop on the future of the European cultural landscape: 83-91.
- Apáti P., Kristó T.Z., Szóke E., Kéry A., Szentmihályi K., Vinkler P. (2003). Comprehensive evaluation of different *Solidaginis herba* extracts, 597, ISHS: 69-73.
- Aquilino K.M., Cardinale B.J., Ives A.R. (2005). Reciprocal effects of host plant and natural enemy diversity on herbivore suppression: an empirical study of a model tritrophic system. *Oikos* 108(2): 275-282.
- Asbakk K., Hrabok J.T., Oksanen A., Nieminen M., Waller P.J. (2006). Prolonged persistence of fecally excreted ivermectin from reindeer in a sub-arctic environment. *Journal of Agricultural and Food Chemistry* 54(24): 9112-9118.
- Ashman T.-L., Knight T.M., Steets J.A., Amarasekare P., Burd M., Campbell D.R., Dudash M.R., Mazer S.J., Johnston M., Mitchell R.J., Morgan M.T., Wilson W.G. (2004). Pollen limitation of plant reproduction: ecological and evolutionary causes and consequences. *Ecology* 85(9): 2408-2421.
- Asner G.P., Elmore A.J., Olander L.P., Martin R.E., Harris A.T. (2004). Grazing systems, ecosystem responses, and global change. *Annual Review of Environment and Resources* 29(1): 261-299.
- Astier C., Rock E., Chardigny J.M., Borel P., Coulon J.B. (2004). Effet de l'origine géographique et du traitement thermique du lait de vache sur sa qualité nutritionnelle, *11. Rencontres Recherches Ruminants*: 75.
- Athanasiadou S., Kyriazakis I., Jackson F., Coop R.L. (2000). Consequences of long-term feeding with condensed tannins on sheep parasitized with *Trichostrongylus colubriformis*. *International Journal for Parasitology* 30(9): 1025-1033.
- Atkinson P.W., Fuller R.J., Vickery J.A., Conway G.J., Tallowin J.R.B., Smith R.E.N., Haysom K.A., Ings T.C., Asteraki E.J., Brown V.K. (2005). Influence of agricultural management, sward structure and food resources on grassland field use by birds in lowland England. *Journal of Applied Ecology* 42(5): 932-942.
- Aubert M., Hedde M., Decaëns T., Bureau F., Margerie P., Alard D. (2003). Effects of tree canopy composition on earthworms and other macro-invertebrates in beech forests of Upper Normandy (France): The 7th international symposium on earthworm ecology · Cardiff · Wales · 2002. *Pedobiologia* 47(5-6): 904-912.

- Aubert M., Hedde M., Decaëns T., Margerie P., Alard D., Bureau F. (2005). Facteurs contrôlant la variabilité spatiale de la macrofaune du sol dans une hêtraie pure et une hêtraie-charmaie. *Compte Rendus - Biologies* 328(1): 57-74.
- Aufrère J., Carrère P., Dudillieu M., Baumont R. (2008). Estimation of nutritive value of grasses from semi-natural grasslands, by biological, chemical and enzymatic methods, 22. *General Meeting of the European Grassland Federation*, Uppsala, Suède.
- Aufrère J., Dudillieu M., Poncet C., Baumont R. (2005). Effect of condensed tannins in sainfoin on *in vitro* protein solubility of lucerne, 20. *International Grassland Congress* Dublin, Ireland, Wageningen Academic Publishers: 248.
- Auge R.M. (2004). Arbuscular mycorrhizae and soil/plant water relations. *Canadian Journal of Soil Science* 84(4): 373-381.
- Aurousseau B. (2002). Les radicaux libres dans l'organisme des animaux d'élevage : conséquences sur la reproduction, la physiologie et la qualité de leurs produits. *INRA Productions animales* 15(1): 67-82.
- Avio L., Pellegrino E., Bonari E., Giovannetti M. (2006). Functional diversity of arbuscular mycorrhizal fungal isolates in relation to extraradical mycelial networks. *New Phytologist* 172: 347-357.
- Axelrod D.I. (1960). The evolution of flowering plants. In *Evolution after Darwin*, (Tax S., ed.), 1, University of Chicago Press, Chicago: 227-305.
- Ayal Y. (2007). Trophic structure and the role of predation in shaping hot desert communities. *Journal of Arid Environments* 68(2): 171-187.
- Babendreier D., Schoch D., Kuske S., Dorn S., Bigler F. (2003). Non-target habitat exploitation by *Trichogramma brassicae*: what are the risks for endemic butterflies? . *Agricultural and Forest Entomology* 5(3): 199-208.
- Bachthaler G. (1968). Die Entwicklung der Ackerunkrautflora in Abhängigkeit von veränderten Feldbaumethoden. *Zeitschrift fuer Acker- und Pflanzenbau* 127: 327-358.
- Bachthaler G. (1968). Die Entwicklung der Ackerunkrautflora in Abhängigkeit von veränderten Feldbaumethoden. 1-Der Einfluss einer veränderten Feldbaumethoden. *Zeitschrift für Acker- und Pflanzenbau* 127(2): 149-170.
- Bahuaud D., Martínez-Ortiz de Montellano C., Chauveau S., Prevot F., Torres-Acosta F., Forraste I., Hoste H. (2006). Effects of four tanniferous plant extracts on the *in vitro* exsheathment of third-stage larvae of parasitic nematodes. *Parasitology* 132: 545-554.
- Bailey K.L., Lazarovits G. (2003). Suppressing soil-borne diseases with residue management and organic amendments. *Soil & Tillage Research* 72(2): 169-180.
- Bailey V.L., Smith J.L., Bolton H. (2002). Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration. *Soil Biology & Biochemistry* 34(7): 997-1007.
- Bakker E.S., Olf H., Boekhoff M., Gleichman J.M., Berendse F. (2004). Impact of herbivores on nitrogen cycling: contrasting effects of small and large species. *Oecologia* 138(1): 91-101.
- Baldrige C.D., Moran M.D. (2001). Behavioral means of coexisting in old fields by heterospecific arthropod predators (Araneae: Lycosidae, Salticidae; Insecta: Coleoptera, Carabidae). *Proceedings of the Entomological Society of Washington* 103(1): 81-88.
- Balesdent J., Chenu C., Balabane M. (2000). Relationship of soil organic matter dynamics to physical protection and tillage. *Soil and Tillage Research* 53(3/4): 215-230.
- Ballabeni P., Conconi D., Gateff S., Rahier M. (2001). Spatial proximity between two host plant species influences oviposition and larval distribution in a leaf beetle. *Oikos* 92(2): 225-234.
- Balvanera P., Daily G.C., Ehrlich P.R., Ricketts T.H., Bailey S.-A., Kark S., Kremen C., Pereira H. (2001). Conserving Biodiversity and Ecosystem Services. *Science* 291(5511): 2047.
- Balvanera P., Pfisterer A.B., Buchmann N., He J.-S., Nakashizuka T., Raffaelli D., Schmid B. (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecology Letters* 9(10): 1146-1156.
- Banaszak J. (1995). Changes in fauna of wild bees in Europe, Pedagogical Univeristy, Bydgoszcz, Poland, 220 p.
- Banks J.E. (2000). Effects of weedy field margins on *Myzus persicae* (Hemiptera : Aphididae) in a broccoli agroecosystem. *Pan-Pacific Entomologist* 76(2): 95-101.
- Barbault R. (2006). Un éléphant dans un jeu de quille: l'homme dans la biodiversité, Science ouverte, Le Seuil, Paris France, 265 p.
- Bärber P., Lo Cascio B. (2001). Long-term tillage and crop rotation effects on weed seedbank size and composition. *Weed Research* 41: 325-340.
- Bardgett R.D., Bowman W.D., Kaufmann R., Schmidt S.K. (2005). A temporal approach to linking aboveground and belowground ecology. *Trends in Ecology & Evolution* 20(11): 634-641.
- Bardgett R.D., Chan K.F. (1999). Experimental evidence that soil fauna enhance nutrient mineralization and plant nutrient uptake in montane grassland ecosystems. *Soil Biology & Biochemistry* 31(7): 1007-1014.
- Bardgett R.D., Cook R. (1998). Functional aspects of soil animal diversity in agricultural grasslands. *Applied Soil Ecology* 10(3): 263-276.
- Bardgett R.D., Jones A.C., Jones D.L., Kemmitt S.J., Cook R., Hobbs P.J. (2001). Soil microbial community patterns related to the history and intensity of grazing in sub-montane ecosystems. *Soil Biology & Biochemistry* 33(12-13): 1653-1664.
- Bardgett R.D., Smith R.S., Shiel R.S., Peacock S., Simkin J.M., Quirk H., Hobbs P.J. (2006). Parasitic plants indirectly regulate belowground properties in grassland ecosystems. *Nature* 439: 969-972.
- Bardgett R.D., Wardle D.A. (2003). Herbivore-mediated linkages between aboveground and belowground communities. *Ecology* 84: 2258-2268.
- Barker G.M. (1991). Biology of slugs (Agirolimacidae and Arionidae: Mollusca) in New Zealand hill country pastures. *Oecologia* 85(4): 581-595.
- Barros E., Curmi P., Hallaire V., Chauvel A., Lavelle P. (2001). The role of macrofauna in the transformation and reversibility of soil structure of an oxisol in the process of forest to pasture conversion. *Geoderma* 100: 193-213.
- Bartelt-Ryser J., Joshi J., Schmid B., Brandl H., Balsler T. (2005). Soil feedbacks of plant diversity on soil microbial communities and subsequent plant growth. *Perspectives in Plant Ecology, Evolution and Systematics* 7: 27-49.
- Barth D., Karrer M., Heinzemut E.M., Elster N. (1994). Colonization and Degradation of Cattle Dung - Aspects of Sampling, Fecal Composition, and Artificially Formed Pats. *Environmental Entomology* 23(3): 571-578.
- Barthell J.F., Randall J.M., Thorp R.W., Wenner A.M. (2001). Promotion of seed set in yellow star-thistle by honey bees: Evidence of an invasive mutualism. *Ecological Applications* 11(6): 1870-1883.

- Bascompte J., Jordano P., Melián C.J., Olesen J.M. (2003). The nested assembly of plant–animal mutualistic networks. *PNAS* 100(16): 9383-9387.
- Basset Y., Burckhardt D. (1992). Abundance, species richness, host utilization and host specificity of insect folivores from a woodland site, with particular reference to host architecture. *Revue suisse de Zoologie* 99(4): 771–791.
- Bateman A.J. (1948). Intrasexual selection in *Drosophila*. *Heredity* 2: 349-369.
- Bateman I.J., Brouwer R., Davies H., Day B.H., Deflandre A., Di Falco S., Georgiou S., Hadley D., Hutchins M., Jones A.P., Kay D., Leeks G., Lewis M., Lovett A.A., Neal C., Posen P., Rigby D., Turner R.K. (2006). Analysing the agricultural costs and non-market benefits of implementing the Water Framework Directive. *Journal of Agricultural Economics* 57(2): 221-237.
- Batra S.W.T. (1993). Male-fertile potato flowers are selectively buzz-pollinated only by *Bombus terricola* in upstate New York. *Journal of Kansas Entomological Society* 66: 252–254.
- Bauchart D., Roy A., Lorenz S., Ferlay A., Gruffat D., Chardigny J.M., Sébédio J.L., Chilliard Y., Durand D. (2007). Dietary supply of butter rich in trans 18:1 isomers or in 9cis, 11trans conjugated linoleic acid affects plasma lipoproteins in hypercholesterolemic rabbits. *Lipids* 42: 123-133.
- Bauman D.E., Corl B.A., Baumgard L.H., Griinari J.M. (2001). Conjugated linoleic acid (CLA) and the dairy cows. In *Recent advances in animal nutrition*, Nottingham University Press, Nottingham, UK: 221-250.
- Baumgarte S., Tebbe C.C. (2005). Field studies on the environmental fate of the Cry1Ab Bt-toxin produced by transgenic maize (MON810) and its effect on bacterial communities in the maize rhizosphere. *Molecular Ecology* 14(8): 2539-2551.
- Baumont R., Pomiès D. (2004). Feed preferences and voluntary intake of dairy heifers fed grass silage and hays offered singly or as a matter of choice, 20. *General Meeting of the European Grassland Federation*, 2004/06/21-24, Luzern, Switzerland: 1089-1091.
- Bazin G. (2003). La PAC contre la multifonctionnalité ? *Economie rurale* 2003: 236-242.
- Beane K.A., Bugg R.L. (1998). Natural and artificial shelter to enhance arthropod biological control agents. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Bugg C.H.P.R.L., ed.), University of California Press, Berkeley - Los Angeles - London: 239-253.
- Begon M., Harper J.L., Townsend C.R. (1996). *Ecology: Individuals, Populations and Communities*, Blackwell Science, London, 1068 p.
- Beintema A.J., Thissen J.B., Tensen D., Visser G.H. (1991). Feeding ecology of charadriiform chicks in agricultural grasslands. *Ardea* 79: 31-44.
- Bellon S., Guerin G. (1992). Old Holm Oak Coppices ... New Sylvopastoral Practices. *Vegetatio* 100: 307-316.
- Belnap J., Phillips S.L. (2001). Soil biota in an ungrazed grassland: response to annual grass (*Bromus tectorum*, L.) invasion. *Ecological Applications* 11: 1261-1275.
- Belovsky G.E., Slade J.B. (2000). Insect herbivory accelerates nutrient cycling and increases plant production. *Proceedings of the National Academy of Sciences* 97: 14412-14417.
- Bengtsson J., Angelstam P., Elmqvist T., Emanuelsson U., Folke C., Ihse M., Moberg F., Nyström M. (2003). Reserves, Resilience and Dynamic Landscapes. *Ambio* 32(6): 389-396.
- Bennett A.E., Bever J.D. (2007). Mycorrhizal species differentially alter plant growth and response to herbivory . . *Ecology* 88(1): 210-218.
- Bennett E.M., Balvanera P. (2007). The future of production systems in a globalized world. *Frontiers in Ecology and the Environment* 5: 191-198.
- Benton T.G., Bryant D.M., Cole L., Crick H.Q.P. (2002). Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology* 39(4): 673-687.
- Berkelmans R., Ferris H., Tenuta M., van Bruggen A.H.C. (2003). Effects of long-term crop management on nematode trophic levels other than plant feeders disappear after 1 year of disruptive soil management. *Applied Soil Ecology* 23(3): 223-235.
- Berlinger M.J., Dahan R., Mordechi S. (1988). Integrated pest management of organically grown greenhouse tomatoes in Israel. *Applied Agricultural Research* 3(5): 233-238.
- Bernard J.L., Granval P., Pasquet G. (1998). Les bords de champs cultivés pour une approche cohérente des attentes cynégétiques, agronomiques et environnementales. *Le Courrier de l'environnement de l'INRA* 34: 21-32.
- Bernard J.-L., Havet P., Fort M. (2007). *Productions végétales, pratiques agricoles et faune sauvage*, UIPP, ACTA, ONCFS, Paris, 251 p.
- Bernhard-Reversat F., Loumeto J.J., Laclau J.P. (2001). Litterfall, litter quality and decomposition changes with eucalypt hybrids and plantation age. In *Effect of exotic tree plantations on plant diversity and biological soil fertility in the Congo savanna: a reference to eucalypts*, (Bernhard-Reversat F., ed.), CIFOR, Bogor: 23-29.
- Bertiller M.B., Mazzarino M.J., Carrera A.L., Diehl P., Satti P., Gobbi M., Sain C.L. (2006). Leaf strategies and soil N across a regional humidity gradient in Patagonia. *Oecologia* 148: 612-624.
- Bertrand M., Lumaret J.-P. (1984). Réaction des populations de Microarthropodes du sol à l'enfouissement des fèces de mouton par des insectes Scarabaeidae en milieux à fortes contraintes climatiques. *Pedobiologia* 27: 51-66.
- Besle J.M., Lamaison J.L., Pradel P., Fraisse D., Viala D., Martin B. (2004). Les flavonoïdes, des fourrages au lait, 11. *Recherches Ruminants*: 67.
- Bever J.D., Schultz M., Pringle A., Morton J.B. (2001). Arbuscular mycorrhizal fungi: more diverse than meets the eye, and the ecological tale of why. *Bioscience* 51(11): 923-931.
- Bezemer T.M., Lawson C.S., Hedlund K., Edwards A.R., Brook A.J., Igual J.M., Mortimer S.R., van der Putten W.H. (2006). Plant species and functional group effects on abiotic and microbial soil properties and plant-soil feedback responses in two grasslands. *Journal of Ecology* 94: 893-904.
- Bhuiyan M.S.I., Wratten S.D. (1994). Grain aphid populations and their fall-off rate on different cultivars of wheat. *Bulletin OILB/SROP* 17(4): 27-35.
- Bianchi F., Booij C.J.H., Tschamtké T. (2006). Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B-Biological Sciences* 273(1595): 1715-1727.
- Bianchi F., Honek A., van der Werf W. (2007). Changes in agricultural land use can explain population decline in a ladybeetle species in the Czech Republic: evidence from a process-based spatially explicit model. *Landscape Ecology* 22(10): 1541-1554.

- Biesmeijer J.C., Roberts S.P.M., Reemer M., Ohlemuller R., Edwards M., Peeters T., Schaffers A.P., Potts S.G., Kleukers R., Thomas C.D., Settele J., Kunin W.E. (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313(5785): 351-354.
- Bigler F., Babendreier D., Kuske S. (2002). Ökologische Risiken bei der biologischen Bekämpfung des Maiszünslers mit *Trichogramma* Wespen. *AGRARforschung* 316-321.
- Birkett M.A., Campbell C.A.M., Chamberlain K., Guerrieri E., Hick A.J., Martin J.L., Matthes M., Napier J.A., Pettersson J., Pickett J.A., Poppy G.M., Pow E.M., Pye B.J., Smart L.E., Wadhams G.H., Wadhams L.J., Woodcock C.M. (2000). New roles for cis-jasmone as an insect semiochemical and in plant defense. *Proceedings of the National Academy of Sciences of the United States of America* 97(16): 9329-9334.
- Biro B., Villanyi I., Fuzy A., Naar Z. (2005). Bacterial and fungal colonization in the rhizosphere of genetically modified (Bt) and isogenic control maize. *Agrokemia es Talajtan* 54(1/2): 189-202.
- Bischoff A., Cremieux L., Smilauerova M., Lawson C.S., Mortimer S.R., Dolezal J., Lanta V., Edwards A.R., Brook A.J., Macel M., Leps J.A.N., Steinger T., Muller-Scharer H. (2006). Detecting local adaptation in widespread grassland species - the importance of scale and local plant community. *Journal of Ecology* 94(6): 1130-1142.
- Björkman M. (2007). Effects of Intercropping on the Life Cycle of the Turnip Root Fly (*Delia floralis*) - Behaviour, Natural Enemies and Host Plant Quality. Doctoral thesis, Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Faculty of Natural Resources and Agricultural Sciences, Uppsala, 42 p.
- Blackwood C.B., Buyer J.S. (2004). Soil microbial communities associated with Bt and non-Bt corn in three soils. *Journal of Environmental Quality* 33(3): 832-836.
- Blanchart E., Albrecht A., Brown G., Decaens T., Duboiset A., Lavelle P., Mariani L., Roose E. (2004). Effects of tropical endogeic earthworms on soil erosion. *Agriculture, Ecosystems & Environment* 104(2): 303-315.
- Blanchart E., Brown G.G., Chernyanskii S.S., Deleporte P., Feller C., Goulet (2005). Perception et popularité des vers de terre avant et après Darwin. *Étude et Gestion des Sols* 12(2): 145-152.
- Blanchart E., Lavelle P., Braudeau E., LeBissonnais Y., Valentin C. (1997). Regulation of soil structure by geophagous earthworm activities in humid savannas of Cote d'Ivoire. *Soil Biology & Biochemistry* 29(3-4): 431-439.
- Blossey B., Hunt-Joshi T.R. (2003). Belowground herbivory by insects: Influence on plants and aboveground herbivores. *Annual Review of Entomology* 48: 521-547.
- Blot J., Mollet T., Dela Torre C. (2007). *Vespa velutina* - frelon asiatique. *Bull. Tech. Apic.* 34: 205-210.
- Blouin M., Zuily-Fodil Y., Pham-Thi A.-T., Laffray D., Reversat G., Pando A., Tondoh J., Lavelle P. (2005). Belowground organism activities affect plant aboveground phenotype, inducing plant tolerance to parasites. *Ecology Letters* 8(2): 202-208.
- Boag B., Palmer L.F., Neilson R., Legg R., Chambers S.J. (1997). Distribution, prevalence and intensity of earthworm populations in arable land and grassland in Scotland. *Annals of Applied Biology* 130(1): 153-165.
- Boag B., Yeates G.W., Johns P.M. (1998). Limitation to the distribution and spread of terrestrial flatworms with special reference to the New Zealand flatworm (*Arthropostia triangulata*). *Pedobiologia* 42(5-6): 495-503.
- Bockus W.W., Shroyer J.P. (1998). The impact of reduced tillage on soilborne plant pathogens. *Annual Review of Phytopathology* 36: 485-500.
- Bohan D.A., Bohan A.C., Glen D.M., Symondson W.O.C., Wiltshire C.W., Hughes L. (2000). Spatial dynamics of predation by carabid beetles on slugs. *Journal of Animal Ecology* 69(3): 367-379.
- Bohanec M., Cortet J., Griffiths B., Znidarsic M., Debeljak M., Caul S., Thompson J., Krogh P.H. (2007). A qualitative multi-attribute model for assessing the impact of cropping systems on soil quality. *Pedobiologia* 51(3): 239-250.
- Boiffin J., Stengel P. (1999). Réapprendre le sol : nouvel enjeu pour l'agriculture et l'espace rural. In *Déméter 2000 : Economie et Stratégies agricoles, 2000*, Armand Colin, Paris: 147-213.
- Boiteau G. (1986). Native predators and the control of potato aphids. *Canadian Entomologist* 118(11): 1177-1183.
- Boizard H., Richard G., Brancourt-Hulmel M., Guérif J. (2000). Effect of cropping systems on change in bulk density, penetration resistance and hydraulic conductivity in subsoil. In *Subsoil compaction: distribution, processes and consequences*, (Horn R. *et al.*, eds.), Advances in Geocology, 32, Catena Verlag, Reiskirchen (DEU): 233-241.
- Boller E.F., Malavolta C., Jörg E. (1997). Guidelines for integrated production of arable crops in Europe. IOBC Technical Guideline III. *IOBC WPRS BULLETIN* 20(5): 16 p.
- Bommarco R. (1999). Feeding, reproduction and community impact of a predatory carabid in two agricultural habitats. *Oikos* 87(1): 89-96.
- Bonkowski M., Brandt F. (2002). Do soil protozoa enhance plant growth by hormonal effects? *Soil Biology & Biochemistry* 34(11): 1709-1715.
- Bonkowski M., Cheng W.X., Griffiths B.S., Alpei G., Scheu S. (2000). Microbial-faunal interactions in the rhizosphere and effects on plant growth. *European Journal of Soil Biology* 36(3-4): 135-147.
- Bonkowski M., Roy J. (2005). Soil microbial diversity and soil functioning affect competition among grasses in experimental microcosms. *Oecologia* 143(2): 232-240.
- Bonny S. (1998). Quels rendements demain ? Perspectives d'évolution des rendements des grandes cultures. *Notes et Etudes Economiques* 8: 7-77.
- Bonte D. (2005). Anthropogenic induced changes in nesting densities of the dune-specialised digger wasp *Bembix rostrata* (Hymenoptera : Sphecidae). *European Journal of Entomology* 102(4): 809-812.
- Booth B.D., Swanton C.J. (2002). Assembly theory applied to weed communities. *Weed Science* 50: 2-13.
- Borer E.T., Briggs C.J., Murdoch W.W., Swarbrick S.L. (2003). Testing intraguild predation theory in a field system: does numerical dominance shift along a gradient of productivity? *Ecology Letters* 6(10): 929-935.
- Borman M.M., Johnson D.E., Krueger W.C. (1992). Soil moisture extraction by vegetation in a Mediterranean/Maritime climatic regime. *Agronomy Journal* 84(5): 897-904.
- Bornemissza G.F. (1976). The Australian dung beetle project, 1965-1975. *Australian Meat Research Committee Review* 30: 1-30.
- Bornemissza G.F., Williams C.H. (1970). An effect of dung beetle activity on plant yield. *Pedobiologia* 10: 1-7.

- Borre J.v., Bonte D., Maelfait J.P. (2006). Interdemic variation of cannibalism in a wolf spider (*Pardosa monticola*) inhabiting different habitat types. *Ecological Entomology* 31(2): 99-105.
- Bosch J. (1994). Improvement of field management of *Osmia cornuta* (Latreille) (Hymenoptera, Megachilidae) to pollinate almond. *Apidologie* 25(1): 71-83.
- Bosch J., Blas M. (1994). Foraging behaviour and pollinating efficiency of *Osmia cornuta* and *Apis mellifera* on almond (Hymenoptera, Megachilidae and Apidae). *Applied Entomology and Zoology* 29: 1-9.
- Bosch J., Kemp W., Trostle G.E. (2006). Bee Population Returns and Cherry Yields in an Orchard Pollinated with *Osmia lignaria* (Hymenoptera: Megachilidae). *Journal of Economic Entomology* 99(2): 408-413.
- Boschi C., Baur B. (2007). The effect of horse, cattle and sheep grazing on the diversity and abundance of land snails in nutrient-poor calcareous grasslands. *Basic and Applied Ecology* 8(1): 55-65.
- Bossuyt H., Deneff K., Six J., Frey S.D., Merckx R., Paustian K. (2001). Influence of microbial populations and residue quality on aggregate stability. *Applied Soil Ecology* 16(3): 195-208.
- Bossuyt H., Six J., Hendrix P.F. (2005). Protection of soil carbon by microaggregates within earthworm casts. *Soil Biology & Biochemistry* 37(2): 251-258.
- Bottenberg H., Tamo M. (1998). Occurrence of phytophagous insects on wild *Vigna* sp. and cultivated cowpea: comparing the relative importance of host-plant resistance and millet intercropping. *Agriculture Ecosystems & Environment* 70(2-3): 217-229.
- Boubakri H., Beuf M., Simonet P., Vogel T.M. (2006). Development of metagenomic DNA shuffling for the construction of a xenobiotic gene. *Gene* 375: 87-94.
- Bounfour M., Jebbour F., Wadjiny J. (2005). Biological traits of invasive insect species harmful to Moroccan agriculture. In Plant protection and plant health in Europe: introduction and spread of invasive species, held at Humboldt University, Berlin, Germany, 9-11 June 2005, (Alford D.V., Backhaus G.F., eds.), Alton UK: 95-100.
- Bourbouze A., Donadieu P. (1987). L'élevage sur parcours en régions méditerranéennes, Options Méditerranéennes, CIHEAM-IAM, Montpellier, 104 p.
- Boutin C., Jobin B., Belanger L. (2003). Importance of riparian habitats to flora conservation in farming landscapes of southern Quebec, Canada. *Agriculture, Ecosystems & Environment* 94(1): 73-87.
- Bouvier J.C., Toubon J.F., Boivin T., Sauphanor B. (2005). Effects of apple orchard management strategies on the great tit (*Parus major*) in southeastern France. *Environmental Toxicology and Chemistry* 24(11): 2846-2852.
- Bouwman E.G. (1996). Direct emission of nitrous oxide from agricultural soils. *Nutrient Cycling in Agroecosystems* 46: 53-70.
- Bouwmeester H.J., Smid H.G. (1995). Seed yield in caraway (*Carum carvi*). 1. Role of pollination. *Journal of Agricultural Science* 124: 235-244.
- Boyd J.M. (1958). The ecology of earthworms in cattle grazed machair in Tiree, Argyll. *Journal of Animal Ecology* 27: 147-157.
- Bradford M.A., Tordoff G.M., Eggers T., Jones T.H., Newington J.E. (2002). Microbiota, fauna, and mesh size interactions in litter decomposition. *Oikos* 99(2): 317-323.
- Brandenburg R.L. (1985). The effect of field applications of insecticides for variegated cutworm, *Peridroma saucia* (Hubner) (Noctuidae: Lepidoptera) control on non-target arthropods in alfalfa. *Journal of the Kansas Entomological Society* 58(3): 437-441.
- Bräuer I., Müssner R., Marsden K., Oosterhuis F., Rayment M., Miller C., Dodoková A. (2006). The Use of Market Incentives to Preserve Biodiversity : Final Report, A project under the Framework contract for economic analysis, Ecologic, ENV.G.1/FRA/2004/0081: 51 p. <http://ec.europa.eu/environment/enveco/others/pdf/mbi.pdf>
- Brauman K.A., Daily G.C., Duarte T.K., Mooney H.A. (2007). The nature and value of ecosystem services: an overview highlighting hydrological services. *Annual Reviews of Environment and Resources* 32: 67-98
- Breland T.A. (1994). Measured and Predicted Mineralization of Clover Green Manure at Low-Temperatures at Different Depths in 2 Soils. *Plant and Soil* 166(1): 13-20.
- Bretagnolle V. (2004). Prairies et cultures fourragères : Quels enjeux pour les oiseaux dans les milieux céréaliers intensifs ? . *Fourrages* 178: 171-178.
- Bretagnolle V., Houte S. (2006). L'outarde canepetière en plaine céréalière : une extinction inéluctable. *Revue Espaces Naturels* 15(Juillet 2006): 28.
- Brett M.T., Goldman C.R. (1996). A meta-analysis of the freshwater trophic cascade, *National Academy of Sciences of the USA*, 93: 7723-7726.
- Brewer M.J., Noma T., Elliott N.C. (2005). Hymenopteran parasitoids and dipteran predators of the invasive aphid *Diuraphis noxia* after enemy introductions: temporal variation and implication for future aphid invasions. *Biological Control* 33(3): 315-323.
- Brinckerhoff-Jackson J. (1984). Discovering the vernacular landscape. (Traduction française (2003) : A la découverte du paysage vernaculaire), (Actes Sud-ENSP A.-P., ed.), Yale University Press, New Haven, USA., 278.
- Briones M.J.I., Ostle N.J., Garnett M.H. (2007). Invertebrates increase the sensitivity of non-labile soil carbon to climate change. *Soil Biology & Biochemistry* 39(3): 816-818.
- Brodeur J., Rosenheim J.A. (2000). Intraguild interactions in aphid parasitoids. *Entomologia Experimentalis et Applicata* 97(1): 93-108.
- Broughton L.C., Gross K.L. (2000). Patterns of diversity in plant and soil microbial communities along a productivity gradient in a Michigan old-field. *Oecologia* 125(3): 420-427.
- Brown G.G., Barois I., Lavelle P. (2000). Regulation of soil organic matter dynamics and microbial activity in the drilosphere and the role of interactions with other edaphic functional domains. *European Journal of Soil Biology* 36(3-4): 177-198.
- Brown M.W. (2001). Flowering ground cover plants for pest management in peach and apple orchards. *Bulletin OILB/SROP* 24(5): 379-382.
- Brown M.W., Miller S.S. (1998). Coccinellidae (Coleoptera) in apple orchards of eastern West Virginia and the impact of invasion by *Harmonia axyridis*. *Entomological News* 109(2): 143-151.
- Brown M.W., Tworowski T. (2004). Pest management benefits of compost mulch in apple orchards. *Agriculture, Ecosystems & Environment* 103(3): 465-472.
- Brown R.L., Fridley J.D. (2003). Control of plant species diversity and community by species immigration: seed richness versus seed density. *Oikos* 102: 15-24.

- Bru D., Sarr A., Philippot L. (2007). Relative abundances of proteobacterial membrane-bound and periplasmic nitrate reductases in selected environments. *Applied and Environmental Microbiology* 73(18): 5971-5974.
- Bruce T.J.A., Martin J.L., Pickett J.A., Pye B.J., Smart L.E., Wadhams L.J. (2003). cis-Jasmone treatment induces resistance in wheat plants against the grain aphid, *Sitobion avenae* (Fabricius) (Homoptera: Aphididae). *Pest Management Science* 59(9): 1031-1036.
- Bruinenberg M.H., Valk H., Korevaar H., Struik P.C. (2002). Factors affecting digestibility of temperate forages from seminatural grasslands: a review. *Grass & Forage Science* 57: 292-301.
- Bruinenberg M.J., Valk H., Struik P.C. (2003). Voluntary intake and *in vivo* digestibility of forages from semi-natural grasslands in dairy cows. *Netherlands journal of agricultural science* 51(3): 219-235.
- Brunet B., Aufrere J., El Babili F., Fouraste I., Hoste H. (2007). The kinetics of exsheathment of infective nematode larvae is disturbed in the presence of a tannin-rich plant extract (sainfoin) both *in vitro* and *in vivo*. *Parasitology* 134: 1253-1262.
- Brunet S., Hoste H. (2006). Monomers of Condensed Tannins Affect the Larval Exsheathment of Parasitic Nematodes of Ruminants. *Journal of Agricultural and Food Chemistry* 54(20): 7481-7487.
- Brussaard L., BehanPelletier V.M., Bignell D.E., Brown V.K., Didden W., Folgarait P., Fragoso C., Freckman D.W., Gupta V.V.S.R., Hattori T., Hawsworth D.L., Klopatek C., Lavelle P., Malloch D.W., Rusek J., Soderstrom B., Tiedje J.M., Virginia R.A. (1997). Biodiversity and ecosystem functioning in soil. *Ambio* 26(8): 563-570.
- Brussaard L., de Ruiter P.C., Brown G.G. (2007). Soil biodiversity for agricultural sustainability. *Agriculture, Ecosystems & Environment* 121(3): 233-244.
- Brussaard L., Kuyper T.W., Didden W.A.M., de Goede R.G.M., Bloem J. (2004). Biological soil quality from biomass to biodiversity - importance and resilience to management stress and disturbance. In *Managing soil quality: challenges in modern agriculture*, (Schjonning P. *et al.*, eds.), CABI: 139-161.
- Brust G.E. (1994). Natural enemies in straw-mulch reduce Colorado potato beetle populations and damage in potato. *Biological Control* 4(2): 163-169.
- Büchi R. (2002). Mortality of pollen beetle (*Meligethes* spp.) larvae due to predators and parasitoids in rape fields and the effect of conservation strips. *Agriculture, Ecosystems & Environment* 90(3): 255-263.
- Buchin S., Martin B., Dupont D., Bornard A., Achilleos C. (1999). Influence of the composition of alpine highland pasture on the chemical, rheological and sensory properties of cheese. *Journal of Dairy Science* 66: 579-588.
- Buchmann S.L., Nabhan G.P. (1996). *The forgotten pollinators*, Island Press, Washington USA, 292 p.
- Buckland S.M., Grime J.P. (2000). The effects of trophic structure and soil fertility on the assembly of plant communities: a microcosm experiment. *Oikos* 91(2): 336-352.
- Buddle C.M., Walker S.E., Rypstra A.L. (2003). Cannibalism and density-dependent mortality in the wolf spider *Pardosa milvina* (Araneae: Lycosidae). *Canadian Journal of Zoology* 81(8): 1293-1297.
- Bugaud C., Buchin S., Coulon J.B., Hauwuy A., Dupont D. (2001). Influence of the nature of alpine pastures on plasmin activity, fatty acid and volatile compound composition of milk. *Lait* 81: 401-414.
- Bugaud C., Buchin S., Hauwuy A., Coulon J.B. (2002). Texture et flaveur du fromage selon la nature du pâturage : cas du fromages d'Abondance. *INRA Productions animales* 15(1): 31-36.
- Bugaud C., Doreau M., Chabrot J., Hauwuy A., Buchin S. (2000). Composition en acides gras des laits alpins. Relation avec la composition en acides gras des pâtures, 11. *Réunion du sous-réseau Pâturage de montagnes*, Luz-Saint-Sauveur F., ed., FAO - CIHEAM: 3 p.
- Bugg R.L., Pickett C.H. (1998). Introduction: Enhancing biological control - Habitat management to promote natural enemies of agricultural pests. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Pickett C.H., Bugg R.L., eds.), University of California Press, Berkeley - Los Angeles - London: 1-23.
- Bukovinsky T., Potting R.P.J., Clough Y., van Lenteren J.C., Vet L.E.M. (2005). The role of pre- and post-alighting detection mechanisms in the responses to patch size by specialist herbivores. *Oikos* 109(3): 435-446.
- Buller H. (2000). Re-Creating Rural Territories: LEADER in France. *Sociologia Ruralis* 40: 191-199.
- Bullock D.G. (1992). Crop Rotation. *Critical Reviews in Plant Sciences* 11: 309-326.
- Bullock J.M., Pywell R.E., Walker K.J. (2007). Long-term enhancement of agricultural production by restoration of biodiversity. *Journal of Applied Ecology* 44: 6-12.
- Bullock J.M., Pywell R.F., Burke M.J.W., Walker K.J. (2001). Restoration of biodiversity enhances agricultural production. *Ecology Letters* 4: 185-189.
- Bünemann E.K., Schwenke G.D., Van Zwieten L. (2006). Impact of agricultural inputs on soil organisms - a review. *Australian Journal of Soil Research* 44(4): 379-406.
- Buntin G.D. (1998). Cabbage seedpod weevil (*Ceutorhynchus assimilis*, Paykull) management by trap cropping and its effect on parasitism by *Trichomalus perfectus* (Walker) in oilseed rape. *Crop Protection* 17: 299-305.
- Burel F., Baudry J. (1999). *Ecologie du paysage : concepts, méthodes et applications*, Tec & Doc, Paris, 359 p.
- Burgio G., Ferrari R., Boriani L., Pozzati M., Lenteren J.v. (2006). The role of ecological infrastructures on Coccinellidae (Coleoptera) and other predators in weedy field margins within northern Italy agroecosystems. *Bulletin of Insectology* 59(1): 59-67.
- Burgio G., Ferrari R., Pozzati M., Boriani L. (2004). The role of ecological compensation areas on predator populations: an analysis on biodiversity and phenology of Coccinellidae (Coleoptera) on non-crop plants within hedgerows in Northern Italy. *Bulletin of Insectology* 57(1): 1-10.
- Burn A.J. (1988). Assessment of the impact of pesticides on invertebrate predation in cereal crops. *Aspects of Applied Biology* 17(1): 279-288.
- Burn A.J. (1992). Interactions between cereal pests and their predators and parasites. In *Pesticides, cereal farming and the environment: the Boxworth project*, (Greig-Smith P.W. *et al.*, eds.), Stationery Office Books London 110-131.
- Bush L., Slosser J.E., Worrall W.D., Horner N.V. (1991). Potential of wheat cultivar mixtures for greenbug (Homoptera: Aphididae) management. *Journal of Economic Entomology* 84(5): 1619-1624.
- Caamal-Maldonado J.A., Jimenez-Osornio J.J., Torres-Barragan A., Anaya A.L. (2001). The use of allelopathic legume cover and mulch species for weed control in cropping systems. *Agronomy Journal* 93: 27-36.

- CABI (2004). *Solidago canadensis* L. data sheet, Crop Protection Compendium, CAB International Wallingford, UK, 8 p.
- Cadisch G., Rowe E., Suprayogo D., van Noordwijk M. (2004). Safety-nets and filter functions of tropical agroforestry systems. *In* Controlling Nitrogen Flows and Losses, (D.J. Hatch D.R.C., S.C. Jarvis and J.A. Roker, ed.): 406-414.
- Caldeira M.C., Hector A., Loreau M., Pereira J.S. (2005). Species richness, temporal variability and resistance of biomass production in a Mediterranean grassland. *Oikos* 110: 115-123.
- Caldeira M.C., Ryel R.J., Lawton J.H., Pereira J.S. (2001). Mechanisms of Positive Biodiversity-production Relationships: Insights Provided by ¹³C Analysis in Experimental Mediterranean Grassland Plots. *Ecology Letters* 4: 439-443.
- Calderon F., Chauveau-Duriot B., Martin B., Graulet B., Doreau M., Nozières P. (2007). Variations in carotenoids, vitamins A and E, and color in cow's plasma and milk during late pregnancy and the first three months of lactation. *Journal of Dairy Science* 90: 2335-2346.
- Campbell C.A.M., Cone W.W. (1994). Influence of predators on population development of *Phorodon humuli* (Homoptera: Aphididae) on hops. *Environmental Entomology* 23(6): 1391-1396.
- Canals R.M., Eviner V.T., Herman D.J., Chapin F.S.I. (2005). Plant colonizers shape early N-dynamics in gopher-mounds *Plant and Soil* 276: 327-334.
- Cannon R.J.C. (1998). The implications of predicted climate change for insect pests in the UK, with emphasis on non-indigenous species. *Global Change Biology* 4(7): 785-796.
- Cantelo W.W., Sanford L.L. (1984). Insect population response to mixed and uniform plantings of resistant and susceptible plant material. *Environmental Entomology* 13(5): 1443-1445.
- Caraveli H. (2000). A comparative analysis on intensification and extensification in mediterranean agriculture: dilemmas for LFAs policy. *Journal of Rural Studies* 16(2): 231-242.
- Cardinale B.J., Srivastava D., Emmett Duffy J., Wright I.J., Downing A.L., Sankaran M., Jouseau C. (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature* 443: 989-992.
- Cardinale B.J., Wright J.P., Cadotte M.W., Carroll I.T., Hector A., Srivastava D.S., Loreau M., Weis J.J. (2007). Impacts of plant diversity on biomass production increase through time because of species complementarity. *Proceedings of the National Academy of Sciences* 104(46): 18123-18128.
- Cardon Z.G., Gage D.J. (2006). Resource exchange in the rhizosphere: Molecular tools and the microbial perspective. *Annual Review of Ecology Evolution and Systematics* 37: 459-488.
- Cardwell C., Hassall M., White P. (1994). Effects of headland management on carabid beetle communities in Breckland cereal fields. *Pedobiologia* 38(1): 50-62.
- Carena M.J., Glogoza P. (2004). Resistance of maize to the corn leaf aphid: a review. *Maydica* 49(4): 241-254.
- Carney K.M., Hungate B.A., Drake B.G., Megonigal J.P. (2007). Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of Sciences of the United States of America* 104(12): 4990-4995.
- Carof M., de Tourdonnet S., Saulas P., Le Floch D., Roger-Estrade J. (2007). Undersowing wheat with different living mulches in a no-till system. II. Competition for light and nitrogen. *Agronomy for Sustainable Development* 27(4): 357-365.
- Carr P.M., Kandel H.J., Porter P.M., Horsley R.D., Zwinger S.F. (2006). Wheat cultivar performance on certified organic fields in Minnesota and North Dakota. *Crop Science* 46(5): 1963-1971.
- Carreck N.L., Williams I.H. (2002). Food for insect pollinators on farmland: insect visits to flowers of annual seed mixtures. *Journal of Insect Conservation* 6(1): 13-23.
- Carreck N.L., Williams I.H., Little D.J. (1997). The movement of honey bee colonies for crop pollination and honey production by beekeepers in Great Britain. *Bee world* 78(2): 67-77.
- Carsoulle J. (1986). Influence de quelques techniques culturales sur le ruissellement et l'érosion en vignoble de coteaux (Beaujolais), 2. *Symposium International sur la Non Culture de la Vigne*, Montpellier (France), 1986/11/26-28, ANPP, ed.: 345-352.
- Carsoulle J. (1995). L'enherbement permanent du vignoble - Influence sur la production viticole et son environnement. *Phytoma, La Défense des Végétaux* 478: 38-41.
- Carter P.E., Rypstra A.L. (1995). Top-Down Effects in Soybean Agroecosystems - Spider Density Affects Herbivore Damage. *Oikos* 72(3): 433-439.
- Carvell C., Meek W.R., Pywell R.F., Goulson D., Nowakowski M. (2007). Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *Journal of Applied Ecology* 44(1): 29-40.
- Carvell C., Meek W.R., Pywell R.F., Nowakowski M. (2004). The response of foraging bumblebees to successional change in newly created arable field margins. *Biological Conservation* 118: 327-339.
- Castane C., Alomar O., Goula M., Gabarra R. (2004). Colonization of tomato greenhouses by the predatory mirid bugs *Macrolophus caliginosus* and *Dicyphus tamaninii*. *Biological Control* 30(3): 591-597.
- Casula P., Wilby A., Thomas M.B. (2006). Understanding biodiversity effects on prey in multi-enemy systems. *Ecology Letters* 9(9): 995-1004.
- Catroux G., Hartmann A., Revellin C. (2001). Trends in rhizobial inoculant production and use. *Plant and Soil* 230(1): 21-30.
- Caubel G., Chaubet B. (1985). Hatching and multiplication of *Heterodera schachtii* in the presence of rape or radishes. *Agronomie* 5(5): 463-466.
- Caubel-Forget V., Grimaldi C., Rouault F. (2001). Contrasted dynamics of nitrate and chloride in groundwater submitted to the influence of a hedge *Compte Rendus de l'Académie des Sciences, Série IIA* 332: 107-113.
- Caussanel J.P. (1989). Nuisibilité et seuil de nuisibilité des mauvaises herbes dans une culture annuelle : relation de concurrence bispécifique. *Agronomie* 9: 219-240.
- Cava R., Ruiz J., López-Bote C., Martín L., García C., Ventanas J., Antequera T. (1997). Influence of finishing diet on fatty acid profiles of intramuscular lipids, triglycerides and phospholipids in muscles of the Iberian pig. *Meat Science* 45(2): 263-270.
- Cavagnaro T.R., Smith F.A., Smith S.E., Jakobsen I. (2005). Functional diversity in arbuscular mycorrhizas: exploitation of soil patches with different phosphate enrichment differs among fungal species. *Plant, Cell and Environment* 28(5): 642-650.
- Chabert A., Gandrey J. (2005). Impact of some insecticides on Carabidae and consequences for slug populations. *Bulletin OILB/SROP* 28(6): 111-113.

- Chaboussou F. (1970). Influence des pesticides sur la plante : conséquences écologiques. *Bulletin de la Société Française d'Ecologie* 3: 146-158.
- Chagnon M., Gingras J., de Oliveira D. (1993). Complementary aspects of strawberry pollination by honey and indigenous bees (Hymenoptera). *Journal of Economic Entomology* 86(2): 416-420.
- Chambers S. (1998). Short- and Long-Term Effects of Clearing Native Vegetation for Agricultural Purposes, Flinders University of South Australia, Adelaide.
- Chambon J.-P. (1984). Structure d'une biocénose céréalière. In *Faune et flore auxiliaires en agriculture*, (ACTA, ed.): 19-27.
- Champion R.A., Orr R.J., Penning P.D., Rutter S.M. (2004). The effect of the spatial scale of heterogeneity of two herbage species on the grazing behaviour of lactating sheep. *Applied Animal Behaviour Science* 88: 61-76.
- Chan K.M.A., Shaw M.R., Cameron D.R., Underwood E.C., Daily G.C. (2006). Conservation planning for ecosystem services. *Plos Biology* 4(11): 2138-2152.
- Chang G.C. (1996). Comparison of single versus multiple species of generalist predators for biological control. *Environmental Entomology* 25(1): 207-212.
- Chang G.C., Eigenbrode S.D. (2004). Delineating the effects of a plant trait on interactions among associated insects. *Oecologia* 139(1): 123-130.
- Chantelot E. (2003). L'enherbement de la vigne, ITAB, 4 p.
- Chapin F.S. (2003). Effects of plant traits on ecosystem and regional processes: a conceptual framework for predicting the consequences of global change. *Annals of Botany* 91: 1-9.
- Chapin F.S., Zavaleta E.S., Eviner V.T., Naylor R.L., Vitousek P.M., Reynolds H.L., Hooper D.U., Lavorel S., Sala O.E., Hobbie S.E., Mack M.C., Diaz S. (2000). Consequences of changing biodiversity. *Nature* 405: 234-242.
- Chardigny J.M. (2007). *Trans* fatty acids from partially hydrogenated oils banned from New York restaurants. What about ruminant *trans* fatty acids? *European journal of lipid science and technology* 109(7): 645-646.
- Chaudhary O.P. (2006). Diversity, foraging behaviour of floral visitors and pollination ecology of fennel (*Foeniculum vulgare* Mill.). *Journal of spices and aromatic crops* 15(1): 34-41.
- Chauvel A., Grimaldi M., Barros E., Blanchart E., Sarrazin M., Lavelle P. (1999). Pasture damage by an Amazonian earthworm. *Nature* 398(6722): 32-33.
- Chauvel B., Dessaint F., Lonchamp J.-P., Gasquez J. (2005). Cinq élues et des candidates - Enquête sur les mauvaises herbes envahissantes en grandes cultures en France. *Phytoma, La Défense des Végétaux* 578: 16-20.
- Chauvel B., Gasquez J. (1993). Le coquelicot est-il symbole de naturel ?, *Faut-il sauver les mauvaises herbes?*, Gap, France, Ed. Conservatoire Botanique National de Gap-Charance: 237.
- Chauzat M.P., Higes M., Martín-Hernández R., Meana A., Cougoule N., Faucon J.P. (2007). Presence of *Nosema ceranae* in French honey bee colonies. *J. Apic. Res.* 46: 127-128.
- Cheesman O.D. (1998). The impact of some field boundary management practices on the development of *Dipsacus fullonum* L. flowering stems, and implications for conservation. *Agriculture, Ecosystems & Environment* 68(1-2): 41-49.
- Chen B.R., Wise D.H. (1999). Bottom-up limitation of predaceous arthropods in a detritus-based terrestrial food web. *Ecology* 80(3): 761-772.
- Chen C.J. (1990). Integrated management of pine caterpillars in China, Chinese Forestry Publication House, Beijing.
- Chen Y.H., Welter S.C. (2002). Abundance of a native moth *Homoeosoma electellum* (Lepidoptera : Pyralidae) and activity of indigenous parasitoids in native and agricultural sunflower habitats. *Environmental Entomology* 31(4): 626-636.
- Chen Y.H., Welter S.C. (2005). Crop domestication disrupts a native tritrophic interaction associated with the sunflower, *Helianthus annuus* (Asterales : Asteraceae). *Ecological Entomology* 30(6): 673-683.
- Cheng X., Baumgartner K. (2004). Arbuscular mycorrhizal fungi-mediated nitrogen transfer from vineyard cover crops to grapevines. *Biology and Fertility of Soils* 40(6): 406-412.
- Chesson P.L., Pacala S., Neuhauser C. (2002). Environmental niches and ecosystem functioning. In *Functional consequences of biodiversity*, (Kinzig A. et al., eds.), Princeton University Press, Princeton: 213-245.
- Chevallier A. (2006). Lutte biologique par conservation et gestion des habitats : étude des éléments fixes du paysage comme réservoirs d'auxiliaires des cultures. Mémoire d'Ingénieur Agronome ENSAT, Institut National Polytechnique, Toulouse, 69 p.
- Chilliard Y., Bauchart D., Lessire M., Schmidely P., Mourou J. (2008). Qualité des produits : modulation par l'alimentation des animaux de la composition en acides gras du lait et de la viande. *INRA Productions animales* 21(Numéro spécial Anniversaire «20 ans de recherche en productions animales à l'INRA»): 95-106.
- Chilliard Y., Ferlay A., Doreau M. (2001). Effect of different types of forages, animal fat or marine oils in cow's diet on milk fat secretion and composition, especially conjugated linoleic acid (CLA) and polyunsaturated fatty acids. *Livestock Production Science* 70: 31-48.
- Chilliard Y., Ferlay A., Mansbridge R.M., Doreau M. (2000). Ruminant milk fat plasticity: nutritional control of saturated, polyunsaturated, *trans* and conjugated fatty acids. *Annales de zootechnie* 49(3): 181-205.
- Chilliard Y., Glasser F., Ferlay A., Bernard L., Rouel J., Doreau M. (2007). Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *European journal of lipid science and technology* 109(8): 828-855.
- Chirico J., Wikteliuss S., Waller P.J. (2003). Dung beetle activity and the development of trichostrongylid eggs into infective larvae in cattle faeces. *Veterinary Parasitology* 118(1-2): 157-163.
- Chittka L., Thompson J.D. (2001). Cognitive ecology of pollination: Animal behaviour and floral evolution, University Press, Cambridge UK, 344 p.
- Chiverton P.A. (1986). Predator density manipulation and its effects on populations of *Rhopalosiphum padi* (Hom.: Aphididae) in spring barley. *Annals of Applied Biology* 109(1): 49-60.
- Chu G.X., Shen Q.R., Cao J.L. (2004). Nitrogen fixation and transfer from peanut to rice cultivated in aerobic soil in an intercropping system and its effect on soil N fertility. *Plant and Soil* 263: 17-27.
- Ciotola M., Watson, A.K, Hallet, S.G, (1995). Discovery of an isolate of *Fusarium oxysporum* with potential to control *Striga hermonthica* in Africa. *Weed Research* 35: 303-309.

- Cisneros J.J., Rosenheim J.A. (1997). Ontogenetic change of prey preference in the generalist predator *Zelus renardii* and its influence on predator-predator interactions. *Ecological Entomology* 22(4): 399-407.
- Clapham W.M., Foster J.G., Neel J.P.S., Fedders J.M. (2005). Fatty acid composition of traditional and novel forages. *Journal of Agricultural and Food Chemistry* 53: 10068-10073.
- Clapperton M.J., Lee N.O., Binet F., Conner R.L. (2001). Earthworms indirectly reduce the effects of take-all (*Gaeumannomyces graminis* var. *tritici*) on soft white spring wheat (*Triticum aestivum* cv. Fielder). *Soil Biology & Biochemistry* 33(11): 1531-1538.
- Clavien Y., Delabays N. (2006). Inventaire floristique des vignes de Suisse romande: connaître la flore pour mieux la gérer. *Revue suisse de Viticulture, d'Arboriculture et d'Horticulture* 38(6): 335-341.
- Clermont-Dauphin C. (1995). Fertilisation minérale d'une association Haricot-Mais en zone tropicale humide. Diagnostic des effets des systèmes de culture. Thèse, INA Parie Grignon, 2997 p.
- Clifford P.T.P., Scott D. (1989). Inflorescence, bumble bee, and climate interactions in seed crops of a tetraploid red clover (*Trifolium pratense* L.). *Journal of Applied Seed Production* 7: 38-45.
- Colbach N., Debaeke P., Orlando, D. (1998). Integrating crop management and crop rotation effects into models of weed population dynamics: a review. *Weed Science* 46(6): 717-728.
- Cole L., Dromph K.M., Boaglio V., Bardgett R.D. (2004). Effect of density and species richness of soil mesofauna on nutrient mineralisation and plant growth. *Biology and fertility of Soils* 39(5): 337-343.
- Coleman D.C., Whitman W.B. (2005). Linking species richness, biodiversity and ecosystem function in soil systems. *Pedobiologia* 49(6): 479-497.
- Coleno F.C., Duru M. (1998). Gestion de production en systèmes d'élevage utilisateurs d'herbe. Une approche par atelier. *Etudes et Recherches sur les Systèmes Agraires et le Développement* 31: 45-61.
- Colin M.E., Faucon J.P., Heinrich A., Ferry R., Giauffret A. (1983). Etude du premier foyer français de varrobose de l'abeille. *Bull. Acad. Vét. France* 56: 89-93.
- Coll M., Bottrell D.G. (1995). Predator-prey association in mono- and dicultures: effect of maize and bean vegetation. *Agriculture, Ecosystems & Environment* 54(1-2): 115-125.
- Coll M., Bottrell D.G. (1996). Movement of an insect parasitoid in simple and diverse plant assemblages. *Ecological Entomology* 21(2): 141-149.
- Collinge W., E (1913). The destruction and dispersal of weeds by wild birds. *Jour. Board Agr. (Great Britain)* 20: 15-26.
- Collins K.L., Boatman N.D., Wilcox A., Holland J.M. (2003). A 5-year comparison of overwintering polyphagous predator densities within a beetle bank and two conventional hedgebanks. *Annals of Applied Biology* 143(1): 63-71.
- Collins K.L., Boatman N.D., Wilcox A., Holland J.M. (2003). Effects of different grass treatments used to create overwintering habitat for predatory arthropods on arable farmland. *Agriculture, Ecosystems & Environment* 96(1-3): 59-67.
- Collins K.L., Boatman N.D., Wilcox A., Holland J.M., Chaney K. (2002). Influence of beetle banks on cereal aphid predation in winter wheat. *Agriculture, Ecosystems & Environment* 93(1-3): 337-350.
- Collins K.L., Wilcox A., Chaney K., Boatman N.D., Holland J.M. (1997). The influence of beetle banks on aphid population predation in winter wheat. *Aspects of Applied Biology* 50: 341-346.
- Collins S.L. (1995). The measurement of stability in grasslands. *Trends in Ecology and Evolution* 10(3): 95-96.
- Collomb M., Butikofer U., Maurer J., Sieber R. (2006). Composition en acides gras du lait de brebis produit à diverses altitudes. *Revue Suisse d'Agriculture* 38: 335-339.
- Collomb M., Butikofer U., Sieber R., Jeangros B., Bosset J.O. (2002). Correlation between fatty acids in cows' milk fat produced in the Lowlands, Mountains and Highlands of Switzerland and botanical composition of the fodder. *International Dairy Journal* 12(8): 661-666.
- Collomb M., Butikofer U., Spahni M., Jeangros B., Bosset J.O. (1999). Fatty acid and glyceride composition of cow's milk fat in highland lowland regions. *Sciences des Aliments* 19: 97-110.
- Commission des communautés européennes (2006). Enrayer la diminution de la biodiversité à l'horizon 2010 et au-delà - Préserver les services écosystémiques pour le bien-être humain: 17 p. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=COM:2006:0216:FIN:FR:PDF>
- Connor D.J. (2001). Optimizing crop diversification. In *Crop science: progress and prospects*, (Nosberger J. et al., eds.), Cab International, Bristol, UK: 191-211.
- Connor E.F., McCoy E.D. (1979). The statistics and biology of the species-area relationship. *American Naturalist* 113(6): 791-833.
- Cook R., Baker K.F. (1983). The nature and practice of biological control of plant pathogens, Am. Phytopathol. Soc., St Paul, Minnesota, 539 p.
- Coomes D.S., Sotherton N.W. (1986). The dispersal and distribution of polyphagous predatory Coleoptera in cereals. *Annals of Applied Biology* 108(3): 461-474.
- Copeland P.J., Crookston R.K. (1992). Crop sequence affects nutrient composition of corn and soybean grown under high fertility. *Agronomy Journal* 84: 503-509.
- Corbet S.A. (1997). Role of pollinators in species preservation, conservation, ecosystem stability and genetic diversity, *Acta Horticulturae*, 437: 219-229.
- Corbett A. (1998). The importance of movement in the response of natural enemies to habitat manipulation. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Pickett C.H., Bugg R.L., eds.), University of California Press, Berkeley - Los Angeles - London: 25-48.
- Corre-Hellou G., Dibet A., Aveline A., Crozat Y. (2004). Le pois dans des systèmes à faibles intrants : culture pure ou associée ? *Perspectives Agricoles* 306: 8-10.
- Côrtes C., Damasceno J.C., Jamot J., Prache S., Aulagnier S. (2006). Ewes increase their intake when offered a choice of herbage species at pasture. *Animal Science* 82: 183-191.
- Cortet J., Joffre R., Elmholt S., Coeurdassier M., Scheiffler R., Krogh P.H. (2006). Interspecific relationships among soil invertebrates influence pollutant effects of phenanthrene. *Environmental Toxicology and Chemistry* 25(1): 120-127.

- Cortet J., Ronce D., Poinot-Balaguer N., Beaufreton C., Chabert A., Viaux P., de Fonseca J.P.C. (2002). Impacts of different agricultural practices on the biodiversity of microarthropod communities in arable crop systems. *European Journal of Soil Biology* 38(3-4): 239-244.
- Cortez J., Garnier E., Pérez-Harguindeguy N., Debussche M., Gillon D. (2007). Plant traits, litter quality and decomposition in a Mediterranean old-field succession. *Plant and Soil* 296: 19-34.
- Cosentino D., Chenu C., Bissonnais L.Y. (2006). Aggregate stability and microbial community dynamics under drying-wetting cycles in a silt loam soil. *Soil Biology & Biochemistry* 38(8): 2053-2062.
- Costamagna A.C., Landis D.A. (2006). Predators exert top-down control of soybean aphid across a gradient of agricultural management systems. *Ecological Applications* 16(4): 1619-1628.
- Costamagna A.C., Landis D.A., Difonzo C.D. (2007). Suppression of soybean aphid by generalist predators results in a trophic cascade in soybeans. *Ecological Applications* 17(2): 441-451.
- Costanza R., Arge R.d., Groot R.d., Farber S., Grasso M., Hannon B., Limburg K., Naeem S., O'Neill R.V., Paruelo J., Raskin R.G., Sutton P., Belt M.v.d. (1997). The value of the world's ecosystem services and natural capital. *Nature* 387(6630): 253-260.
- Costello M., Altieri M.A. (1995). Abundance, growth rate and parasitism of *Brevicoryne brassicae* and *Myzus persicae* (Homoptera: Aphididae) on broccoli grown in living mulches. *Agriculture, Ecosystems & Environment* 52(2-3): 187-196.
- Cottingham K.L., Brown B.L., Lennon J.T. (2001). Biodiversity may regulate the temporal variability of ecological systems. *Ecology Letters* 4: 72-85.
- Cottrell T.E. (2005). Predation and cannibalism of lady beetle eggs by adult lady beetles. *Biological Control* 34(2): 159-164.
- Coulon J.B., Delacroix-Buchet A., Martin B., Pirisi A. (2005). Facteurs de production et qualité sensorielle des fromages. *INRA Productions animales* 18: 49-62.
- Coulon J.B., Priolo A. (2002). La qualité sensorielle des produits laitiers et de la viande dépend des fourrages consommés par les animaux. *INRA Productions animales* 15: 333-342.
- Courault D., Drobinski P., Brunet Y., Lacarrère P., Talbot C. (2007). Impact of surface heterogeneity on a buoyancy-driven convective boundary layer. *Boundary Layer Meteorology* 127(3): 383-403.
- Cousens R., Mortimer M. (1995). Dynamics of weed population, Cambridge University Press, Cambridge, 332 p.
- Couteaux M.M., Monrozier L.J., Bottner P. (1996). Increased atmospheric CO₂: Chemical changes in decomposing sweet chestnut (*Castanea sativa*) leaf litter incubated in microcosms under increasing food web complexity. *Oikos* 76(3): 553-563.
- Cowger C., Mundt C.C. (2002). Aggressiveness of *Mycosphaerella graminicola* isolates from susceptible and partially resistant wheat cultivars. *Phytopathology* 92(6): 624-630.
- Cox-Foster D.L., Conlan S., Holmes E.C., Palacios G., Evans J.D., Moran N.A., Quan P.L., Briese T., Hornig M., Geiser D.M., Martinson V., vanEngelsdorp D., Kalkstein A.L., Drysdale A., Hui J., Zhai J., Cui L., Hutchinson S.K., Simons J.F., Egholm M., Pettis J.S., Lipkin W.I. (2007). A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* 318: 283-287.
- Cragg J.B., Bardgett R.D. (2001). How changes in soil faunal diversity and composition within a trophic group influence decomposition processes. *Soil Biology & Biochemistry* 33: 2073-2082.
- Craine J.M., Wedin D.A., Chapin F.S.I., Reich P.B. (2003). Relationship between the structure of root systems and resource use for 11 North American grassland plants. *Plant Ecology* 165: 85-100.
- Craine J.M., Wedin D.A., Reich P.B. (2001). The response of soil CO₂ flux to changes in atmospheric CO₂, nitrogen supply and plant diversity. *Global Change Biology* 7: 947-953.
- Cremene C., Groza G., Rakosy L., Schileyko A.A., Baur A., Erhardt A., Baur B. (2005). Alterations of steppe-like grasslands in Eastern Europe: a threat to regional biodiversity hotspots. *Conservation Biology* 19(5): 1606-1618.
- Croft B.A. (1982). Management of apple orchard weeds to improve biological control of spider mites, *Annual Meeting of Weed Science Society of America*, 257: 134 p.
- Cromartie W.J. (1991). The environmental control of insects using crop diversity. In CRC handbook of pest management in agriculture, (Pimentel D., ed.), 1, CRC Press, Boca Raton, FL: 223-251.
- Cross J.V., Solomon M.G., Babandriker D., Blommers L., Easterbrook M.A., Jay C.N., Jenser G., Jolly R.L., Kuhlmann U., Lilley R., Olivella E., Toepfer S., Vidal S. (1999). Biocontrol of pests of apples and pears in northern and central Europe: 2. Parasitoids. *Biocontrol Science and Technology* 9(3): 277-314.
- Crovetto Lamarca C. (2000). Les fondements d'une agriculture durable, Teknea, 300 p.
- Crozier P. (2004). Influence de l'enherbement sur le vignes et les vins de pinots noirs en Bourgogne, *AFPP - Dix neuvième Conférence du COLUMA Journées Internationales sur la lutte contre les Mauvaises Herbes*, Dijon, France, CD ROM.
- Crutsinger G.M., Collins M.D., Fordyce J.A., Gompert Z., Nice C.C., Sanders N.J. (2006). Plant genotypic diversity predicts community structure and governs an ecosystem process. *Science* 313: 966-968.
- Cruz P., Duru M., Théron O., Theau J.P., Ducourtieux C., Jouany C., Al Haj Khaled R., Ansquer P. (2002). Une nouvelle approche pour caractériser les prairies naturelles et leur valeur d'usage. *Fourrages* 172: 335-354.
- Cuddeford V. (1999). Beneficial bugs at risk from pesticides, Wild Worldlife Fund, Toronto, Canada, <http://www.ca/satellite/prip/resources/bugs-at-risk.pdf> : 68 p.
- Currie C.R., Spence J.R., Niemela J. (1996). Competition, cannibalism and intraguild predation among ground beetles (Coleoptera: Carabidae): a laboratory study. *Coleopterists Bulletin* 50(2): 135-148.
- Curry J.P., Boyle K.E., Farrell E.P. (1989). The invertebrate fauna of reclaimed cutaway peat in central Ireland and its influence on soil fertility and plant growth. *Agriculture, Ecosystems & Environment* 27(1-4): 217-225.
- Curtis I.S., He C., Scott R., Power J.B., Davey M.R. (1996). Genomic male sterility in lettuce, a baseline for the production of F1 hybrids. *Plant Science* 113: 113-119.
- Da Silva Pontes L., Soussana J.F., Louault F., Andueza D., Carrère P. (2007). Leaf traits affect the above-ground productivity and quality of grasses. *Functional Ecology* 21: 844-853.
- Daboussi M. (1996). Fungal transposable elements: generators of diversity and genetic tools. *Journal of Genetics* 75(3): 325-339.

- Daccord R., Wyss U., Kessler J., Arrigo Y., Rouel M., Lehman J., Jeangros B. (2006). Valeur nutritive des fourrages. In Apports alimentaires recommandés et tables de la valeur nutritive des aliments pour les ruminants, Station de recherche Agroscope Liebefeld-Posieux ALP, Posieux, Suisse: 18 p.
- Dag A., Zipori I., Pleser Y. (2006). Using bumblebees to improve almond pollination by the honeybee. *Journal of Apicultural Research* 45(3): 215-216.
- Dalmastri C., Chiarini L., Cantale C., Bevivino A., Tabacchioni S. (1999). Soil type and maize cultivar affect the genetic diversity of maize root-associated Burkholderia cepacia populations. *Microbial Ecology* 38(3): 273-284.
- Dambreville C., Hallet S., Nguyen C., Morvan T., Germon J.C., Philippot L. (2006). Structure and activity of the denitrifying community in a maize-cropped field fertilized with composted pig manure or ammonium nitrate. *FEMS Microbiology Ecology* 56(1): 119-131.
- Daniel T.C. (2001). Whither scenic beauty? Visual landscape quality assessment in the 21st century. *Landscape and Urban Planning* 54: 267-281.
- D'Antonio C.M. (2000). Fire, plant invasions, and global changes. In *Invasive species in a changing world*, (Mooney H.A., Hobbs R.J., eds.), Island Press, Covela: 65-94.
- D'Antonio C.M., Vitousek P.M. (1992). Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23: 63-87.
- Darwin C.H. (1862). On the various contrivances by which British and foreign orchids are fertilized by insects, and on the good effects of intercrossing, John Murray, London UK.
- Davidson D.A., Grieve I.C. (2006). Relationships between biodiversity and soil structure and function: Evidence from laboratory and field experiments. *Applied Soil Ecology* 33(2): 176-185.
- Davis M.A., Grime J.P., Thompson K. (2000). Fluctuating resources in plant communities: a general theory of invasibility. *Journal of Ecology* 88: 528-534.
- Dawson L.A., Duff E.I., Campbell C.D., Hirst D.J. (2001). Depth distribution of cherry (*Prunus Avium* L.) tree roots as influenced by grass root competition. *Plant and Soil* 231: 11-19.
- Daza-Bustamante P., Fuentes-Contreras E., Rodriguez L.C., Figueroa C.C., Niemeyer H.M. (2002). Behavioural differences between *Aphidius ervi* populations from two tritrophic systems are due to phenotypic plasticity. *Entomologia Experimentalis et Applicata* 104(2/3): 321-328.
- de Chazal J. (2003). Perspectives on Ecosystem Health. PhD Thesis, PhD Thesis, Australian National University, Canberra, Australia.
- De Deyn G.B., Cornelissen H.C., Bardgett R.D. (2008). Plant functional traits and soil carbon sequestration in contrasting biomes. *Ecology Letters* 11(5): 516-531.
- de Deyn G.B., Raaijmakers C.E., van Ruijven J., Berendse F., van der Putten W.H. (2004). Plant species identity and diversity effects on different trophic levels of nematodes in the soil food web. *Oikos* 106(3): 576-586.
- de Deyn G.B., Raaijmakers C.E., Zoomer H.R., Berg M.P., de Ruiter P.C., Verhoef H.A., Bezemer T.M., van der Putten W.H. (2003). Soil invertebrate fauna enhances grassland succession and diversity. *Nature* 422: 711-713.
- de la Torre A., Debiton E., Durand D., Chardigny J.M., Berdeaux O., Barthomeuf C., Bauchart D., Gruffat D. (2005). Conjugated linoleic acid isomers and their conjugated derivatives inhibit growth of human cancer cell lines. *Anticancer Research* 25: 3943-3950.
- de la Torre A., Debiton E., Durand D., Juanéda P., Durand D., Chardigny J.M., Barthomeuf C., Bauchart D., Gruffat D. (2006). Beef conjugated linoleic acid isomers reduce human cancer cell growth even when associated to other beef fatty acids. *British Journal of Nutrition* 95: 346-352.
- de Moraes C.M., Cortesero A.M., Stapel J.O., Lewis W.J. (1999). Intrinsic and extrinsic competitive interactions between two larval parasitoids of *Heliothis virescens*. *Ecological Entomology* 24(4): 402-410.
- de Oliveira D., Pion S., Paradis R.O. (1984). Pollination and the yield of apples. *Technical Bulletin, Research Station, Saint-Jean-sur-Richelieu, Quebec* 19: 51-61.
- Deacon L.J., Pryce-Miller E.J., Frankland J.C., Bainbridge B.W., Moore P.D., Robinson C.H. (2006). Diversity and function of decomposer fungi from a grassland soil. *Soil Biology & Biochemistry* 38(1): 7-20.
- Deak A., Hall M.H., Sanderson M.A., Archibald D.D. (2007). Production and nutritive value of grazed simple and complex forage mixtures. *Agronomy Journal* 99(3): 814-821.
- Debaeke P., Orlando D. (1991). Simplification du travail du sol et évolution de la flore adventice : conséquences pour le désherbage à l'échelle de la rotation. In *Simplification du travail du sol.*, (INRA, ed.): 35-62.
- Debras J.F., Rieux R., Arcier F.F.d., Kretschmar A., Simon S. (1998). Regulation of pear psyllid: importance and role of the orchard environment. *Phytoma, la défense des végétaux* 510: 53-55.
- Debras J.-F., Dussaud A., Rieux R., Dutoit T. (2007). Recherche prospective sur le rôle " source " des haies en production fruitière intégrée. Le cas des perce-oreilles : *Forficula auricularia* L. et *Forficula pubescens* Gené. *Comptes rendus de l'Académie des Sciences* 330(9): 664-673.
- Debry G. (2001). Lait, nutrition et santé, Tec & Doc, Paris, France, 566 p.
- Debussche M., Rambal S., Lepart J. (1987). Les changements de l'occupation des terres en région méditerranéenne humide: évaluation des conséquences hydrologiques. *Acta Oecologica* 8(4): 317-332.
- Decaens T., Dutoit T., Alard D. (1997). Earthworm community characteristics during afforestation of abandoned chalk grasslands (Upper Normandy, France). *European Journal of Soil Biology* 33(1): 1-11.
- Decaens T., Dutoit T., Alard D., Lavelle P. (1998). Factors influencing soil macrofaunal communities in post-pastoral successions of western France. *Applied Soil Ecology* 9(1-3): 361-367.
- Decaens T., Galvis J.H., Amézquita E. (2001). Propriétés des structures produites par les ingénieurs écologiques à la surface du sol d'une savane colombienne. *Comptes rendus de l'Académie des sciences. Série 3, Sciences de la vie* 324(5): 465-478.
- Decaens T., Jimenez J.J., Gioia C., Measey G.J., Lavelle P. (2006). The values of soil animals for conservation biology. *European Journal of Soil Biology* 42: S23-S38.
- Decaens T., Lavelle P., Jimenez J.J. (2008). Priorities for conservation of soil animals. *CAB Reviews: perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 3: 1-18.

- Decaens T., Rangel A.F., Asakawa N., Thomas R.J. (1999). Carbon and nitrogen dynamics in ageing earthworm casts in grasslands of the eastern plains of Colombia. *Biology and Fertility of Soils* 30(1-2): 20-28.
- Decourtye A., Devillers J., Cluzeau S., Charretton M., Pham-Delègue M. (2004). Effect of Imidacloprid and Deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. *Ecotoxicology and Environmental Safety* 57: 410-419.
- Decourtye A., Lecompte P., Pierre J., Chauzat M.-P., Thiébeau P. (2007). Introduction de jachères florales en zones de grandes cultures : comment mieux concilier agriculture, biodiversité et apiculture ? . *Le Courrier de l'environnement de l'INRA* 54: 33-56.
- Dedieu B., Louault F., Tournadre H., Benoit M. (2008). Réponse de systèmes d'élevage innovants à la variabilité climatique : une expérimentation en production extensive ovin viande intégrant des préoccupations environnementales. In *Les exploitations d'élevage en mouvement. Flexibilités et dynamiques face aux aléas et aux nouveaux enjeux des filières et territoires*, (Dedieu B. *et al.*, eds.), Editions Quae, Paris: 161-178.
- Degrandi-Hoffman G., Watkins J.C. (2000). The foraging activity of honey bees *Apis mellifera* and non-*Apis* bees on hybrid sunflowers (*Helianthus annuus*) and its influence on cross-pollination and seed set. *Journal of Apicultural Research* 39: 37-45.
- Deiglmayr K., Philippot L., Kandeler E. (2006). Functional stability of the nitrate-reducing community in grassland soils towards high nitrate supply. *Soil Biology & Biochemistry* 38(9): 2980-2984.
- Delabays N., Mermilod G., de Joffrey J.P., Bohren C. (2004). Demonstration in cultivated fields of the reality of the phenomenon of allelopathy, 12. *Colloque International sur la Biologie des Mauvaises Herbes*, Dijon (France), AFPP: 97-105.
- Delabays N., Mermilod G., Bohren C. (2007). Plantes indésirables dans les jachères florales : résultats d'un réseau national d'observation. *Revue suisse Agriculture* 39(4): 199-203.
- Delaplane K.S., Mayer D.F. (2000). *Crop pollination by bees*, CABI Publishing, New York, 360 p.
- Delorme H. (2004). *La politique agricole commune : anatomie d'une transformation*, Presses de Sciences Po, Paris, 402 p.
- Delpuech J.M., Bardon C., Bouletreau M. (2005). Increase of the behavioral response to kairomones by the parasitoid Wasp *Leptopilina heterotoma* surviving insecticides. *Archives of Environmental Contamination and Toxicology* 49(2): 186-191.
- Delpuech J.M., Meyet J. (2003). Reduction in the sex ratio of the progeny of a parasitoid wasp (*Trichogramma brassicae*) surviving the insecticide chlorpyrifos. *Archives of Environmental Contamination and Toxicology* 45(2): 203-208.
- Denison R.F., Bryant D.C., Kearney T.E. (2004). Crop yields over the first nine years of LTRAS, a long-term comparison of field crop systems in a Mediterranean climate. *Field Crops Research* 86: 267-277.
- Dennis P., Aspinall R.J., Gordon I.J. (2002). Spatial distribution of upland beetles in relation to landform, vegetation and grazing management. *Basic and Applied Ecology* 3(2): 183-193.
- Dennis P., Fry G.L.A. (1992). Field margins: can they enhance natural enemy population densities and general arthropod diversity on farmland? *Agriculture, Ecosystems & Environment* 40(1-4): 95-115.
- Dennis P., Wratten S.D. (1991). Field manipulation of populations of individual staphylinid species in cereals and their impact on aphid populations. *Ecological Entomology* 16(1): 17-24.
- Denno R.F., Mitter M.S., Langellotto G.A., Gratton C., Finke D.L. (2004). Interactions between a hunting spider and a web-builder: consequences of intraguild predation and cannibalism for prey suppression. *Ecological Entomology* 29(5): 566-577.
- Denys C., Tschamtké T. (2002). Plant-insect communities and predator-prey ratios in field margin strips, adjacent crop fields, and fallows. *Oecologia* 130(2): 315-324.
- Depret G., Houot S., Allard M.R., Breuil M.C., Nouaim R., Laguerre G. (2004). Long-term effects of crop management on *Rhizobium leguminosarum* biovar *viciae* populations. *FEMS Microbiology Ecology* 51(1): 87-97.
- Derner J.D., Hart R.H. (2007). Grazing-induced modifications to peak standing crop in Northern mixed-grass prairie. *Rangeland Ecology & Management* 60(3): 270-276.
- Deruiter P.C., Vanveen J.A., Moore J.C., Brussaard L., Hunt H.W. (1993). Calculation of Nitrogen Mineralization in Soil Food Webs. *Plant and Soil* 157(2): 263-273.
- Desneux N., Decourtye A., Delpuech J.M. (2007). The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology* 52: 81-106.
- Desneux N., Denoyelle R., Kaiser L. (2006). A multi-step bioassay to assess the effect of the deltamethrin on the parasitic wasp *Aphidius ervi*. *Chemosphere* 65(10): 1697-1706.
- Deutsch B., Paulian M., Thierry D., Canard M. (2005). Quantifying biodiversity in ecosystems with green lacewing assemblages. *Agronomy for Sustainable Development* 25(3): 337-343.
- Devers M., Henry S., Hartmann A., Martin-Laurent F. (2005). Horizontal gene transfer of atrazine-degrading genes (*atz*) from *Agrobacterium tumefaciens* St96-4 pADP1 : Tn5 to bacteria of maize-cultivated soil. *Pest Management Science* 61(9): 870-880.
- Dewar A.M., Haylock L.A., Garner B.H., Sands R.J.N., Baker P. (2005). The environmental impact of GM herbicide-tolerant sugarbeet. *Aspects of Applied Biology* 74: 119-126.
- Di Falco S., Chavas J.P. (2006). Crop genetic diversity, farm productivity and the management of environmental risk in rainfed agriculture. *European Review of Agricultural Economics* 33(3): 289-314.
- Di Falco S., Chavas J.P., Smale M. (2007). Farmer management of production risk on degraded lands: the role of wheat variety diversity in the Tigray region, Ethiopia. *Agricultural Economics* 36(2): 147-156.
- Di Pietro J.P., Caillaud C.M., Chaubet B., Pierre J.S., Trottet M. (1998). Variation in resistance to the grain aphid, *Sitobion avenae* (Sternorhynca: Aphididae), among diploid wheat genotypes : multivariate analysis of agronomic data. *Plant Breeding* 117: 407-412.
- Díaz S., Cabido M. (2001). Vive la différence: plant functional diversity matters to ecosystem processes. *Trends in Ecology & Evolution* 16(11): 646-655.
- Díaz S., Fargione J., Chapin F.S., Tilman D. (2006). Biodiversity loss threatens human well-being. *PLoS Biology* 4(8): e277.
- Díaz S., Lavorel S., De Bello F., Quétier F., Grigulis K., Robson T.M. (2007). Incorporating plant functional diversity effects in ecosystem service assessments. *Proceedings of the National Academy of Sciences* 104: 20684-20689.
- Díaz S., Symstad A.J., Chapin III F.S., Wardle D.A., Huenneke L.F. (2003). Functional diversity revealed by removal experiments. *Trends in Ecology and Evolution* 18: 140-146.

- Diaz S., Tilman D., Fargione J., Chapin III F.S., Dirzo R., Kitzberger T., Gemmill B., Zobel M., Vilà M., Mitchell C., Wilby A., Daily G.C., Galetti M., Laurance W.F., Pretty J., Naylor R., Power A., Harvell A., Potts S., Kremen C., Griswold T., Eardley C. (2005). Biodiversity Regulation of Ecosystem Services. *In Ecosystems and human well-being: Current state and trends*, (Assessment M.E., ed.), Island Press, Washington, D.C.: 297-329.
- Diaz-Zorita M., Perfect E., Grove J.H. (2002). Disruptive methods for assessing soil structure. *Soil & Tillage Research* 64: 3-22.
- Dicke M., Takabayashi J., Posthumus M.A., Schutte C., Krips O.E. (1998). Plant-phytoseiid interactions mediated by herbivore-induced plant volatiles: variation in production of cues and in responses of predatory mites. *Experimental & Applied Acarology* 22(6): 311-333.
- Dickie I.A. (2007). Host preference, niches and fungal diversity. *New Phytologist* 174(2): 230-233.
- Diden W., Rombke J. (2001). Enchytraeids as indicator organisms for chemical stress in terrestrial ecosystems. *Ecotoxicology and Environmental Safety* 50(1): 25-43.
- Diehl S., Feissel M. (2000). Effects of enrichment on three-level food chains with omnivory. *American Naturalist* 155(2): 200-218.
- Dilly O., Bloem J., Vos A., Munch J.C. (2004). Bacterial diversity in agricultural soils during litter decomposition. *Applied and Environmental Microbiology* 70(1): 468-474.
- Dimitrakopoulos A.P., Schmid B. (2004). Biodiversity effects increase linearly with biotope space. *Ecology Letters* 7: 574-583.
- Dinter A. (2002). Microcosm studies on intraguild predation between female erigonid spiders and lacewing larvae and influence of single versus multiple predators on cereal aphids. *Journal of Applied Entomology* 126(5): 249-257.
- Dix N.J., Webster J. (1995). *Fungal Ecology*, 1st ed., Chapman & Hall, London, UK, 499 p.
- Dobremez L., Josien E., Camacho O., Andrieu N. (2008). La sécurisation des systèmes fourragers et la réponse aux enjeux agri-environnementaux. *In Les exploitations d'élevage en mouvement. Flexibilités et dynamiques face aux aléas et aux nouveaux enjeux des filières et territoires* (Dedieu B., Chia, E., Moulin, C.H., Leclerc, B., Tichit, M., ed.), QUAE, Paris: 225-238.
- Dodd M., Silvertown J., McConway K., Potts J., Crawley M. (1995). Community stability : a 60-year record of trends and outbreaks in the occurrence of species in the Park Grass experiment. *Journal of Ecology* 83(2): 277-285.
- Don R. (1997). Weed seed contaminants in cereal seed, *The 1997 Brighton Crop Conference - Weeds*, Brighton (UK), British Crop Protection Council 255-262.
- Doreau M., Lee M.R.F., Ueda K., Scollan N.D. (2005). Métabolisme ruminal et digestibilité des acides gras des fourrages, *12. Rencontres Recherches Ruminants*: 101.
- Dormaar J.F., Naeth M.A., Willms W.D., Chanasyk D.S. (1995). Effect of native prairie, crested wheatgrass (*Agropyron cristatum* (L.) Gaertn.) and Russian wildrye (*Elymus junceus* Fisch.) on soil chemical properties. *Journal of Range Management* 48: 258-263.
- Dormann C.F., Schweiger O., Arens P., Augenstein I., Aviron S., Bailey D., Baudry J., Billeter R., Bugter R., Bukacek R., Burel F., Cerny M., de Cock R., de Blust G., DeFilippi R., Diekötter T., Dirksen J., Durka W., Edwards P.J., Frenzel M., Hamersky R., Hendrickx F., Herzog F., Klotz S., Koolstra B., Lausch A., Le Coeur D., Liira J., Maelfait J.P., Opdam P., Roubalova M., Schermann-Legionnet A., Schermann N., Schmidt T., Smulders M.J.M., Speelmans M., Simova P., Verboom J., van Wingerden W., Zobel M. (2008). Prediction uncertainty of environmental change effects on temperate European biodiversity. *Ecology Letters* 11(3): 235-244.
- Dorrepaal E. (2007). Are plant growth-form-based classifications useful in predicting northern ecosystem carbon cycling feedbacks to climate change? *Journal of Ecology* 95: 1167-1180.
- Dougherty M.C., Thevathasan N.V., Gordon A.M., Lee H., Kort J. (2007). Nitrate and E. Coli NAR analysis in tile drain effluent from a mixed tree intercrop and monocrop system, *When Trees and Crops Get Together. Proceedings of the 10th North American Agroforestry Conference*, Québec City, Canada, Campeau A.O.a.S., ed.
- Dover J., Sparks T., Clarke S., Gobbett K., Glossop S. (2000). Linear features and butterflies: the importance of green lanes. *Agriculture, Ecosystems & Environment* 80(3): 227-242.
- Dover J.W. (1997). Conservation headlands: effects on butterfly distribution and behaviour. *Agriculture Ecosystems & Environment* 63(1): 31-49.
- Downie J.A. (2007). Rhizobia. *In Encyclopedia of Life Sciences*, John Wiley & Sons, Ltd: 1-8.
- Dramstad W., Fry G. (1995). Foraging Activity of Bumblebees (*Bombus*) in Relation to Flower Resources on Arable Land. *Agriculture, Ecosystems & Environment* 53(2): 123-135.
- Drew E.A., Murray R.S., Smith S.E., Jakobsen I. (2003). Beyond the rhizosphere: growth and function of arbuscular mycorrhizal external hyphae in sands of varying pore size. *Plant and Soil* 251: 105-114.
- Drissner D., Blum H., Tschirko D., Kandeler E. (2007). Nine years of enriched CO₂ changes the function and structural diversity of soil microorganisms in a grassland. *European Journal of Soil Science* 58(1): 260-269.
- Duchemin M. (2007). Réduction de la pollution diffuse d'origine agricole à l'aide de bandes enherbées et arborées, *When Trees and Crops Get Together. Proceedings of the 10th North American Agroforestry Conference*, Québec City, Canada, Campeau A.O.a.S., ed.
- Duda J.J., Freeman D.C., Emlen J.M., Belnap J., Kitchen S.G., Zak J.C., Sobek E., Tracy M., Montante J. (2003). Differences in native soil ecology associated with invasion of the exotic annual chenopod, *Halogeton glomeratus*. *Biology and Fertility of Soils* 38(2): 72.
- Duelli P., Obrist M.K. (1998). In search of the best correlates for local organismal biodiversity in cultivated areas. *Biodiversity and Conservation* 7: 297-309.
- Duffield S. (1991). Does the size of area you spray with insecticides influence the extent of side-effects on invertebrate populations? *Game Conservancy Review* 22: 64-65.
- Duffield S.J., Bryson R.J., Young J.E.B., Sylvester-Bradley R., Scott R.K. (1997). The influence of nitrogen fertiliser on the population development of the cereal aphids *Sitobion avenae* (F.) and *Metopolophium dirhodum* (Wlk.) on field grown winter wheat. *Annals of Applied Biology* 130(1): 13-26.
- Duijff B.J., Recorbet G., Bakker P.A.H.M., Loper J.E., Lemanceau P. (1999). Microbial antagonism at the root level is involved in the suppression of *Fusarium* wilt by the combination of nonpathogenic *Fusarium oxysporum* Fo47 and *Pseudomonas putida* WCS358. *Phytopathology* 89(11): 1073-1079.
- Dukes J.S., Mooney H.A. (1999). Does global change increase the success of biological invaders? *Trends in Ecology and Evolution* 14(4): 135-139.

- Dukes J.S., Mooney H.A. (2004). Disruption of ecosystem processes in western North America by invasive species, *How landscapes change: human disturbances and ecosystem disruptions in the Americas*, Revista Chilena de Historia Natural, 77, Bradshaw G.A. et al., eds., Springer-Verlag: 411-437.
- Duncan A.J., Ginane C., Gordon I.J., Orskov E.R. (2003). Why do herbivores select mixed diets? , 6. *International Symposium on the nutrition of herbivores* ; , Mexico, 2003/10/19-24: 225-228.
- Dunning J.B., Danielson B.J., Pulliam H.R. (1992). Ecological processes that affect populations in complex landscapes. *Oikos* 65: 169-175.
- Dupraz C. (2005). Entre agronomie et écologie: vers la gestion d'écosystèmes cultivés. *Cahier d'étude DEMETER 2006*: 73-89.
- Dupraz C., Burgess P., Gavaland A., Graves A., Herzog F., Incoll L.D., Jackson N., Keesman K., Lawson G., Lecomte I., Liagre F., Mantzanas K., Mayus M., Moreno G., Palma J., Papanastasis V., Paris P., Pilbeam D.J., Reisner Y., Van Noordwijk M., Vincent G., Werf Van der W. (2005). Rapport final, Silvoarable Agroforestry For Europe (SAFE), European Research contract QLK5-CT-2001-00560. 254 p.
- Dupraz C., Liagre F. (2007). Innover en associant arbres et cultures: les atouts de l'agroforesterie moderne. *Forêt entreprise* 175: 56-60.
- Dupraz C., Liagre F. (2008). Agroforesterie, des arbres et des cultures, (Agricole E.F., ed.), 416.
- Durand D., Scislawski V., Gruffat D., Chilliard Y., Bauchart D. (2005). High-fat rations and lipid peroxidation in ruminants: Consequences on the health of animals and quality of their products. *In Indicators of Milk and Beef Quality*, (Hocquette J.F., Gigli S., eds.), Wageningen Academic, Wageningen, The Netherlands: 137-150.
- Duru M., Cruz P., Magda D. (2008). La conduite des couverts prairiaux, source de flexibilité. *In Les exploitations d'élevage en mouvement. Flexibilités et dynamiques face aux aléas et aux nouveaux enjeux des filières et territoires*, (Dedieu B. et al., eds.), Editions Quae, Paris: 57-71.
- Duru M., Gibon A. (1988). Prévoir la valeur nutritive des foin et des regains dans les Pyrénées centrales. I - Principaux facteur de variation de la composition chimique. *Fourrages* 114: 143-165.
- Duru M., Tallowin J., Cruz P. (2005). Functional diversity in low-input grassland farming systems: characterisation, effect and management, *Integrated efficient grassland farming and biodiversity*, Tartu, Estonia, Grassland Science in Europe, 10, Lillak R. et al., eds., European Grassland Federation: 199-210.
- Duso C., Malagnini V., Paganelli A., Aldegheri L., Bottini M., Otto S. (2004). Pollen availability and abundance of predatory phytoseiid mites on natural and secondary hedgerows. *Biocontrol* 49(4): 397-415.
- Duso C., Pasini M., Pellegrini M. (2003). Distribution of the predatory mite Typhlodromus pyri (Acari: Phytoseiidae) on different apple cultivars. *Biocontrol Science and Technology* 13(7): 671-681.
- Duthil J. (1973). *Eléments d'Ecologie et d'Agronomie*. Tome III. Exploitation et amélioration du milieu, Ed. Baillière, 656 p.
- Dyer L.A., Stireman J.O. (2003). Community-wide trophic cascades and other indirect interactions in an agricultural community. *Basic and Applied Ecology* 4(5): 423-432.
- Eastman J.L., Coughenour M.B., Pielke Sr. R.A. (2001). Does grazing affect regional climate? *Journal of Hydrometeorology* 2: 243-253.
- Edwards C.A., Brust G.E., Stinner B.R., McCartney D.A. (1992). Work in the United States on the use of cropping patterns to promote natural enemies of pests. *Aspects of Applied Biology* 31: 139-148.
- Ehler L.E. (1998). Invasion biology and biological control. *Biological Control* 13(2): 127-133.
- Ehlers B.K., Thompson J.D. (2004). Do co-occurring plant species adapt to one another ? The response of Bromus erectus to the presence of different Thymus vulgaris chemotypes. *Oecologia* 141: 511-518.
- Ehrenfeld J.G. (2003). Effects of exotic plants on soil nutrient cycling processes. *Ecosystems* 6: 503-523.
- Ehrenfeld J.G., Ravit B., Elgersma K. (2005). Feedbacks in the plant-soil system. *Annual Review of Ecology and Systematics* 30(1): 75-115.
- Eichhorn M., Paris P., Herzog F., Incoll L., Liagre F., Mantzanas K., Mayus M., Moreno G., Papanastasis V., Pilbeam D., Pisanelli A., Dupraz C. (2006). Silvoarable Systems in Europe : Past, Present and Future Prospects. *Agroforestry Systems* 67: 29-50.
- Ekesi S., Shah P.A., Clark S.J., Pell J.K. (2005). Conservation biological control with the fungal pathogen Pandora neoaphidis: implications of aphid species, host plant and predator foraging. *Agricultural and Forest Entomology* 7(1): 21-30.
- Elands B.H.M., Wiersum F.K. (2000). Forestry and rural development in Europe: an exploration of socio-political discourses. *Forest Policy and Economics* 3: 5-16.
- Elliott N.C., Kieckhefer R.W., Lee J.H., French B.W. (1999). Influence of within-field and landscape factors on aphid predator populations in wheat. *Landscape Ecology* 14(3): 239-252.
- Elmstrom K.M., Andow D.A., Barclay W.W. (1988). Flea-beetle movement in a broccoli monoculture and diculture. *Environmental Entomology* 17: 299-305.
- Elton C.S. (1958). *The Ecology of Invasions by Animals and Plants*, Methuen, London, 1996 p.
- Elzinga J.A., van Nouhuys S., van Leeuwen D.J., Biere A. (2007). Distribution and colonization ability of three parasitoids and their herbivorous host in a fragmented landscape. *Basic and Applied Ecology* 8: 75-88.
- English-Loeb G., Norton A.P., Walker M.A. (2002). Behavioral and population consequences of acarodomatia in grapes on phytoseiid mites (Mesostigmata) and implications for plant breeding. *Entomologia Experimentalis et Applicata* 104(2/3): 307-319.
- Entwistle J.C., Dixon A.F.G. (1989). The effect of augmenting grain aphid (Sitobion avenae) numbers in a field of winter wheat in spring on the aphid's abundance in summer and its relevance to the forecasting of outbreaks. *Annals of Applied Biology* 114(3): 397-408.
- Eriksson O., Wikstrom S., Eriksson A., Lindborg R. (2006). Species-rich Scandinavian grasslands are inherently open to invasion. *Biological Invasions* 8(2): 355-363.
- Ester A., van Rozen K. (2002). Earthworms (Aporrectodea spp.; Lumbricidae) cause soil structure problems in young Dutch polders. *European Journal of Soil Biology* 38(2): 181-185.
- Estes J.R., Amos B.B., Sullivan J.R. (1983). Pollination from two perspectives: the agricultural and biological sciences, *Handbook of experimental pollination biology*, (Jones C.E.L.R.J., ed.), 536-554.
- Eviner V.T., Chapin F.S.I. (2003). Biogeochemical interactions and biodiversity. *In Global change and human impacts*, (Melillo J.M. et al., eds.), Island Press, Washington: 151-173.
- Eviner V.T., Chapin F.S.I. (2005). Selective gopher disturbance influences plant species effects on nitrogen cycling. *Oikos* 109: 154-166.

- Eviner V.T., Chapin III F.S. (2003). Functional matrix: a conceptual framework for predicting multiple plant effects on ecosystem processes. *Annual Review of Ecology and Systematics* 34: 455-485.
- Faber J.H., Verhoef H.A. (1991). Functional differences between closely-related soil arthropods with respect to decomposition processes in the presence or absence of pine tree roots. *Soil Biology & Biochemistry* 23(1): 15-23.
- Faegri K., van der Pijl L. (1979). The principles of pollination ecology, Pergamon, Oxford.
- Fahrig L. (2003). Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology Evolution and Systematics* 34: 487-515.
- Fang M., Motavalli P.P., Kremer R.J., Nelson K.A. (2007). Assessing changes in soil microbial communities and carbon mineralization in Bt and non-Bt corn residue-amended soils. *Applied Soil Ecology* 37(1-2): 150-160.
- Fargione J., Brown C.S., Tilman D. (2003). Community assembly and invasion: An experimental test of neutral versus niche processes. *Proceedings of the National Academy of Sciences (U.S.A.)* 100: 8916-8920.
- Farruggia A., Martin B., Baumont S., Prache S., Doreau M., Hoste H., Durand D. (2008). Quels intérêts de la diversité floristique des prairies permanentes pour les ruminants et les produits animaux ? *INRA Productions Animales* 21(2): 181-200.
- Feldman T.S., Morris W.F., Wilson W.G. (2004). When can two plant species facilitate each other's pollination? *Oikos* 105(1): 197-207.
- Ferguson S.H., Joly D.O. (2002). Dynamics of springtail and mite populations: the role of density dependence, predation, and weather. *Ecological Entomology* 27(5): 565-573.
- Ferron P., Deguine J.P. (2005). Crop protection, biological control, habitat management and integrated farming. A review. *Agronomy for Sustainable Development* 25(1): 17-24.
- Filhol X., Baillet X. (1864). Expériences faites sur des lapins et sur des chiens avec le grain de *Lolium temulentum*. *Journal de Chimie Médicale de Pharmacie et de Toxicologie* VII(4): 395-397.
- Finch S., Collier R.H. (2000). Host-plant selection by insects – a theory based on 'appropriate/inappropriate landings' by pest insects of cruciferous plants. *Entomologia Experimentalis et Applicata* 96: 91-102.
- Finch S., Collier R.M. (2003). Insects can see clearly now the weeds have gone. *Biologist* 50(3): 132-135.
- Finckh M.R., Mundt C.C. (1992). Stripe Rust, Yield, and Plant Competition in Wheat Cultivar Mixtures. *Phytopathology* 82(9): 905-913.
- Finckh M.R., Wolfe M.S. (1998). Diversification strategies. In *The Epidemiology of Plant Diseases*, (Jones D.G., ed.), Chapman and Hall, London: 231-259.
- Finke D.L., Denno R.F. (2002). Intraguild predation diminished in complex-structured vegetation: implications for prey suppression. *Ecology* 83(3): 643-652.
- Finke D.L., Denno R.F. (2006). Spatial refuge from intraguild predation: implications for prey suppression and trophic cascades. *Oecologia* 149(2): 265-275.
- Finkes L.K., Cady A.B., Mulroy J.C., Clay K., Rudgers J.A. (2006). Plant-fungus mutualism affects spider composition in successional fields. *Ecology Letters* 9(3): 347-356.
- Fischer D.G., Hart S.C., Whitham T.G., Martinsen G.D., Keim P. (2004). Ecosystem implications of genetic variation in water-use of a dominant riparian tree. *Oecologia* 139: 288-297.
- Fitter A.H., Gilligan C.A., Hollingworth K., Kleczkowski A., Twyman R.M., Pitchford J.W. (2005). Biodiversity and ecosystem function in soil. *Functional Ecology* 19(3): 369-377.
- Fitzgerald J.D., Solomon M.G. (2004). Can flowering plants enhance numbers of beneficial arthropods in UK apple and pear orchards? *Biocontrol Science and Technology* 14(3): 291-300.
- Flexner J.L., Westigard P.H., Gonzalves P., Hilton R. (1991). The Effect of Groundcover and Herbicide Treatment on 2 Spotted Spider-Mite Density and Dispersal in Southern Oregon Pear Orchards. *Entomologia Experimentalis Et Applicata* 60(2): 111-123.
- Floate K.D., Wardhaugh K.G., Boxall A.B.A., Sherratt T.N. (2005). Fecal residues of veterinary parasiticides: Nontarget effects in the pasture environment. *Annual Review of Entomology* 50: 153-179.
- Flores J. (1997). Mediterranean vs. northern European meat products. Processing technologies and main differences. *Food Chemistry* 59(4): 505-510.
- Fontaine C., Collin C.L., Dajoz I. (2008). Generalist foraging of pollinators: diet expansion at high density. *Journal of Ecology* 96(5): 1002-1010.
- Fontaine C., Dajoz I., Meriguet J., Loreau M. (2006). Functional diversity of plant-pollinator interaction webs enhances the persistence of plant communities. *Plos Biology* 4(1): 129-135.
- Forman R.T.T., Baudry J. (1984). Hedgerows and hedgerow networks in landscape ecology. *Environmental Management* 8(6): 495-510.
- Fomara D.A., Tilman D. (2008). Plant functional composition influences rates of soil carbon and nitrogen accumulation. *Journal of Ecology* 96(2): 314-322.
- Fortuna M.A., Bascompte J. (2006). Habitat loss and the structure of plant-animal mutualistic networks. *Ecology Letters* 9(3): 281-286.
- Fortunel C., Garnier E., Joffre R., Kazakou E., Quested H., Grigulis K., Lavorel S., VISTA consortium (in press). Plant functional traits capture the effects of land use change and climate on litter decomposability of herbaceous communities in Europe and Israel. *Ecology*.
- Forup M.L., Memmott J. (2005). The relationship between the abundances of bumblebees and honeybees in a native habitat. *Ecological Entomology* 30(1): 47-57.
- Foster B.L., Smith V.H., Dickson T.L., Hildebrand T. (2002). Invasibility and compositional stability in a grassland community: relationship to diversity and extrinsic factors. *Oikos* 99: 300-307.
- Fournier E., Loreau M. (1999). Effects of newly planted hedges on ground-beetle diversity (Coleoptera, Carabidae) in an agricultural landscape. *Ecography* 22(1): 87-97.
- Fox J.W. (2005). Interpreting the 'selection effect' of biodiversity on ecosystem function *Ecology Letters* 8: 846-856.
- Fox T.B., Landis D.A., Cardoso F.F., Difonzo C.D. (2004). Predators suppress Aphis glycines Matsumura population growth in soybean. *Environmental Entomology* 33(3): 608-618.
- Fragoso C., Kanyonyo J., Moreno A., Senapati B.K., Blanchart E., Rodriguez C. (1999). A survey of tropical earthworms: taxonomy, biogeography and environmental plasticity. In *Earthworm management in tropical agroecosystems*, (Lavelle P. et al., eds.), CAB International: 27-55.

- Fraisse D., Carnat A., Viala D., Pradel P., Besle J.M., Coulon J.B., Felgines C., Lamaison J.L. (2007). Polyphenolic composition of a permanent pasture: Variations related to the period of harvesting. *Journal of the Science of Food and Agriculture* 87(13): 2427-2435.
- Frank D.A., Evans R.D. (1997). Effects of Native Grazers on Grassland N Cycling in Yellowstone National Park *Ecology* 78: 2238-2248.
- Frank D.A., Groffman P.M. (1998). Ungulate vs. Landscape Control of Soil C and N Processes in Grasslands of Yellowstone National Park *Ecology* 79: 2229-2241.
- Frank D.A., McNaughton S.W. (1991). Stability Increases with Diversity in Plant Communities: Empirical Evidence from the 1988 Yellowstone Drought *Oikos* 62: 360-362.
- Frank J., Volkmar W. (2006). *Episyrphus balteatus*: an example of pollination efficiency in hover flies, 91. Annual Meeting ESA, Memphis, Tennessee, August 2006, Ecological Society of America
- Frank S.D., Wratten S.D., Sandhu H.S., Shrewsbury P.M. (2007). Video analysis to determine how habitat strata affects predator diversity and predation of *Epiphyas postvittana* (Lepidoptera: Tortricidae) in a vineyard. *Biological Control* 41(2): 230-236.
- Frank T., Kehrl P., Germann C. (2007). Density and nutritional condition of carabid beetles in wildflower areas of different age. *Agriculture, Ecosystems & Environment* 120(2-4): 377-383.
- Frank T., Nentwig W. (1995). Ground dwelling spiders (Araneae) in sown weed strips and adjacent fields. *Acta Oecologica* 16(2): 179-193.
- Frankel E.N. (2007). Antioxidants in food and biology, facts and fiction, The Oily Press, Bridgwater, England, 254 p.
- Frantzen J., Rossi F. (2002). Integration of biological control of common groundsel (*Senecio vulgaris*) and chemical control. *Weed Science* 50: 787-793.
- Free (1993). Insect pollination of crops, 2nd ed., Academic Press, London UK, 684.
- Free J.B. (1993). Insect pollination of crops, 2 ed., Academic Press, London UK, 768 p.
- Fridley J.D. (2001). The influence of species diversity on ecosystem productivity: how, where and why? *Oikos* 93: 514-526.
- Fridley J.D. (2002). Resource availability dominates and alters the relationship between species diversity and ecosystem productivity in experimental plant communities. *Oecologia* 132: 271-277.
- Fridley J.D. (2003). Diversity effects on production in different light and fertility environments: an experiment with communities of annual plants. *Journal of Ecology* 91: 396-406.
- Fridley J.D., Brown R.L., Bruno J.F. (2004). Null models of ecological invasion and scale-dependent patterns of native and exotic species richness. *Ecology* 85: 3215-3222.
- Fridley J.D., Grime J.P., Bilton M. (2007). Genetic identity of interspecific neighbours mediates plant responses to competition and environmental variation in a species-rich grassland. *Journal of Ecology* 95: 908-915.
- Fried G., Chauvel B., Reboud X. (in press). A functional analysis of large scale temporal shifts in the sunflower weed assemblages in France between 1970 and 2000. *Journal of Vegetation Science*.
- Frouws J. (1998). The contested redefinition of the countryside. An analysis of rural discourses in the Netherlands. *Sociologia Ruralis* 38(1): 54-68.
- Fujiyoshi P.T., Gliessman S.R., Langenheim J.H. (2007). Factors in the suppression of weeds by squash interplanted in corn. *Weed Biology and Management* 7(2): 105-114.
- Funderburk J.E., Rhoads F.M., Teare I.D. (1994). Modifying Soil Nutrient Level Affects Soybean Insect Predators. *Agronomy Journal* 86(3): 581-585.
- Furlong M.J., Shi Z., Liu S., Zalucki M.P. (2004). Evaluation of the impact of natural enemies on *Plutella xylostella* L. (Lepidoptera: Yponomeutidae) populations on commercial Brassica farms. *Agricultural and Forest Entomology* 6(4): 311-322.
- Fye R.E. (1983). Cover Crop Manipulation for Building Pear Psylla (Homoptera, Psyllidae) Predator Populations in Pear Orchards. *Journal of Economic Entomology* 76(2): 306-310.
- Gadoum S., Terzo M., Rasmont P. (2007). Jachères apicoles et jachères fleuries : la biodiversité au menu de quelles abeilles ? *Le Courrier de l'environnement de l'INRA* 54: 57-63.
- Galloni M., Podda L., Vivarelli D., Quaranta M., Cristofolini G. (2008). Visitor diversity and pollinator specialization in Mediterranean legumes. *Flora* 203: 94-102.
- Gamfeldt L., Hillebrand H., Jonsson P.R. (2008). Multiple functions increase the importance of biodiversity for overall ecosystem functioning. *Ecology* 89: 1223-1231.
- Gange A.C., Brown V.K., Farmer L.M. (1990). A test of mycorrhizal benefit in an early successional plant community. *New Phytologist* 115(1): 85-91.
- Gange A.C., Llewellyn M. (1989). Factors affecting orchard colonisation by the black-kneed capsid (*Blepharidopterus angulatus* (Hemiptera: Miridae)) from alder windbreaks. *Annals of Applied Biology* 114(2): 221-230.
- Gange A.C., Smith A.K. (2005). Arbuscular mycorrhizal fungi influence visitation rates of pollinating insects *Ecological Entomology* 30(5): 600-606.
- Gao L.L., Horbury R., Nair R.M., Singh K.B., Edwards O.R. (2007). Characterization of resistance to multiple aphid species (Hemiptera: Aphididae) in *Medicago truncatula*. *Bulletin of entomological research* 97(1): 41-48.
- Gao M., Liu H., Qin G., Qiao Y., Wang Z. (2000). Influence of mulching in apple orchard on the azotobacter contents in soil. *Journal of fruit science* 17(3): 185-187.
- Garbeva P., van Elsas J.D., van Veen J.A. (2007). Rhizosphere microbial community and its response to plant species and soil history. *Plant and Soil* 302(1-2): 19-32.
- Garbeva P., van Veen J.A., van Elsas J.D. (2004). Assessment of the diversity, and antagonism towards *Rhizoctonia solani* AG3, of *Pseudomonas* species in soil from different agricultural regimes. *FEMS Microbiology Ecology* 47(1): 51-64.
- Gardiner M.M., Landis D.A. (2007). Impact of intraguild predation by adult *Harmonia axyridis* (Coleoptera: Coccinellidae) on *Aphis glycines* (Hemiptera: Aphididae) biological control in cage studies. *Biological Control* 40(3): 386-395.
- Gardner R., Milne B.T., Turner M.G., O'Neill R.V. (1987). Neutral models for the analysis of broad-scale landscape pattern. *Landscape Ecology* 1(1): 19-28.
- Gardner R.H., Romme W.H., Turner M.G. (1999). Predicting forest fire effects at landscape scales. In *Spatial modelling of forest landscapes: Approaches and applications*, (Mladenoff D.J., Baker W.L., eds.), Cambridge University Press, Cambridge: 163-185.

- Garnier E., Cortez J., Billès G., Navas M.-L., Roumet C., Debussche M., Laurent G., Blanchard A., Aubry D., Bellmann A., Neill C., Toussaint J.-P. (2004). Plant functional markers capture ecosystem properties during secondary succession. *Ecology* 85(9): 2630-2637.
- Garrett K.A., Nelson R.J., Mundt C.C., Chacón G., Jaramillo R.E., Forbes G.A. (2001). The effects of host diversity and other management components on epidemics of potato late blight in the humid highland tropics. *Phytopathology* 91(10): 993-1000.
- Gartner T.B., Cardon Z.G. (2004). Decomposition dynamics in mixed-species leaf litter. *Oikos* 104: 230-246.
- Garvin M.H., Martinez F., Jesus J.B., Gutierrez M., Ruiz P., Cosin D.J.D. (2001). Effect of *Hormogaster elisae* (Oligochaeta: Hormogastridae) on the stability of soil aggregates. *European Journal of Soil Biology* 37(4): 273-276.
- Gastine A., Scherer-Lorenzen M., Leadley P.W. (2003). No consistent effects of plant diversity on root biomass, soil biota and soil abiotic conditions in temperate grassland communities. *Applied Soil Ecology* 24(1): 101-111.
- Gavaland A., Burnel L. (2005). Croissance et biomasse aérienne de noyers noirs en parcelle agroforestière à Les Eduts (Charente-Maritime). *Revue des Chambres d'Agriculture, Dossier "agroforesterie: produire autrement"* 945: 20-22.
- Geiger F., Bianchi F., Wackers F.L. (2005). Winter ecology of the cabbage aphid *Brevicoryne brassicae* (L.) (Homo., Aphididae) and its parasitoid *Diaeretiella rapae* (McIntosh) (Hym., Braconidae : Aphidiidae). *Journal of Applied Entomology* 129(9-10): 563-566.
- Gelsomino A., Badalucco L., Ambrosoli R., Crecchio C., Puglisi E., Meli S.M. (2006). Changes in chemical and biological soil properties as induced by anthropogenic disturbance: A case study of an agricultural soil under recurrent flooding by wastewaters. *Soil Biology & Biochemistry* 38(8): 2069-2080.
- Gem-Oniflor (2005). Audit de la filière miel - réactualisation des données économiques issues de l'audit 1997 121.
- Genersch E., Yue C., Fries I., de Miranda J.R. (2006). Detection of deformed wing virus, a honey bee viral pathogen, in bumble bees (*Bombus terrestris* and *Bombus pascuorum*) with wing deformities. *J. Invert. Pathol.* 91: 61-63.
- Geoffrion R. (2000). Les plantes toxiques dans les cultures : A redouter dans des parcelles non ou mal désherbées. *Phytoma, la défense des végétaux* 532: 8-11.
- Gerowitt B., Isselstein J., Marggraf R. (2003). Rewards for ecological goods—requirements and perspectives for agricultural land use. *Agriculture, Ecosystems & Environment* 98(1-3): 541-547.
- Ghazoul J. (2006). Floral diversity and the facilitation of pollination. *Journal of Ecology* 94(2): 295-304.
- Ghiloufi M., Saulas P., Picard D. (2001). Towards cropping systems with permanent ground cover and zero tillage in open field cropping systems in France, 1. *World Congress on Conservation Agriculture*, Madrid, 2001/10/01-05.
- Ghiorbani R., Seel, W., Litterick, A., Leifert, C. (2000). Evaluation of *Alternaria alternata* for biological control of *Amaranthus retroflexus*. *Weed Science* 48: 474-480.
- Ghosheh H.Z. (2005). Constraints in implementing biological weed control: a review. *Weed Biology and Management* 5: 83-92.
- Gianoli E., Ramos I., Alfaro-Tapia A., Valdez Y., Echegaray E.R., Yabar E. (2006). Benefits of a maize-bean-weeds mixed cropping system in Urubamba Valley, Peruvian Andes. *International Journal of Pest Management* 52(4): 283-289.
- Gibot-Leclerc S., Brault, M., Pinochet, X., Sallé, G. (2003). Rôle potentiel des plantes adventices du colza d'hiver dans l'extension de l'orobanche rameuse en Poitou-Charentes. *Compte Rendus de l'Académie des Sciences, Biologies* 326: 645-658.
- Gibson R.H., Nelson I.L., Hopkins G.W., Hamlett B.J., Memmott J. (2006). Pollinator webs, plant communities and the conservation of rare plants: arable weeds as a case study. *Journal of Applied Ecology* 43(2): 246-257.
- Giller K.E., Beare M.H., Lavelle P., Izac A.M.N., Swift M.J. (1997). Agricultural intensification, soil biodiversity and agroecosystem function. *Applied Soil Ecology* 6(1): 3-16.
- Giller K.E., Day J.M. (1985). Nitrogen fixation in the rhizosphere: significance in natural and agricultural systems. In *Ecological interactions in soil*, (Fitter A.H. et al., eds.), Blackwell, Oxford, UK: 127-147.
- Gillespie D.R., Quiring D.J.M. (1992). Competition between *Orius tristicolor* (White) (Hemiptera: Anthracoridae) and *Amblyseius cucumeris* (Oudemans) (Acari: Phytoseiidae) feeding on *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae). *Canadian Entomologist* 124(6): 1123-1128.
- Ginane C., Baumont R., Lassalas J., Petit M. (2002). Feeding behaviour and intake of heifers fed on hays of various quality, offered alone or in a choice situation. *Animal Research* 51: 177-188.
- Girard N., Bellon S., Hubert B., Lardon S., Moulin C.H., Osty P.L. (2001). Categorising combinations of farmers' land use practices: an approach based on examples of sheep farms in the south of France. *Agronomie* 21(5): 435-459.
- Girma H., Rao M.R., Sithanatham S. (2000). Insect pests and beneficial arthropod populations under different hedgerow intercropping systems in semiarid Kenya. *Agroforestry Systems* 50(3): 279-292.
- Gladine C., Rock E., Morand C., Bauchart D., Durand D. (2007). Bioavailability and antioxidant capacity of plant extracts rich in polyphenols, given as a single acute dose, in sheep made highly susceptible to lipoperoxidation. *British Journal of Nutrition* 98(4): 691-701.
- Gladis T. (1997). Bees versus flies? - Rearing methods and effectiveness of pollinators in crop germplasm regeneration, *Acta Horticulturae*, 437: 235-238.
- Glasser F., Schmidely P., Sauviant D., Doreau M. (2008). Digestion of fatty acids in ruminants: a meta-analysis of flows and variation factors: 2. C18 fatty acids. *Animal* 2: 691-704.
- Gobat J.M., Aragno M., Matthey W. (1998). Le sol vivant : base de pédologie, biologie des sols, 2ème ed., Collection Gérer l'Environnement, Presses Polytechniques et Universitaires Romandes, Lausanne, 519 p.
- Goh K.M., Ridgen G.E., Daly M.J. (1995). Understorey biomass production and biological nitrogen fixation in an organic apple orchard in Canterbury. *Communications in soil science and plant analysis* 26(19-20): 3261-3273
- Gomez-Limon J., Fernandez J.V.D.L. (1999). Changes in use and landscape preferences on the agricultural-livestock landscapes of the central Iberian Peninsula (Madrid, Spain). *Landscape and Urban Planning* 44(4): 165-175.
- Gordon P.L., McEwen F.L. (1984). Insecticide-stimulated reproduction of *Myzus persicae*, the green peach aphid (Homoptera: Aphididae). *Canadian Entomologist* 116(5): 783-784.
- Gormsen D., Olsson P.A., Hedlund K. (2004). The influence of collembolans and earthworms on AM fungal mycelium. *Applied Soil Ecology* 27(3): 211-220.
- Goulson D. (2003). *Bumblebees : their behaviour and ecology*, Oxford University Press, 235 p.

- Goulson D., Hughes W.O.H., Derwent L.C., Stout J.C. (2002). Colony growth of the bumblebee, *Bombus terrestris*, in improved and conventional agricultural and suburban habitats. *Oecologia* 130(2): 267-273.
- Gowling G.R., Emden H.F.v. (1994). Falling aphids enhance impact of biological control by parasitoids on partially aphid-resistant plant varieties. *Annals of Applied Biology* 125(2): 233-242.
- Graham J.H. (2000). Assessing cost of arbuscular mycorrhizal symbiosis in agroecosystems. In Current advances in mycorrhizal research, (Podila G.K., Douds Jr D.D., eds.), 105, APS Press, St Paul NM: 127-140.
- Grant W.B. (1998). Milk and other dietary influences on coronary heart disease. *Alternative medicine review* 3(4): 281-294.
- Grasswitz T.R., Burts E.C. (1995). Effect of native natural enemies on the population dynamics of the grape mealybug, *Pseudococcus maritimus* (Hom.: Pseudococcidae), in apple and pear orchards. *Entomophaga* 40(1): 105-117.
- Graulet B., Chauveau-Duriot B., Noziere P., Doreau M. (2006). Teneur et composition en caroténoïdes des contenus digestifs et du plasma chez la vache au pâturage, 13. *Rencontres Recherches Ruminants*: 193.
- Grayston S.J., Campbell C.D., Bardgett R.D., Mawdsley J.L., Clegg C.D., Ritz K., Griffiths B.S., Rodwell J.S., Edwards S.J., Davies W.J., Elston D.J., Millard P. (2004). Assessing shifts in microbial community structure across a range of grasslands of differing management intensity using CLPP, PLFA and community DNA techniques. *Applied Soil Ecology* 25(1): 63-84.
- Green J.L., Holmes A.J., Westoby M., Oliver I., Briscoe D., Dangerfield M., Gillings M., Beattie A.J. (2004). Spatial scaling of microbial eukaryote diversity. *Nature* 432(7018): 747-750.
- Greenleaf S., Kremen C. (2006). Wild bee species increase tomato production and respond differently to surrounding land use in Northern California. *Biological Conservation* 133(1): 81-87.
- Greenleaf S.S., Kremen C. (2006). Wild bees enhance honey bees' pollination of hybrid sunflower. *Proceedings of the National Academy of Sciences of the United States of America* 103: 13890-13895.
- Grekul C., W, Bork, E,W, (2004). Herbage yield losses in perennial pasture due to canada thistle (*Cirsium arvense*). *Weed Technology* 18: 784-794.
- Griffiths B.S., Bonkowski M., Roy J., Ritz K. (2001). Functional stability, substrate utilisation and biological indicators of soils following environmental impacts. *Applied Soil Ecology* 16(1): 49-61.
- Griffiths B.S., Ritz K., Bardgett R.D., Cook R., Christensen S., Ekelund F., Sorensen S.J., Baath E., Bloem J., de Ruiter P.C., J. D., Nicolardot B. (2000). Ecosystem response of pasture soil communities to fumigation-induced microbial diversity reductions: an examination of the biodiversity-ecosystem function relationship. *Oikos* 90(2): 279-294.
- Griffiths B.S., Ritz K., Wheatley R., Kuan H.L., Boag B., Christensen S., Ekelund F., Sorensen S.J., Muller S., Bloem J. (2001). An examination of the biodiversity-ecosystem function relationship in arable soil microbial communities. *Soil Biology & Biochemistry* 33(12-13): 1713-1722.
- Griffiths G.J.K., Winder L., Holland J.M., Thomas C.F.G., Williams E. (2007). The representation and functional composition of carabid and staphylinid beetles in different field boundary types at a farm-scale. *Biological Conservation* 135(1): 145-152.
- Grigulis K., Lavorel S., Davies I.D., Dossantos A., Lloret F., Vilà M. (2005). Landscape-scale positive feedbacks between fire and expansion of the large tussock grass, *Ampelodesmos mauritanica*, in Catalan shrublands. *Global Change Biology* 11: 1042-1053.
- Griinari J.M., Corl B.A., Lacy S.H., Chouinard P.Y., Nurmela K.V.V., Bauman D.E. (2000). Conjugated linoleic acid is synthesized endogenously in lactating dairy cows by D9- desaturase. *Journal of nutrition* 130: 2285-2291.
- Grime J.P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology* 86: 902-906.
- Grime J.P. (2000). Plant strategies, vegetation processes and ecosystem properties, John Wiley & Sons Ltd, Chichester, UK, 456 p.
- Grime J.P. (2002). Declining plant diversity: empty niches or functional shifts? *Journal of Vegetation Science* 13: 457-460.
- Grime J.P., Mackey J.M.L., Hiller S.H., Read D.J. (1987). Floristic diversity in a model system using experimental microcosms. *Nature* 328: 420-422.
- Gross K.L., Mittelbach G.G., Reynolds H.L. (2005). Grassland invasibility and diversity: responses to nutrients, seed input, and disturbance. *Ecology* 86: 476-486.
- Gross N., Robson T.M., Lavorel S., Albert C., Le Bagousse-Pinguet Y., Guillemin R. (2008). Plant response traits mediate the effects of subalpine grasslands on soil moisture. *New Phytologist* 180(3): 652-662.
- Grosskopf G. (2005). Biology and life history of *Cheilosia urbana* (Meigen) and *Cheilosia psilophthalma* (Becker), two sympatric hoverflies approved for the biological control of hawkweeds (*Hieracium* spp.) in New Zealand. *Biological Control* 35(2): 142-154.
- Grundmann G.L., Debouzie D. (2000). Geostatistical analysis of the distribution of NH₄ and NO₂ oxidizing bacteria and serotypes at the millimetric scale along a soil transect. *FEMS Microbiology Ecology* 1170: 1-6.
- Gruner D.S. (2005). Biotic resistance to an invasive spider conferred by generalist insectivorous birds on Hawaiï Island. *Biological Invasions* 7(3): 541-546.
- Grüter D., Schmid B., Brandl H. (2006). Influence of plant diversity and elevated atmospheric carbon dioxide levels on belowground bacterial diversity. *BMC Microbiology* 6: 68, .
- Gu D., Wright D.J., Waage J.K. (1995). Influence of sublethal effects of insecticides on the population dynamics of *Plutella xylostella* and its parasitoid *Diadegma eucerocephala* in the field. *Entomologia Sinica* 2(1): 77-82.
- Guérin G., Pfmilin A., Léger F. (1994). Stratégie d'alimentation. Méthodologie d'analyse et de diagnostic de l'utilisation et de la gestion des surfaces fourragères et pastorales, (Lignes C., ed.), Institut d'Elevage, 36 p.
- Guichard H., Leconte D., Picoche B., Pages J., Simon J.C. (2006). Influence de la composition floristique des prairies permanentes normandes sur les caractéristiques des laits crus dérivés. *Fourrages* 188: 457-475.
- Gurr G.M., van Emden H.F., Wratten S.D. (1998). Habitat manipulation and natural enemy efficiency: implications for the control of pests. In Conservation Biological Control, (Barbosa P., ed.), Academic Press, San Diego London Boston New York Sydney Tokyo Toronto: 155-183.
- Gurr G.M., Wratten S.D., Altieri M.A. (2004). Ecological engineering for pest management - Advances in habitat manipulation for arthropods, CSIRO Publishing - CABI Publishing, Collingwood - Oxon, 232 p.
- Gurr G.M., Wratten S.D., Barbosa P. (2000). Success in Conservation Biological Control of arthropods. In Biological Control: Measures of success, (Gurr G.M., Wratten S.D., eds.), Kluwer Academic Pub.: 105-132.

- Gurr G.M., Wratten S.D., Luna J.M. (2003). Multi-function agricultural biodiversity: pest management and other benefits. *Basic and Applied Ecology* 4(2): 107-116.
- Güsewell S., Zuberbühler N., Clerc C. (2005). Distribution and functional traits of *Solidago gigantea* in a Swiss lakeshore wetland. *Botanica Helvetica* 115: 63-75.
- Gutierrez J.R., Meserve P.L., Herrera S., Contreras L.C., Jaksic F.M. (1997). Effects of small mammals and vertebrate predators on vegetation in the Chilean semi-arid zone. *Oecologia* 109(3): 398-406.
- Hacker S.D., Bertness M.D. (1995). A herbivore paradox: why salt marsh aphids live on poor-quality plants. *American Naturalist* 145(2): 192-210.
- Haddad N.M., Tilman D., Haarstad J., Ritchie M.E., Knops J.M.H. (2001). Contrasting effects of plant richness and composition on insect communities: a field experiment. *American Naturalist* 158(1): 17-35.
- Hagn A., Pritsch K., Schloter M., Munch J.C. (2003). Fungal diversity in agricultural soil under different farming management systems, with special reference to biocontrol strains of *Trichoderma* spp. *Biology and Fertility of Soils* 38(4): 236-244.
- Halaj J., Wise D.H. (2001). Terrestrial trophic cascades: How much do they trickle? *American Naturalist* 157(3): 262-281.
- Halley J.M., Thomas C.F.G., Jepson P.C. (1996). A model for the spatial dynamics of linyphiid spiders in farmland. *Journal of Applied Ecology* 33(3): 471-492.
- Halliwell B. (1990). How to characterize a biological antioxidant. *Free Radic Res Commun* 9(1): 1-32.
- Hance T. (1986). Experiments on the population control of *Aphis fabae* by different densities of Carabidae (Coleoptera: Carabidae). *Annales de la Societe Royale Zoologique de Belgique* 116(1): 15-24.
- Häni F. (1990). Farming systems research at Ipsach, Switzerland - The "Third Way" Project. *Recherche agronomique en Suisse* 29(4): 257-271.
- Häni F.J., Boller E.F., Keller S. (1998). Natural regulation at the farm level. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Pickett C.H., Bugg R.L., eds.), University of California Press, Berkeley - Los Angeles - London: 161-210.
- Hanley M.E. (2004). Seedling herbivory and the influence of plant species richness in seedling neighbourhoods. *Plant Ecology* 170: 35-41.
- Hanley M.E., Goulson D. (2003). Introduced weeds pollinated by introduced bees: cause or effect? *Weed Biology and Management* 3(4): 204-212.
- Hanna R., Zalom F.G., Roltsch W.J. (2003). Relative impact of spider predation and cover crop on population dynamics of *Erythroneura variabilis* in a raisin grape vineyard. *Entomologia Experimentalis et Applicata* 107(3): 177-191.
- Hardeman F., Sjolting S. (2007). Metagenomic approach for the isolation of a novel low-temperature-active lipase from uncultured bacteria of marine sediment. *FEMS Microbiology Ecology* 59(2): 524-534.
- Hargrove W.L. (1986). Winter legumes as a nitrogen source for No-Till grain sorghum. *Agronomy Journal* 78(1): 70-74.
- Harmon J.P., Hladilek E.E., Hinton J.L., Stodola T.J., Andow D.A. (2003). Herbivore response to vegetational diversity: spatial interaction of resources and natural enemies. *Population Ecology* 45(2): 75-81.
- Harrison S.K., Regnier E.E., Schmoll J.T. (2003). Postdispersal predation of giant ragweed (*Ambrosia trifida*) seed in no-tillage corn. *Weed Science* 51(6): 955-964.
- Hart M.M., Reader R.J. (2002). Taxonomic basis for variation in the colonization strategy in arbuscular mycorrhizal fungi. *New Phytologist* 153: 335-344.
- Hartmann A., Giraud J.J., Catroux G. (1998). Genotypic diversity of *Sinorhizobium* (formerly *Rhizobium*) meliloti strains isolated directly from a soil and from nodules of alfalfa (*Medicago sativa*) grown in the same soil. *FEMS Microbiology Ecology* 25(2): 107-116.
- Hartmann A., Rothballer M., Schmid M. (2008). Lorenz Hiltner, a pioneer in rhizosphere microbial ecology and soil bacteriology research. *Plant and Soil* 312(1-2): 7-14.
- Hartnett D.C., Wilson G.W.T. (1999). Mycorrhizae influence plant community structure and diversity in tallgrass prairie. *Ecology* 80(4): 1187-1195.
- Hartwig N.L., Ammon H.U. (2002). Cover crops and living mulches. *Weed Science* 50(6): 688-699.
- Harwood J.D., Obyrck J.J. (2005). Web-construction behavior of linyphiid spiders (Araneae, Linyphiidae): Competition and co-existence within a generalist predator guild. *Journal of Insect Behavior* 18(5): 593-607.
- Hasken K.H., Poehling H.M. (1995). Effects of different intensities of fertilisers and pesticides on aphids and aphid predators in winter wheat. *Agriculture, Ecosystems & Environment* 52(1): 45-50.
- Hassall M., Hawthorne A., Maudsley M., White P., Cardwell C. (1992). Effects of headland management on invertebrate communities in cereal fields. *Agriculture, Ecosystems & Environment* 40(1-4): 155-178.
- Hatfield J.C., Karlen D.C. (1994). *Sustainable Agriculture Systems*, CRC Press, 316 p.
- Hattenschwiler S., Tiunov A.V., Scheu S. (2005). Biodiversity and litter decomposition in terrestrial ecosystems. *Annual Review of Ecology and Systematics* 36: 191-218.
- Hauggaard-Nielsen H., Jømsgaard B., Kinane J., Jensen E.S. (2008). Grain legume-cereal intercropping: The practical application of diversity, competition and facilitation in arable and organic cropping systems. *Renewable Agriculture and Food Systems* 23: 3-12.
- Hausammann A. (1996). The effects of weed strip-management on pests and beneficial arthropods in winter wheat fields. *Journal of Plant Diseases and Protection* 103(1): 70-81.
- Hawes C., Houghton A.J., Osborne J.L., Roy D.B., Clark S.J., Perry J.N., Rothery P., Bohan D.A., Brooks D.R., Champion G.T., Dewar A.M., Heard M.S., Woiwod I.P., Daniels R.E., Young M.W., Parish A.M., Scott R.J., Firbank L.G., Squire G.R. (2003). Responses of plants and invertebrate trophic groups to contrasting herbicide regimes in the farm scale evaluations of genetically modified herbicide-tolerant crops. *Philosophical Transactions of the Royal Society B* 358: 1899-1913.
- Hawkes C., DeAngelis K., Firestone M.K. (2006). Root interactions with microbial communities and processes. In *The Rhizosphere: an ecological perspective*, (Whitbeck J.L., Cardon Z.G., eds.), Academic Press, Inc, San Diego CA: 1-31.
- He J.-S., Bazzaz F.A., Schmid B. (2002). Interactive effects of diversity, nutrients and elevated CO₂ on experimental plant communities. *Oikos* 97: 337-348.
- He J.-S., Wolfe-Bellin K.S., Schmid B., Bazzaz F.A. (2005). Density may alter diversity-productivity relationships in experimental plant communities. *Basic and Applied Ecology* 6: 505-517.

- He X.H., Critchley C., Bledsoe C. (2003). Nitrogen transfer within and between plants through common mycorrhizal networks (CMNs). *Critical Reviews in Plant Sciences* 22: 531-567.
- Heckendorn F., Häring D.A., Maurer V., Zinsstag J., Langhans W., Hertzberg H. (2006). Effect of sainfoin (*Onobrychis viciifolia*) silage and hay on established populations of *Haemonchus contortus* and *Cooperia curticei* in lambs. *Veterinary Parasitology* 142(3-4): 293-300.
- Hector A., Bagchi R. (2007). Biodiversity and ecosystem multifunctionality. *Nature* 448: 188-191.
- Hector A., Bazeley-White E., Loreau M., Otway S., Schmid B. (2002). Overyielding in grassland communities: testing the sampling effect hypothesis with replicated biodiversity experiment. *Ecology Letters* 5: 502-511.
- Hector A., Dobson K., Minns A., Bazeley-White E., Lawton J. (2001). Community diversity and invasion resistance: an experimental test in a grassland ecosystem and a review of comparable studies. *Ecological Research* 16(5): 819-831.
- Heemsbergen D.A., Berg M.P., Loreau M., van Haj J.R., Faber J.H., Verhoef H.A. (2004). Biodiversity effects on soil processes explained by interspecific functional dissimilarity. *Science* 306(5698): 1019-1020.
- Hein L., van Koppen K., de Groot R.S., van Ierland E.C. (2005). Spatial scales, stakeholders and the valuation of ecosystem services. *Ecological Economics* 57(2): 209-228.
- Helenius J. (1989). The influence of mixed intercropping of oats with field beans on the abundance and spatial distribution of cereal aphids (Homoptera, Aphididae). *Agriculture, Ecosystems & Environment* 25(1): 53-73.
- Helland S.J., Holland J.B. (2001). Blend response and stability and cultivar blending ability in oat. *Crop Science* 41(6): 1689-1696.
- Helland S.J., Holland J.B. (2003). Genome-wide genetic diversity among components does not cause cultivar blend responses. *Crop Science* 43(5): 1618-1627.
- Hernandez J.B., Cisneros C.V., Moreno C.M., Claramunt J.L.F., Casco J.M.G. (1998). Manejo en extensivo del ganado porcino autoctono en España. *Anaporc* 183: 74-84.
- Herrera C.M. (1987). Components of pollinator "quality": comparative analysis of a diverse insect assemblage. *Oikos* 50: 79-90.
- Herrera C.M. (1989). Pollinator abundance, morphology, and flower visitation rate : analysis of the "quantity" component in a plant-pollinator system. *Oecologia* 80: 241-248.
- Herrmann F., Westphal C., Moritz R.F.A., Steffan-Dewenter I. (2007). Genetic diversity and mass resources promote colony size and forager densities of a social bee (*Bombus pascuorum*) in agricultural landscapes. *Molecular Ecology* 16(6): 1167-1178.
- Herzog F., Dreier S., Hofer G., Marfurt C., Schubach B., Spiess M., Walter T. (2005). Effect of ecological compensation areas on floristic and breeding bird diversity in Swiss agricultural landscapes. *Agriculture, Ecosystems & Environment* 108(3): 189-204.
- Hickerson C.A.M., Anthony C.D., Walton B.M. (2005). Edge effects and intraguild predation in native and introduced centipedes: evidence from the field and from laboratory microcosms. *Oecologia* 146(1): 110-119.
- Hickman J.M., Wratten S.D. (1994). Use of *Phacelia tanacetifolia* (Hydrophyllaceae) as a pollen resource to enhance hoverfly (Diptera: Syrphidae) populations in sweetcorn fields. *Bulletin OILB/SROP* 17(4): 156-167.
- Hickman J.M., Wratten S.D. (1996). Use of *Phacelia tanacetifolia* strips to enhance biological control of aphids by hover fly larvae in cereal fields. *Journal of Economic Entomology* 89(4): 832-840.
- Hiddink G.A., van Bruggen A.H.C., Termorshuizen A.J., Raaijmakers J.M., Semenov A.V. (2005). Effect of organic management of soils on suppressiveness to *Gaeumannomyces graminis* var. *tritici* and its antagonist, *Pseudomonas fluorescens*. *European Journal of Plant Pathology* 113(4): 417-435.
- Higes M., Pilar G.P., Raquel M.H., Aránzazu M. (2007). Experimental infection of *Apis mellifera* honeybees with *Nosema ceranae* (Microsporidia). *J. Invert. Pathol.* 94: 211-217.
- Hilbeck A., Eckel C., Kennedy G.G. (1997). Predation on Colorado potato beetle eggs by generalist predators in research and commercial potato plantings. *Biological Control* 8(3): 191-196.
- Hill M.P., Cilliers C.J. (1999). *Azolla filiculoides* Lamarck (Pteridophyta : Azollaceae), its status in South Africa and control. *Hydrobiologia* 415: 203-206.
- Hillebrand H., Bennett D.M., Cadotte M.W. (2008). Consequences of dominance: A review of evenness effects on local and regional ecosystem processes. *Ecology* 89(6): 1510-1520.
- Hindayana D., Meyhofer R., Scholz D., Poehling H.M. (2001). Intraguild predation among the hoverfly *Episyrphus balteatus* de Geer (Diptera : Syrphidae) and other aphidophagous predators. *Biological Control* 20(3): 236-246.
- Hippa H., Koponen S., Osmonen O. (1978). Role of bees (Hym., Apidae) in pollination of the cloudberry (*Rubus chamaemorus* L.) in northern Fennoscandia. *Rep. Kevo Subarctic Res. Stat.* 14: 31-37.
- Hiremath A.J., Ewel J.J. (2001). Ecosystem Nutrient Use Efficiency, Productivity, and Nutrient Accrual in Model Tropical Communities. *Ecosystems* 4(7): 669-682.
- Hironori Y., Katsuhiko S. (1997). Cannibalism and interspecific predation in two predatory ladybirds in relation to prey abundance in the field. *Entomophaga* 42(1-2): 153-163.
- Hodgson J.G., Montserrat-Martí G., Tallwin J., Thompson K., Diaz S., Cabido M., Grime J.P., Wilson P.J., Band S.R., Bogard A., Cabido R., Caceres D., Castro-Díez P., Ferrer C., Maestro-Martínez M., Perez-Rontome M.C., Charles M., Cornelissen J.H.C., Dabbert S., Perez-Harguindeguy N., Krimly T., Sijtsma F.J., Strijker D., Vendramini F., Guerrero-Campo J., Hynd A., Jones G., Romo-Díez A., de Torres Espuny L., Villar-Salvador P., Zak M.R. (2005). How much will it cost to save grassland diversity? *Biological Conservation* 122(2): 263-273.
- Hoggart H., Buller H., Black R. (1995). *Rural Europe: Identities and Change*, Arnold, London, UK, 319 p.
- Hogh-Jensen H. (2006). Commentary: The nitrogen transfer between plants: an important but difficult flux to quantify. *Plant and Soil* 282(1-2): 1-5.
- Hogh-Jensen H., Schjoerring J.K. (2000). Below-ground nitrogen transfer between different grassland species: direct quantification by ¹⁵N leaf feeding compared with indirect dilution of soil ¹⁵N. *Plant and Soil* 227: 171-183.
- Hogh-Jensen H., Schjoerring J.K. (2001). Rhizodeposition of nitrogen by red clover, white clover and ryegrass leys. *Soil Biology & Biochemistry* 33: 439-448.
- Hokkanen H., Husberg G.B., Soderblom M. (1988). Natural enemy conservation for the integrated control of the rape blossom beetle *Meligethes aeneus* F. *Annales Agriculturae Fenniae* 27(4): 281-194.

- Holland J.M., Hutchison M.A.S., Smith B., Aebischer N.J. (2006). A review of invertebrates and seed-bearing plants as food for farmland birds in Europe. *Annals of Applied Biology* 148: 49-71.
- Holland J.M., Thomas C.F.G., Birkett T., Southway S., Oaten H. (2005). Farm-scale spatiotemporal dynamics of predatory beetles in arable crops. *Journal of Applied Ecology* 42(6): 1140-1152.
- Holland J.M., Thomas S.R. (1997). Assessing the role of beneficial invertebrates in conventional and integrated farming systems during an outbreak of *Sitobion avenae*. *Biological Agriculture & Horticulture* 15(1-4): 73-82.
- Holloway B.A. (1976). Pollen-feeding in hover-flies (Diptera: Syrphidae). *New Zealand Journal of Zoology* 3: 339-350.
- Holmes T.H., Rice K.J. (1996). Patterns of growth and soil-water utilization in some exotic annuals and native perennial bunchgrasses of California. *Annals of Botany* 78: 233-243.
- Holt R.D., Lawton J.H., Polis G.A., Martinez N.D. (1999). Trophic rank and the species-area relationship. *Ecology* 80: 1495-1504.
- Holzschuh A., Steffan-Dewenter I., Kleijn D., Tschamtké T. (2007). Diversity of flower-visiting bees in cereal fields: effects of farming system, landscape composition and regional context. *Journal of Applied Ecology* 44(1): 41-49.
- Hoogerkamp M., Rogaar H., Eijssackers H.J.P. (1983). Effect of earthworms on grassland on recently reclaimed polder soil in the Netherlands. In *Earthworm ecology. From Darwin to vermiculture*, (Satchell J.E., ed.), Chapman & Hall, London 85-105.
- Hooks C.R.R., Johnson M.W. (2003). Impact of agricultural diversification on the insect community of cruciferous crops. *Crop Protection* 22(2): 223-238.
- Hooper D. (1998). The role of complementarity and competition in ecosystem responses to variation in plant diversity. *Ecology* 79: 704-719.
- Hooper D.U., Chapin F.S., Ewel J.J., Hector A., Inchausti P., Lavorel S., Lawton J.H., Lodge D.M., Loreau M., Naeem S., Schmid B., Setälä H., Symstad A.J., Vandermeer J., Wardle D.A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs* 75(1): 3-35.
- Hooper D.U., Dukes J.S. (2004). Overyielding among plant functional groups in a long-term experiment. *Ecology Letters* 7: 95-105.
- Hooper D.U., Solan M., Symstad A.J., Diaz S., Gessner M.O., Buchmann N., Degrange V., Grime J.P., Hulot F., Mermillod-Blondin F., Roy J., Spehn E.M., Van Peer L. (2002). Species diversity, functional diversity, and ecosystem functioning. In *Biodiversity and ecosystem functioning: synthesis and perspectives*, (Loreau M. et al., eds.), Oxford University Press, Oxford, UK: 195-208.
- Hooper D.U., Vitousek P.M. (1997). The effects of plant composition and diversity on ecosystem processes. *Science* 277: 1302-1305.
- Hooper D.U., Vitousek P.M. (1998). Effects of plant composition and diversity on nutrient cycling. *Ecological Monographs* 68: 121-149.
- Horner-Devine M.C., Leibold M.A., Smith V.H., Bohannan B.J.M. (2003). Bacterial diversity patterns along a gradient of primary productivity. *Ecology Letters* 6(7): 613-622.
- Horton D.R., Broers D.A., Lewis R.R., Granatstein D., Zack R.S., Unruh T.R., Moldenke A.R., Brown J.J. (2003). Effects of mowing frequency on densities of natural enemies in three Pacific Northwest pear orchards. *Entomologia Experimentalis Et Applicata* 106(2): 135-145.
- Horton D.R., Lewis T.M., Broers D.A. (2004). Ecological and geographic range expansion of the introduced predator: *Anthocoris nemoralis* (Heteroptera: Anthocoridae) in North America: potential for nontarget effects? *American Entomologist* 50(1): 18-30.
- Hoste H., Gaillard L., Le Frileux Y. (2005). Consequences of the regular distribution of sainfoin hay on gastrointestinal parasitism with nematodes and milk production in dairy goats. *Small Ruminant Research* 59(2-3): 265-271.
- Hoste H., Jackson F., Athanasiadou S., Thamsborg S.M., Hoskin S.O. (2006). The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends in Parasitology* 22(6): 253-261.
- Howarth F.G. (2000). Non-target effects of biological control agents. In *Biological Control: Measures of success*, (Gurr G.M., Wratten S.D., eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands: 369-403.
- Huang H.T., Yang P. (1987). The ancient cultured Citrus ant, a tropical ant is used to control insect pests in southern China. *Bioscience* 37: 665-671.
- Hubbell S.P. (2001). A unified neutral theory of biodiversity and biogeography, *Monographs in population biology*, 32, Princeton University Press, 396 p.
- Huhta V., Sulkava P., Viberg K. (1998). Interactions between enchytraeid (*Cognettia sphagnetorum*), microarthropod and nematode populations in forest soil at different moistures. *Applied Soil Ecology* 9(1-3): 53-58.
- Hulme P.E., Bremner E.T. (2006). Assessing the impact of *Impatiens glandulifera* on riparian habitats: partitioning diversity components following species removal. *Journal of Applied Ecology* 43: 43-50.
- Hunt H.W., Coleman D.C., Ingham E.R., Ingham R.E., Elliott E.T., Moore J.C., Rose S.L., Reid C.P.P., Morley C.R. (1987). The detrital food web in a shortgrass prairie. *Biology and Fertility of Soils* 3(1): 57-68.
- Hunt H.W., Wall D.H. (2002). Modelling the effects of loss of soil biodiversity on ecosystem function. *Global Change Biology* 8: 33-50.
- Hunter Jr M.L. (1996). *Fundamentals of conservation biology*, Blackwell Publishing, Cambridge, 547 p.
- Hunter M.D. (2001). Insect population dynamics meets ecosystem ecology: effects of herbivory on soil nutrient dynamics. *Agriculture and Forest Entomology* 3(2): 77-84.
- Hunt-Joshi T.R., Root R.B., Blossey B. (2005). Disruption of weed biological control by an opportunistic mirid predator. *Ecological Applications* 15(3): 861-870.
- Hunziker M. (1995). The spontaneous reforestation in abandoned agricultural lands: perception and aesthetic assessment by locals and tourists. *Landscape and Urban Planning* 31(1-3): 399-410.
- Hussain I., Olsen K.R., Siemens J.C. (1998). Long-term tillage effects on physical properties of eroded soil. *Soil Science* 163(12): 970-981.
- Huston M.A. (2004). Management strategies for plant invasions: manipulating productivity, disturbance, and competition. *Diversity and Distributions* 10: 167-178.
- Huston M.A., Aarssen L.W., Austin M.P., Cade B.S., Fridley J.D., Garnier E., Grime J.P., Hodgson J., Lauenroth W.K., Thompson K., Vandermeer J.H., Wardle D.A. (2000). No consistent effect of plant diversity on productivity. *Science* 289: 1255-1257.
- Iason G.R., Lennon J.J., Pakeman R.J., Thoss V., Beaton J.K., Sim D.A., Elston D.A. (2005). Does chemical composition of individual Scots pine trees determine the biodiversity of their associated ground vegetation? *Ecology Letters* 8(4): 364-369.
- Inderjit, Dashini M.M. (1995). Allelopathic potential of an annual weed, *Polypogon monspeliensis*, in crops in India. *Plant and Soil* 173(2): 251-257.

- Innes L., Hobbs P.J., Bardgett R.D. (2004). The impacts of individual plant species on rhizosphere microbial communities varies in soils of different fertility. *Biology and Fertility of Soils* 40: 7-13.
- Irvin N.A., Scarratt S.L., Wratten S.D., Frampton C.M., Chapman R.B., Tilyanakis J.M. (2006). The effects of floral understoreys on parasitism of leafrollers (Lepidoptera : Tortricidae) on apples in New Zealand. *Agricultural and Forest Entomology* 8(1): 25-34.
- Ives A.R., Klug J.L., Gross K. (2000). Stability and species richness in complex communities. *Ecology Letters* 3: 399-411.
- Jackson L.E., Pascual U., Hodgkin T. (2007). Utilizing and conserving agrobiodiversity in agricultural landscapes. *Agriculture, Ecosystems & Environment* 121(3): 196-210.
- Jackson R.B., Banner J.L., Jobbagy E.G., Pockman W.T., Wall D.H. (2002). Ecosystem carbon loss with woody plant invasion of grasslands. *Nature* 418: 623-626.
- Jacobson D.A., Kring T.J. (1995). Efficacy of predators attacking *Helicoverpa zea* (Boddie) (Lepidoptera: Noctuidae) eggs on grain sorghum in the field. *Journal of Entomological Science* 30(2): 251-257.
- Jactel H., Brockerhoff E., Duelli P. (2005). A test of the biodiversity-stability theory: meta-analysis of tree species diversity effects on insect pest infestations, and re-examination of responsible factors. In *Forest diversity and function: temperate and boreal systems*, (Scherer-Lorenzen M. K.C., Schulze E.D., ed.), 176, Springer, Berlin Heidelberg New York: 235-262.
- James D.G. (2003). Synthetic herbivore-induced plant volatiles as field attractants for beneficial insects. *Environmental Entomology* 32(5): 977-982.
- Jankauskas B., Jankauskiene G. (2003). Erosion-preventive crop rotations for landscape ecological stability in upland regions of Lithuania. *Agriculture, Ecosystems & Environment* 95(1): 129-142.
- Jansen J.P. (1999). Selectivity of plant protection products used in wheat in summer on the major cereal aphid natural enemies. *Mededelingen - Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, Universiteit Gent* 64(3a): 25-40.
- Janson J., P., de Fontanges A., Deneufbourg F., Jouy L. (2004). Nuisibilité des adventices en production de semences fourragères : cas du vulpin dans le ray-grass anglais, *XIIème Colloque International sur la Biologie des Mauvaises Herbes*, Dijon (France): 81-88.
- Jarlan A., de Oliveira D., Gingras J. (1997). Effects of *Eristalis tenax* (Diptera: Syrphidae) pollination on characteristics of greenhouse sweet pepper fruits. *Journal of Economic Entomology* 90(6): 1650-1654.
- Jarlan A., de Oliveira D., Gingras J. (1997). Pollination by *Eristalis tenax* (Diptera: Syrphidae) and seed set of greenhouse sweet pepper. *Horticultural Entomology* 90: 1646-1649.
- Jauzein P. (1995). Flore des champs cultivés, INRA, SOPRA, 898.
- Jauzein P. (2001). Biodiversité des champs cultivés : l'enrichissement floristique. In *Dossiers de l'Environnement de l'INRA - Agriculture et biodiversité des plantes*, (Le Perchec S. et al., eds.), 21: 43-64.
- Jauzein P. (2001). L'appauvrissement floristiques des champs cultivés. In *Dossiers de l'Environnement de l'INRA - Agriculture et biodiversité des plantes*, (Le Perchec S. et al., eds.), 21: 65-78.
- Jeanros B., Scehovic J., Troxler J., Bachmann H.J., Bosset J.O. (1999). Comparaison des caractéristiques botaniques et chimiques d'herbages pâturés en plaine et en montagne. *Fourrages* 159: 277-292.
- Jégou D., Schrader S., Diestel H., Cluzeau D. (2001). Morphological, physical and biochemical characteristics of burrow walls formed by earthworms. *Applied Soil Ecology* 17(2): 165-174.
- Jenkinson D.S. (1977). The Nitrogen Economy of the Broadbalk Experiments. 1. Nitrogen Balance in the Experiments. *Rothamsted Experimental Report for 1976 Part 2*: 103-109.
- Jensen E.S., Ambus P., Bellostas N., Boisen S., Brisson N., Corre-Hellou G., Crozat Y., Dahmann C., Dibet A., von Fragstein P., Gooding M., Hauggaard-Nielsen H., Kasyanova E., Launay M., Monti M., Pristeri A. (2006). Intercropping of cereals and grain legumes for increased production, weed control, improved product quality and prevention of N-losses in European organic farming systems (INTERCROP), *Paper presented at Joint Organic Congress, Odense, Denmark, May 30-31, 2006. Org Print 7659*.
- Jiang L., Pu Z., Nemergut D.R. (2008). On the importance of the negative selection effect for the relationship between biodiversity and ecosystem functioning. *Oikos* 117(4): 488-493.
- Joffre R., Rambal S. (2006). Tree-grass interactions in the south-western Iberian Peninsula dehesas and montados. *Science et Changements Planétaires / Sécheresse* 17: 340-342.
- Joffre R., Rambal S., Ratte J.P. (1999). The dehesa system of southern Spain and Portugal as a natural ecosystem mimic. *Agroforestry Systems* 45: 57-79.
- Johansen A., Jensen E.S. (1996). Transfer of N and P from intact or decomposing roots of pea to barley interconnected by an arbuscular mycorrhizal fungus. *Soil Biology & Biochemistry* 28(1): 73-81.
- Johnsen K., Jacobsen C., Torsvik V., Sørensen J. (2001). Pesticide effects on bacterial diversity in agricultural soils - a review. *Biology and Fertility of Soils* 33(6): 443-453.
- Johnson M.T.G., Agrawal A.A. (2005). Plant genotype and environment interact to shape a diverse arthropod community on evening primrose (*Oenothera biennis*). *Ecology* 86(4): 874-885.
- Johnson M.T.G., Agrawal A.A. (2007). Covariation and composition of arthropod species across plant genotypes of evening primrose, *Oenothera biennis*. *Oikos* 116: 941-956.
- Johnson M.T.G., Lajeunesse M.J., Agrawal A.A. (2006). Additive and interactive effects of plant genotypic diversity on arthropod communities and plant fitness. *Ecology Letters* 9: 24-34.
- Johnston J.M. (2000). The contribution of microarthropods to aboveground food webs: A review and model of belowground transfer in a coniferous forest. *American Midland Naturalist* 143(1): 226-238.
- Jones G.A., Sieving K.E., Avery M.L., Meagher R.L. (2005). Parasitized and non-parasitized prey selectivity by an insectivorous bird. *Crop Protection* 24(2): 185-189.
- Jones G.A., Sieving K.E., Jacobson S.K. (2005). Avian diversity and functional insectivory on north-central Florida farmlands. *Conservation Biology* 19(4): 1234-1245.
- Jones J. (2006). A case study based assessment of the financial impact of agri-environment scheme participation: pre- and post-decoupling. *Journal of Farm Management* 12(8): 477-498.
- Jongmans A.G., Pulleman M.M., Balabane M., van Oort F., Marinissen J.C.Y. (2003). Soil structure and characteristics of organic matter in two orchards differing in earthworm activity. *Applied Soil Ecology* 24(3): 219-232.

- Joshi J., Otway S.J., Koricheva J., Pfisterer A.B., Alpei J., Roy B.A., Scherer-Lorenzen M., Schmid B., Spehn E.M., Hector A. (2004). Bottom-up effects and feedbacks in simple and diverse experimental grassland communities. *In* *Insects and Ecosystem Function* (Weisser W.W., Seimann E., eds.), Springer Verlag, Berlin: 115-134.
- Jouglar J.Y., Pflimlin A., Enjambert F. (1983). Les indigestions spumeuses chez les bovins consommant du trèfle blanc : facteurs de risques et contrôle. *Fourrages* 95: 189-202.
- Jullien J. (1999). Le zabre des céréales : il fait surface dans les pays de la Loire et en Bretagne. *Phytoma, La Défense des Végétaux* 520: 31-32.
- Jumpponen A., Högborg P., Huss-Danell K., Mulder C.P.H. (2002). Interspecific and spatial differences in nitrogen uptake in monocultures and two-species mixtures in north European grasslands. *Functional Ecology* 16: 454-461.
- Jurt L. (2000). Ne pas oublier les agriculteurs! *Hotspot* 2(Novembre 2002): 5.
- Jutila H.M. (2003). Germination in Baltic coastal wetland meadows: similarities and differences between vegetation and seed bank. *Plant Ecology* 166(2): 275-293.
- Jyoti J.L., Shelton A.M., Earle E.D. (2001). Identifying sources and mechanisms of resistance in crucifers for control of cabbage maggot (Diptera: Anthomyiidae). *Journal of Economic Entomology* 94: 942-949.
- Kahmen A., Perner J., Buchmann N. (2005). Diversity-dependent productivity in semi-natural grasslands following climate perturbations. *Functional Ecology* 19: 594-601.
- Kahmen A., Renker C., Unsicker S.B., Buchmann N. (2006). Niche complementarity for nitrogen: an explanation for the biodiversity and ecosystem functioning relationship? *Ecology* 87: 1244-1255.
- Kajita Y., Takano F., Yasuda H., Agarwala B.K. (2000). Effects of indigenous ladybird species (Coleoptera: Coccinellidae) on the survival of an exotic species in relation to prey abundance. *Applied Entomology and Zoology* 35(4): 473-479.
- Kardol P., Bezemer T.M., van der Wal A., Van der Putten W.H. (2005). Successional trajectories of soil nematode and plant communities in a chronosequence of ex-arable lands. *Biological Conservation* 126: 317-327.
- Karlen D.L., Sharpley A.N. (1994). Management strategies for sustainable soil fertility. *In* *Sustainable Agriculture Systems*, (Hatfield J.L., Karlen D.L., eds.), Lewis Publisher CRC Press, Boca Raton: 47-108.
- Karlen D.L., Varvel, G.E., Bullock D.G., Cruse R.M. (1994). Crop rotations for the 21st century. *Advances in Agronomy* 53: 1-45.
- Karlen D.L., Wollenhaupt N.C., Erbach D.C., Berry E.C., Swan J.B., Eash N.S., Jordahl J.L. (1994). Long-term tillage effects on soil quality. *Soil & Tillage Research* 32: 313-327.
- Kass D., Sylvester-Bradley R., Nygren P. (1997). The role of nitrogen fixation and nutrient supply in some agroforestry systems of the Americas. *Soil Biology & Biochemistry* 29(5-6): 775-785.
- Kay B.D., Angers D.A., Groenevelt P.H., Baldock J.A. (1988). Quantifying the influence of cropping history on soil structure. *Canadian Journal of Soil Science* 68: 359-368.
- Kazakou E., Garnier E., Gimenez O. (2007). Contribution of leaf life span and nutrient resorption to mean residence time: elasticity analysis. *Ecology* 88(7): 1857-1863.
- Keesing F., Holt R.D., Ostfeld R.S. (2006). Effects of species diversity on disease risk. *Ecology Letters* 9(4): 485-498.
- Keller M., Kollman, J., Edwards, P, J, (2000). Genetic introgression from distant provenances reduces fitness in local weed populations. *Journal of Applied Ecology* 37: 647-659.
- Keller S., Häni F. (2000). Ansprüche von Nützlingen und Schädlingen an den Lebensraum. *In* *Streifenförmige ökologische Ausgleichsflächen in der Kulturlandschaft: Ackerkrautstreifen, Buntbrache, Feldränder*, (Nentwig W., ed.), Verlag Agrarökologie., Bern, Hannover: 199-217.
- Kells A.R., Goulson D. (2003). Preferred nesting sites of bumblebee queens (Hymenoptera : Apidae) in agroecosystems in the UK. *Biological Conservation* 109(2): 165-174.
- Kendall D.A., Chinn N.E., Smith B.D., Tidboald C., Winstone L., Western N.M. (1991). Effects of straw disposal and tillage on spread of barley yellow dwarf virus in winter barley. *Annals of Applied Biology* 119(2): 359-364.
- Kennedy A.D., Biggs H., Zambatis N. (2003). Relationship between grass species richness and ecosystem stability in Kruger National Park, South Africa. *African Journal of Ecology* 41(2): 131-140.
- Kennedy T.A., Naeem S., Howe K.M., Knops J.M.H., Tilman D., Reich P. (2002). Biodiversity as a barrier to ecological invasion. *Nature* 417: 636-638.
- Kfir R. (2002). Increase in cereal stem borer populations through partial elimination of natural enemies. *Entomologia Experimentalis et Applicata* 104(2/3): 299-306.
- Khanh T.D., Chung M.I., Xuan T.D., Tawata S. (2005). The exploitation of crop allelopathy in sustainable agricultural production. *Journal of Agronomy & Crop Science* 191: 172-184.
- Kidd P., Prieto-Fernandez A., Monterroso C., Acea M. (2007). Rhizosphere microbial community and hexachlorocyclohexane degradative potential in contrasting plant species. *Plant and Soil* 302(1-2): 233-247.
- Kilian S., Werner D. (1996). Enhanced denitrification in plots of N₂-fixing faba beans compared to plots of a non-fixing legume and non-legumes. *Biology and Fertility of Soils* 21(1): 77-83.
- Kimura M. (1983). *The neutral theory of molecular evolution*, Cambridge University Press, Cambridge, 384 p.
- King A.P., Berry A.M. (2005). Vineyard $\delta^{15}\text{N}$, nitrogen and water status in perennial clover and bunch grass cover crop systems of California's central valley. *Agriculture, Ecosystems & Environment* 109(3-4): 262-272.
- King R.F., Dromph K.M., Bardgett R.D. (2002). Changes in species evenness of litter have no effect on decomposition processes. *Soil Biology & Biochemistry* 34(12): 1959-1963.
- Kingsley P.C., Scriber J.M., Harvey R.G. (1986). Relationship of weed density with leafhopper populations in alfalfa, Wisconsin. *Agriculture, Ecosystems & Environment* 17: 281-286.
- Kinkel L.L., Wilson M., Lindow S.E. (2000). Plant species and plant incubation conditions influence variability in epiphytic bacterial population size. *Microbial Ecology* 39(1): 1-11.
- Kiran, Sharma S.G., Singh S.P. (2006). Effect of monocrotophos and butachlor on N-fixing cyanobacteria and associated biochemical activities. *Annals of Plant Protection Sciences* 14(1): 210-214.

- Kirwan L., Luscher A., Sebastia M.T., Finn J.A., Collins R.P., Porqueddu C., Helgadottir A., Baadshaug O.H., Brophy C., Coran C., Dalmannsdottir S., Delgado I., Elgersma A., Fothergill M., Frankow-Lindberg B.E., Golinski P., Grieu P., Gustavsson A.M., Hoglind M., Huguenin-Elie O., Iliadis C., Jorgensen M., Kadziulene Z., Karyotis T., Lunnan T., Malengier M., Maltoni S., Meyer V., Nyfeler D., Nykanen-Kurki P., Parente J., Smit H.J., Thumm U., Connolly J. (2007). Evenness drives consistent diversity effects in intensive grassland systems across 28 European sites. *Journal of Ecology* 95(3): 530-539.
- Klein A.M., Steffan-Dewenter I., Damayanti B., Tscharntke T. (2002). Effects of land-use intensity in tropical agroforestry systems on coffee flower-visiting and trap-nesting bees and wasps. *Conservation Biology* 16: 1003-1014.
- Klein A.M., Steffan-Dewenter I., Tscharntke T. (2003). Fruit set of highland coffee increases with the diversity of pollinating bees. *Proceedings of the Royal Society B-Biological Sciences* 270: 955-961.
- Klein A.M., Vaissière B., Cane J.H., Steffan-Dewenter I., Cunningham S.A., Kremen C., Tscharntke T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society of London Series B Biological Sciences* 274(1608): 303-313.
- Klein E.K., Lavigne C., Picault H., Renard M., Gouyon P.-H. (2006). Pollen dispersal of oilseed rape: estimation of the dispersal function and effects of field dimension. *Journal of Applied Ecology* 43(1): 141-151.
- Klinger K. (1987). Effects of margin strips along a winter wheat field on the predatory arthropod fauna and cereal aphid infestation. [German]. *Journal of Applied Entomology* 104(1): 47-58.
- Klironomos J.N., Hart M.M. (2001). Food-web dynamics - Animal nitrogen swap for plant carbon. *Nature* 410(6829): 651-652.
- Klironomos J.N., McCune J., Hart M., Neville J. (2000). The influence of arbuscular mycorrhizae on the relation between plant diversity and productivity. *Ecology Letters* 3: 137-141.
- Klironomos J.N., Moutoglou P. (1999). Colonization of nonmycorrhizal plants by mycorrhizal neighbours as influenced by the collembolan, *Folsomia candida*. *Biology and Fertility of Soils* 29(3): 277-281.
- Klose S., Acosta-Martinez V., Ajwa H.A. (2006). Microbial community composition and enzyme activities in a sandy loam soil after fumigation with methyl bromide or alternative biocides. *Soil Biology & Biochemistry* 38(6): 1243-1254.
- Knight T.M., Chase J.M., Hillebrand H., Holt R.D. (2006). Predation on mutualists can reduce the strength of trophic cascades. *Ecology Letters* 9(11): 1173-1178.
- Knight T.M., Steets J.A., Vamosi J.C., Mazer S.J., Burd M., Campbell D.R., Dudash M.R., Johnston M., Mitchell R.J., Ashman T.I. (2005). Pollen limitation of plant reproduction: Ecological and Evolutionary Causes and Consequences. *Annual Review of Ecology, Evolution and Systematics* 36: 467-497.
- Knops J.M.H., Wedin D., Tilman D. (2001). Biodiversity and decomposition in experimental grassland systems. *Oecologia* 126: 429-433.
- Koch R.L., Hutchison W.D., Venette R.C., Heimpel G.E. (2003). Susceptibility of immature monarch butterfly, *Danaus plexippus* (Lepidoptera: Nymphalidae: Danainae), to predation by *Harmonia axyridis* (Coleoptera: Coccinellidae). *Biological Control* 28(2): 265-270.
- Koide R. (1991). Nutrient supply, nutrient demand and plant-response to mycorrhizal infection. *New Phytologist* 117: 365-386.
- Koricheva J., Mulder C.P.H., Schmid B., Joshi J., Huss-Danell K. (2000). Numerical responses of different trophic groups of invertebrates to manipulations of plant diversity in grasslands. *Oecologia* 125(2): 271-282.
- Körner C. (2004). Mountain biodiversity, its causes and function. *Ambio* 13: 11-17.
- Körner C., Stöcklin J., Reuther-Thiébaud L., Pelaez-Riedl S. (2008). Small differences in arrival time influence composition and productivity of plant communities. *New Phytologist* 177: 698-705.
- Korthals G.W., Smilauer P., van Dijk C., van der Putten W.H. (2001). Linking above- and below-ground biodiversity: abundance and trophic complexity in soil as a response to experimental plant communities on abandoned arable land. *Functional Ecology* 15(4): 506-514.
- Koss A.M., Snyder W.E. (2005). Alternative prey disrupt biocontrol by a guild of generalist predators. *Biological Control* 32(2): 243-251.
- Kowalchuk G.A., Buma D.S., de Boer W., Klinkhamer P.G.L., van Veen J.A. (2002). Effects of above-ground plant species composition and diversity on the diversity of soil-borne microorganisms. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* 81(1-4): 509-520.
- Kozdroj J., Trevors J.T., van Elsland J.D. (2004). Influence of introduced potential biocontrol agents on maize seedling growth and bacterial community structure in the rhizosphere. *Soil Biology & Biochemistry* 36(11): 1775-1784.
- Kreiter S., Brian F. (1987). Les Phytoseiidae de la vigne en France, *Proc. Intern. Conf. on "Pests in Agriculture"*, Paris, 1-3 décembre 1987, Ann. ANPP, 6 (3), ANPP, ed.: 241-249.
- Kreiter S., Tixier M.S., Bourgeois T. (2003). Do generalist phytoseiid mites (Gamasida: Phytoseiidae) have interactions with their host plants? *Insect Science and its Application* 23(1): 35-50.
- Kreiter S., Tixier M.S., Croft B.A., Auger P., Barret D. (2002). Plants and leaf characteristics influencing the predaceous mite *Kampimodromus aberrans* (Acari: Phytoseiidae) in habitats surrounding vineyards. *Environmental Entomology* 31(4): 648-660.
- Kremen C., Williams N.M., Aizen M.A., Gemmill-Herren B., LeBuhn G., Minckley R., Packer L., Potts S.G., Roulston T.A., Steffan-Dewenter I., zquez D.P., Winfree R., Adams L., Crone E.E., Greenleaf S.S., Keitt T.H., Klein A.-M., Regetz J., Ricketts T.H. (2007). Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology Letters* 10(4): 299-314.
- Kremen C., Williams N.M., Thorp R.W. (2002). Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences of the United States of America* 99(26): 16812-16816.
- Kremer R.G., Running S.W. (1996). Simulating seasonal soil water balance in contrasting semi-arid vegetation communities. *Ecological Modelling* 84: 151-162.
- Kring T.J., Gilstrap F.E. (1986). Beneficial role of corn leaf aphid, *Rhopalosiphum maidis* (Fitch) (Homoptera: Aphididae), in maintaining *Hippodamia* spp. (Coleoptera: Coccinellidae) in grain sorghum. *Crop Protection* 5(2): 125-128.
- Kromp B. (1999). Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agriculture, Ecosystems & Environment* 74(1-3): 187-228.
- Kropff M.J. (1988). Modelling the effects of weeds on crop production. *Weed Research* 28(6): 465-471.
- Kropff M.J., van Laar H.H. (1993). *Modelling Crop-Weed Interactions*, CAB International, Wallingford (UK), 274.

- Kropff M.J., Vossen F.J.H., Spitters C.J.T., de Groot W. (1984). Competition between a maize crop and a natural population of *Echinochloa crusgalli* (L.). *Netherlands journal of agricultural science* 32: 324-327.
- Krsek M., Wellington E. (2001). Assessment of chitin decomposer diversity within an upland grassland. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* 79(3): 261-267.
- Kruess A., Tschamtké T. (1994). Habitat fragmentation, species loss and biological control. *Science* 264(5165): 1581-1584.
- Kruess A., Tschamtké T. (2000). Species richness and parasitism in a fragmented landscape: experiments and field studies with insects on *Vicia sepium*. *Oecologia* 122(1): 129-137.
- Krupinsky J., Tanaka D., Merrill S., Liebig M., Hanson J. (2006). Crop sequence effects of ten crops in the northern great plains. *Agricultural Systems* 88(2-3): 227-254.
- Kuan H.L., Fenwick C., Glover L.A., Griffiths B.S., Ritz K. (2006). Functional resilience of microbial communities from perturbed upland grassland soils to further persistent or transient stresses. *Soil Biology & Biochemistry* 38(8): 2300-2306.
- Kulmatiski A., Beard K.H., Stevens J.R., Cobbold S.M. (2008). Plant-soil feedbacks: a meta-analytical review. *Ecology Letters* 11(9): 980-992.
- Kuske S., Widmer F., Edwards P.J., Turlings T.C.J., Babendreier D., Bigler F. (2003). Dispersal and persistence of mass released *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) in non-target habitats. *Biological Control* 27(2): 181-193.
- Kyamanywa S., Baliddawa C.W., Omolo E. (1993). Influence of cowpea/maize mixture on generalists predators and their effect on the population density of the legume flower thrips, *Megalurothrips sjostedti* Trybom (Thysanoptera: Thripidae). *Insect Science and its Application* 14(4): 493-499.
- Laakso J., Setälä H. (1999). Population and ecosystem-level effects of predation on nematodes. *Oecologia* 120: 279-286.
- Laakso J., Setälä H. (1999). Sensitivity of primary production to changes in the architecture of belowground food webs. *Oikos* 87(1): 57-64.
- Lacas J.G., Voltz M., Gouy V., Carluier N., Gril J.J. (2005). Using grassed strips to limit pesticide transfer to surface water: a review. *Agronomy for Sustainable Development* 25(2): 253-266.
- Lafay B., Burdon J.J. (2006). Molecular diversity of rhizobia nodulating the invasive legume *Cytisus scoparius* in Australia. *Journal of Applied Microbiology* 100(6): 1228-1238.
- Lagerlof J., Stark J., Svensson B. (1992). Margins of Agricultural Fields as Habitats for Pollinating Insects. *Agriculture, Ecosystems & Environment* 40(1-4): 117-124.
- Lagerlof J., Wallin H. (1993). The Abundance of Arthropods Along 2 Field Margins with Different Types of Vegetation Composition - an Experimental-Study. *Agriculture, Ecosystems & Environment* 43(2): 141-154.
- Laguette G., Depret G., Bourion V., Duc G. (2007). *Rhizobium leguminosarum* bv. *viciae* genotypes interact with pea plants in developmental responses of nodules, roots and shoots. *New Phytologist* 176(3): 680-690.
- Lahr J., van Kats R., Crum S. (2007). Helminthocides in nature. *Vakblad Natuur Bos Landschap* 4(2): 22-23.
- Lançon J. (1978). Les restitutions du bétail au pâturage et leurs effets - I. *Fourrages* 75: 55-88.
- Landis D.A., Menalled F.D., Costamagna A.C., Wilkinson T.K. (2005). Manipulating plant resources to enhance beneficial arthropods in agricultural landscapes. *Weed Science* 53(6): 902-908.
- Landis D.A., Wratten S.D., Gurr G.M. (2000). Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology* 45: 175-201.
- Lang A. (2003). Intraguild interference and biocontrol effects of generalist predators in a winter wheat field. *Oecologia* 134(1): 144-153.
- Langellotto G.A., Denno R.F. (2004). Responses of invertebrate natural enemies to complex-structured habitats: a meta-analytical synthesis. *Oecologia* 139(1): 1-10.
- Langellotto G.A., Rosenheim J.A., Williams M.R. (2006). Assessing trophic interactions in a guild of primary parasitoids and facultative hyperparasitoids: stable isotope analysis. *Oecologia* 150(2): 291-299.
- Lankau R.A., Strauss S.H. (2007). Mutual feedbacks maintain both genetic and species diversity in a plant community. *Science* 317: 1561-1563.
- Larick D.K., Hedrick H.B., Bailey M.E., Williams J.E., Hancock D.L., Garner G.B., Morrow R.E. (1987). Flavor constituents of beef as influenced by forage- and grain-feeding. *Journal of Food Science* 52(2): 245-251.
- Larkin R.P., Griffin T.S. (2007). Control of soilborne potato diseases using *Brassica* green manures. *Crop Protection* 26(7): 1067-1077.
- Larrère R., Brun A., Kalaora B., Nougarede O., Poupardin D. (1981). Forestiers et paysans. Les reboisements en montagne depuis l'Empire. Pratiques et politiques de la nature 1870-1960. *Recherches* 45: 57-84.
- Laurent F., Fontaine A. (2006). Pertes d'azote par lessivage : cultures intermédiaires, une efficacité immédiate et durable. *Perspectives agricoles* 327: 26-30.
- Lavelle P., Bignell D., Lepage M., Wolters V., Roger P., Ineson P., Heal O.W., Dhillon S. (1997). Soil function in a changing world: the role of invertebrate ecosystem engineers. *European Journal of Soil Biology* 33(4): 159-193.
- Lavelle P., Bignell D.E., Austen M.C., Brown V.K., Behan-Pelletier V., Garey J.R., Giller P.S., Hawkins S.J., Brown G.G., St John M., Hunt H.W., Paul E.A. (2004). Connecting soil and sediment biodiversity: the role of scale and implications for management. In *Sustaining Biodiversity and Functioning in Soils and Sediments*, (Wall D.H., ed.), Scientific Committee on Problems of the Environment (SCOPE) Series, Island Press, New York: 193-224.
- Lavelle P., Blouin M., Boyer J., Cadet P., Laffray D., Pham-Thi A.T., Reversat G., Settle W., Zuiluy Y. (2004). Plant parasite control and soil fauna diversity. *Comptes Rendus Biologies* 327(7): 629-638.
- Lavelle P., Decaens T., Aubert M., Barot S., Blouin M., Bureau F., Margerie P., Mora P., Rossi J.P. (2006). Soil invertebrates and ecosystem services. *European Journal of Soil Biology* 42: S3-S15.
- Lavelle P., Spain A.V. (2001). Soil ecology, Kluwer Academic Publishers, Netherlands, 654 p.
- Lavorel S., Gardner R., O'Neill R. (1993). Analysis of patterns in hierarchically structured landscapes. *Oikos* 67: 521-528.
- Lavorel S., Garnier E. (2002). Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology* 16: 545-556.
- Lawton J.H. (1978). Host-plant influences on insect diversity: the effects of space and time. In *Diversity of Insect Faunas*, (N. M.L.A.W., ed.), Blackwell Scientific Publications Oxford, UK: 105-125.

- Lawton J.H., Brown V.K. (1993). Redundancy in ecosystems. *In* Biodiversity and ecosystem function, (Schulze E.-D., Mooney H.A., eds.), Springer: 255-270.
- Le Bayon R.C., Binet F. (1999). Rainfall effects on erosion of earthworm casts and phosphorus transfers by water runoff. *Biology and Fertility of Soils* 30(1-2): 7-13.
- Le Floch S. (1996). Impacts paysagers de la populiculture. *Le Courrier de l'environnement* 29: 39-46.
- Le Gall A., Faverdin P., Thomet P., Verite R. (2001). Le pâturage en Nouvelle-Zélande : des idées pour les régions arrosées d'Europe *Fourrages* 166(2): 137-163.
- Le Gall A., Guernion J.M. (2004). Association graminées - trèfle blanc. Le pâturage gagnant, Collection Synthèse, Institut d'Élevage, Paris, 64 p.
- Le Maitre D.C., Versfeld D.B., Chapman R.A. (2000). The impact of invading alien plants on surface water resources in South Africa: a preliminary assessment. *Water South Africa* 26: 397-408.
- Le Roux X., Poly F., Currey P., Commeaux C., Hai B., Nicol G.W., Prosser J.I., Schloter M., Attard E., Klumpp K. (2008). Effects of aboveground grazing on coupling among nitrifier activity, abundance and community structure. *ISME Journal* 2(2): 221-232.
- Leather S.R. (1992). Aspects of aphid overwintering (Homoptera: Aphidinea: Aphididae). *Entomologia generalis* 17(2): 101-113.
- Ledgard S.F., Steele K.W. (1992). Biological nitrogen fixation in mixed legume/grass pastures. *Plant and Soil* 141(1): 137-153.
- Lee K.E. (1987). Peregrine species of earthworms. *In* On earthworms, (Bonvicini Pagliai A.M., Omodeo P., eds.), Mucchi Editore, Modena: 315-328.
- Lee M.R.F., Connelly P.L., Tweed J.K.S., Dewhurst R.J., Merry R.J., Scollan N.D. (2006). Effects of high-sugar ryegrass silage and mixtures with red clover silage on ruminant digestion. 2. Lipids. *Journal of animal science* 84(11): 3061-3070.
- Lefèbvre S., Clément J.-C., Pinay G., Thenail C., Durand P., Marmonier P. (2007). 15N-Nitrate signature in low-order streams: effects of land cover and agricultural practices. *Ecological Applications* 17: 2333-2346.
- Léger F., Bellon S., Guérin G. (2000). Outils et méthodes pour analyser les ressources au pâturage. *In* Ruptures, nouveaux enjeux, nouvelles fonctions, nouvelle image de l'élevage sur parcours, (Bourbouze A., Qarro M., eds.), CIHEAM - IAMM, Montpellier: 205-215.
- Leininger S., Urich T., Schloter M., Schwark L., Qi J., Nicol G.W., Prosser J.I., Schuster S.C., Schleper C. (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442(7104): 806-809.
- Lemaire G., Wilkins R., Hodgson J. (2005). Challenges for grassland science: managing research priorities. *Agriculture, Ecosystems & Environment* 108: 99-108.
- Lemke R., Goddard T., Hahn D., Burton D., Ellert B., Farrell R., Monreal M., Noot D. (2003). An inter-laboratory comparison of nitrous oxide analysis in Western Canada. *Communications in Soil Science and Plant Analysis* 33(15/18): 2705-2713.
- Lenné J.M., Wood D. (1999). Optimizing biodiversity for productive agriculture. *In* Agrobiodiversity: Characterization, Utilization and Management, (Wood D., Lenné J.M., eds.), CABI Publishing, Oxon UK, New-York USA: 447-470.
- Lepart J., Marty P. (2004). L'objet ou son image ? Science des représentations ou science des paysages. Les enjeux du transfert vers les gestionnaires, *L'évaluation du paysage : une utopie nécessaire ? à la recherche d'indicateurs/marqueurs interdisciplinaires*, Montpellier, France, 15 au 16 janvier 2004, (Eds.) P.D.H.A.R., ed.
- Lepart J., Marty P., Rousset O. (2000). Les conceptions normatives du paysage. Le cas des Grands Causses. *Natures, Sciences, Sociétés* 8: 16-25.
- Leps J. (2004). Variability in population and community biomass in a grassland community affected by environmental productivity and diversity. *Oikos* 107: 64-71.
- Leps J., Brown V.K., Diaz Lenz T.A., Gormsen D., Hedlung K., Kailova J., Korthals G.W., Mortimer S.R., Rodriguze-Barrueco C., Roy J., Santa Regina I., van Dijk C., van der Putten W. (2001). Separating the chance effect from other diversity effects in the functioning of plant communities. *Oikos* 92: 123-134.
- Lepš J., Doležal J., Bezemer T.M., Brown V.K., Hedlung K., Igual A.M., Bracht Jörgensen H., Lawson C.S., Mortimer S.R., Peix Geldart A., Rodriguez Barrueco C., Santa Regina I., Šmilauer P., van der Putten W.H. (2007). Long-term effectiveness of sowing high and low diversity seed mixtures to enhance plant community development on ex-arable fields *Applied Vegetation Science* 10(1): 97-110.
- Lepš J., Osbornová-Kosinová J., Rejmánek M. (1982). Community stability, complexity and species life history strategies. *Plant Ecology* 50(1): 53-63.
- LeRoy C.J., Whitham T.G., Keim P., Marks J.C. (2006). Plant genes link forests and streams. *Ecology* 87(1): 255-261.
- Lesica P., Allendorf F.W. (1999). Ecological genetics and the restoration of plant communities: mix or match? *Restoration Ecology* 7(1): 42-50.
- Letourneau D.K. (1988). Soil management for pest control: a critical appraisal of the concepts, *Global Perspectives on Agroecology and sustainable agricultural systems. Proceedings of the Sixth International Scientific Conference of IFOAM*, Santa Cruz: 581-587.
- Letourneau D.K. (1998). Conservation biology: lessons for conserving natural enemies. *In* Conservation Biological Control, (Barbosa P., ed.), Academic Press, San Diego London Boston New York Sydney Tokyo Toronto: 9-38.
- Letourneau D.K., Dyer L.A. (1998). Experimental test in lowland tropical forest shows top-down effects through four trophic levels. *Ecology* 79: 1678-1687.
- Letourneau D.K., Dyer L.A., Vega G.C. (2004). Indirect effects of a top predator on a rain forest understory plant community. *Ecology* 85(8): 2144-2152.
- Levine J.M., D'Antonio C.M. (1999). Elton revisited: a review of evidence linking diversity and invasibility. *Oikos* 87: 15-26.
- Levine J.M., Vila M., d'Antonio C.M., Dukes J.S., Grigulis K., Lavorel S. (2003). Mechanisms underlying the impacts of exotic plant invasions. *Proceedings of the Royal Society B: Biological Sciences* 270(1517): 775-781.
- Liagre F. (1992). Les pratiques de cultures intercalaires dans les noyeraies fruitières du Dauphiné, Mémoire de fin d'étude de l'Engref.
- Liebman M., Dyck E. (1993). Crop Rotation and Intercropping Strategies for Weed Management. *Ecological Applications* 3(1): 92-122.
- Liiri M., Haimi J., Setälä H. (2002). Community composition of soil microarthropods of acid forest soils as affected by wood ash application. *Pedobiologia* 46(2): 108-124.
- Liiri M., Setälä H., Haimi J., Pennanen T., Fritze H. (2002). Relationship between soil microarthropod species diversity and plant growth does not change when the system is disturbed. *Oikos* 96(1): 137-149.

- Liiri M., Setälä H., Haimi J., Pennanen T., Fritze H. (2002). Soil processes are not influenced by the functional complexity of soil decomposer food webs under disturbance. *Soil Biology & Biochemistry* 34(7): 1009-1020.
- Lindsey A.H. (1984). Reproductive biology of Apiaceae. I: Floral visitors to Thaspium and Zizia and their importance in pollination. *American journal of botany* 71(3): 375-387.
- Lithourgidis A., Dhima K., Vasilakoglou I., Dordas C., Yiakoulaki M. (2007). Sustainable production of barley and wheat by intercropping common vetch. *Agronomy for Sustainable Development* 27(2): 95-99.
- Lloret F., Medail F., Brundu G., Hulme P.E. (2004). Local and regional abundance of exotic plant species on Mediterranean islands: are species traits important? *Global Ecology and Biogeography* 13(1): 37-45.
- Lo P.L. (2004). Toxicity of pesticides to *Halmus chalybeus* (Coleoptera: Coccinellidae) and the effect of three fungicides on their densities in a citrus orchard. *New Zealand Journal of Crop and Horticultural Science* 32(1): 69-76.
- Loiseau P., Louault F., Le Roux X., Bardy M. (2005). Does extensification of rich grasslands alter the C and N cycles, directly or via species composition? *Basic and Applied Ecology* 6(3): 275-287.
- Long Z.T., Mohler C.L., Carson W.P. (2003). Extending the resource concentration hypothesis to plant communities: Effects of litter and herbivores. *Ecology* 84(3): 652-665.
- Lopezaraiza-Mikel M.E., Hayes R.B., Whalley M.R., Memmott J. (2007). The impact of an alien plant on a native plant-pollinator network: an experimental approach. *Ecology Letters* 10(7): 539-550.
- Lopez-Bellido Garrido R.J., Lopez-Bellido L. (2001). Effects of crop rotation and nitrogen fertilization on soil nitrate and wheat yield under rainfed Mediterranean conditions. *Agronomie* 21(6-7): 509-516.
- Loreau M. (2000). Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* 91: 3-17.
- Loreau M. (2001). Microbial diversity, producer-decomposer interactions and ecosystem processes: a theoretical model. *Proceedings of the Royal Society of London Series B-Biological Sciences* 268(1464): 303-309.
- Loreau M., Hector A. (2001). Partitioning selection and complementarity in biodiversity experiments. *Nature* 412: 72-76.
- Loreau M., Mouquet N., Holt R.D. (2003). Meta-ecosystems: a theoretical framework for a spatial ecosystem ecology. *Ecology Letters* 6: 673-679.
- Loreau M., Naeem S., Inchausti P., Bengtsson J., Grime J.P., Hector A., Hooper D.U., Huston M.A., Raffaelli D.G., Schmid B., Tilman D., Wardle D.A. (2001). Biodiversity and ecosystem functioning: Current knowledge and future challenges. *Science* 294(5543): 804-808.
- Loreau M., Yachi S. (1999). Biodiversity and ecosystem productivity in a fluctuating environment : The insurance hypothesis. *Proceedings of the National Academy of Science of the United States of America* 96: 1463-1468.
- Losey J.E., Denno R.F. (1998). Positive predator-predator interactions: enhanced predation rates and synergistic suppression of aphid populations. *Ecology* 79(6): 2143-2152.
- Losey J.E., Denno R.F. (1999). Factors facilitating synergistic predation: The central role of synchrony. *Ecological Applications* 9(2): 378-386.
- Lotter D.W., Seidel R., Liebhardt W. (2003). The performance of organic and conventional cropping systems in an extreme climate year. *American Journal of Alternative Agriculture* 18(2): 146-154.
- Lourenço M., van Ranst G., de Smet S., Raes K., Fievez V. (2007). Effect of grazing pasture with different botanical composition by lambs on rumen fatty acid metabolism and fatty acid pattern of longissimus muscle and subcutaneous fat. *Animal* 1: 537-545.
- Lourenço M., Vlaeminck B., Bruinenberg M., Demeyer D., Fievez V. (2005). Milk fatty acid composition and associated rumen lipolysis and fatty acid hydrogenation when feeding forages from intensively managed or semi-natural grasslands. *Animal Research* 54: 471-484.
- Louvet J., Rouxel F., Alabouvette C. (1976). Recherches sur la résistance des sols aux maladies. I - Mise en évidence de la nature microbiologique de la résistance d'un sol au développement de la fusariose vasculaire du melon. *Annales de Phytopathologie* 8(4): 425-436.
- Lövei G.L., MacDougall D., Bramley G., Hodgson D.J., Wratten S.D. (1992). Floral resources for natural enemies: the effect of *Phacelia tanacetifolia* (Hydrophyllaceae) on within-field distribution of Hoverflies (Diptera: Syrphidae), 45. *New Zealand Plant Protection Conference*: 60-61.
- Lucas E., Alomar O. (2002). Impact of the presence of *Dicyphus tamaninii* Wagner (Heteroptera: Miridae) on whitefly (Homoptera: Aleyrodidae) predation by *Macrolophus caliginosus* (Wagner) (Heteroptera: Miridae). *Biological Control* 25(2): 123-128.
- Ludwig J.A., Tongway D.J., Marsden S.G. (1999). Stripes, strands or stipples: modelling the influence of three landscape banding patterns on resource capture and productivity in semi-arid woodlands, Australia. *Catena* 37: 257-273.
- Ludwig J.A., Wilcox B.P., Breashears D.D., al. e. (2005). Vegetation patches and run-off erosion as interacting hydroecological processes in semiarid landscapes. *Ecology* 86: 288-297.
- Luginbühl Y. (2001). La demande sociale de paysage, Conseil National du Paysage.
- Luka H., Uehlinger G., Pfiffner L., Muhlethaler R., Blick T. (2006). Extended field margins - a new element of ecological compensation in farmed landscapes - deliver positive impacts for Articulata. *Agrarforschung* 13(9): 386-391.
- Lumaret J.P., Galante E., Lumbreras C., Mena J., Bertrand M., Bernal J.L., Cooper J.F., Kadiri N., Crowe D. (1993). Field Effects of Ivermectin Residues on Dung Beetles. *Journal of Applied Ecology* 30(3): 428-436.
- Luxhoj J., Nielsen N.E., Jensen L.S. (2003). Influence of 15N4+ application on gross N turnover rates in soil. *Soil Biology & Biochemistry* 35: 603-606.
- Lyon J.-P. (1965). Influence de quelques facteurs sur l'expression du potentiel de multiplication des Syrphidés aphidiphages. *Ann. Epiphyties* 16(4): 397-398.
- Lyons K.G., Brigham C.A., Traut B.H., Schwartz M.D. (2005). Rare species and ecosystem functioning. *Conservation Biology* 19(4): 1019-1024.
- Ma K., Hao S., Zhao H., Kang L. (2007). Strip cropping wheat and alfalfa to improve the biological control of the wheat aphid *Macrosiphum avenae* by the mite *Allothrombium ovatum*. *Agriculture, Ecosystems & Environment* 119(1/2): 49-52.
- Ma W.C., Immerzeel J., Bodt J. (1995). Earthworm and food interactions on bioaccumulation and disappearance in soil of polycyclic aromatic hydrocarbons: Studies on phenanthrene and fluoranthene. *Ecotoxicology and Environmental Safety* 32(3): 226-232.

- Ma Y., Dickinson N.M., Wong M.H. (2002). Toxicity of Pb/Zn mine tailings to the earthworm *Pheretima* and the effects of burrowing on metal availability. *Biology and Fertility of Soils* 36(1): 79-86.
- Ma Y., Dickinson N.M., Wong M.H. (2006). Beneficial effects of earthworms and arbuscular mycorrhizal fungi on establishment of leguminous trees on Pb/Zn mine tailings. *Soil Biology & Biochemistry* 38(6): 1403-1412.
- Mabbett T. (1991). Straw incorporation trials reveal arable slug damage will increase. *Agriculture International* 43(11): 304-306.
- MacArthur R.H., Wilson E.O. (1967). The theory of island biogeography, Princeton University Press, Princeton, NH, 203 p.
- MacDonald D., Crabtree J.R., Wiesinger G., Dax T., Stamou N., Fleury P., Gutierrez Lazpita J., Gibon A. (2000). Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response. *Journal of Environmental Management*. 59(1): 47-69.
- MacLaren B.E., Peterson B.O. (1994). Wolves, moose, and tree rings on Isle Royale. *Science* 266: 1555-1558.
- MacLeod A., Wratten S.D., Sotherton N.W., Thomas M.B. (2004). 'Beetle banks' as refuges for beneficial arthropods in farmland: long-term changes in predator communities and habitat. *Agricultural and Forest Entomology* 6(2): 147-154.
- Macpherson H. (2004). Landscape's ocular-centrism – and beyond?, *Frontis Workshop From Landscape Research to Landscape Planning: Aspects of Integration, Education and Application*, Wageningen, NL, 1-6 June 2004, Tress B. T.G., Fry G. & Opdam P. (Eds.), ed.: 95-104.
- MacRae I.V., Croft B.A. (1997). Intra- and interspecific predation by adult female *Metaseiulus occidentalis* and *Typhlodromus pyri* (Acari: Phytoseiidae) when provisioned with varying densities and ratios of *Tetranychus urticae* (Acari: Tetranychidae) and phytoseiid larvae. *Experimental & Applied Acarology* 21(4): 235-245.
- Maddox G.D., Root R.B. (1990). Structure of the encounter between Goldenrod (*Solidago altissima*) and its diverse insect fauna. *Ecology* 71(6): 2115-2124.
- Madritch M.D., Hunter M.D. (2002). Phenotypic diversity influences ecosystem functioning in an oak sandhills community. *Ecology* 83: 2084-2090.
- Madritch M.D., Hunter M.D. (2003). Intraspecific litter diversity and nitrogen deposition affect nutrient dynamics and soil respiration. *Oecologia* 136: 124-128.
- Madritch M.D., Hunter M.D. (2004). Phenotypic diversity and litter chemistry affect nutrient dynamics during litter decomposition in a two species mix. *Oikos* 105: 125-131.
- Maestre F.T., Reynolds J.F. (2006). Spatial heterogeneity in soil nutrient supply modulates nutrient and biomass responses to multiple global change drivers in model grassland communities. *Global Change Biology* 12(12): 2431-2441.
- Maherali H., Klironomos J.N. (2007). Influence of Phylogeny on fungal community assembly and ecosystem functioning. *Science* 316(5832): 1746-1748.
- Maigre D., Aerny J. (2001). Enherbement et fumure azotée sur cv. Gamay dans le Valais central. 2. Résultats analytiques et organoleptiques. *Revue Suisse de Viticulture, Arboriculture et d'Horticulture* 33(4): 199-204.
- Maillet J. (1993). Nouvelles pratiques culturales et nouvelles mauvaises herbes, *Faut-il sauver les mauvaises herbes ?*, Gap (France), Gap-Charance C.B.d., ed., AFCEV, BRG, Conservatoire Botanique de Gap-Charance 33-40.
- Mair J., Port G.R. (2002). The influence of mucus production by the slug, *Deroceras reticulatum*, on predation by *Pterostichus madidus* and *Nebria brevicollis* (Coleoptera: Carabidae). *Biocontrol Science and Technology* 12(3): 325-335.
- Mando A., Stroosnijder L., Brussaard L. (1996). Effects of termites on infiltration into crusted soil. *Geoderma* 74(1-2): 107-113.
- Mantle P., G, Shaw, S. (1977). A case study of the aetiology of ergot disease of cereals and grasses. *Plant Pathology* 26: 121-126.
- Marino P.C., Gross K.L., Landis D.A. (1997). Weed seed loss due to predation in Michigan maize fields. *Agriculture, Ecosystems & Environment* 66(3): 189-196.
- Marino P.C., Landis D.A. (1996). Effect of landscape structure on parasitoid diversity in agroecosystems. *Ecological Applications* 6: 276-284.
- Marley C.L., Cook R., Keatinge R., Barrett J., Lampkin N.H. (2003). The effect of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) on parasite intensities and performance of lambs naturally infected with helminth parasites. *Veterinary Journal* 112: 147-155.
- Marrs R.H., Galtress K., Tong C., Cox E.S., Blackbird S.J., Pakeman R.J., Le Duc M.G. (2007). Competing conservation goals, biodiversity or ecosystem services: Element losses and species recruitment in a managed moorland-bracken model system. *Journal of Environmental Management* 85: 1034-1047.
- Marrs R.H., Watt A.S. (2006). Biological flora of the British Isles: *Pteridium aquilinum* (L.) Kuhn. *Journal of Ecology* 94: 1272-1321.
- Marschner H., Dell B. (1994). Nutrient uptake in mycorrhizal symbiosis. *Plant and Soil* 159: 89-102.
- Marshall E.J.P. (2004). Agricultural landscapes: field margin habitats and their interaction with crop production. *Journal of Crop Improvement* 12(1/2): 365-404.
- Marshall E.J.P., Brown V.K., Boatman N.D., Lutman P.J.W., Squire G.R., Ward L.K. (2003). The role of weeds in supporting biological diversity within crop fields. *Weed Research* 43: 77-89.
- Martin B., Verdier-Metz I., Buchin S., Hurtaud C., Coulon J.B. (2005). How do the nature of forages and pasture diversity influence the sensory quality of dairy livestock products? *Animal Science* 81(2): 205-212.
- Martin J.K., Puckridge D.W. (1982). Carbon flow through the rhizosphere of wheat crops in South Australia. In *The cycling of carbon, nitrogen, sulphur and phosphorus in terrestrial and aquatic ecosystems*, (Galbally E., Freney J.R., eds.), Australian Academy of Science, Canberra: 77-81.
- Martin-Laurent F., Barres B., Wagschal I., Piutti S., Devers M., Soulas G., Philippot L. (2006). Impact of the maize rhizosphere on the genetic structure, the diversity and the atrazine-degrading gene composition of cultivable atrazine-degrading communities. *Plant and Soil* 282(1-2): 99-115.
- Martin-Laurent F., Cornet L., Ranjard L., Lopez-Gutierrez J.C., Philippot L., Schwartz C., Chaussod R., Catroux G., Soulas G. (2004). Estimation of atrazine-degrading genetic potential and activity in three French agricultural soils. *FEMS Microbiology Ecology* 48(3): 425-435.
- Marulanda A., Azcon R., Ruiz-Lozano J.M. (2003). Contribution of six arbuscular mycorrhizal fungal isolates to water uptake by *Lactuca sativa* plants under drought stress. *Physiologia Plantarum* 119(4): 526-533.

- Mary B., Laurent F., Beaudoin N. (2002). La gestion durable de la fertilisation azotée, *65. IIRB Congress*, Bruxelles, 2002/02/13-14: 59-65.
- Marzio M.-C., Jolivet C. (1997). À Méron, l'Outarde volera-t-elle au secours des messicoles ? *Crex* 2: 63-73.
- Maskell L.C., Firbank L.G., Thompson K., Bullock J.M., Smart S.M. (2006). Interactions between non-native plant species and the floristic composition of common habitats. *Journal of Ecology* 94(6): 1052-1060.
- Mateille T., Cadet P., Fargette M. (2008). Control and management of plant parasitic nematode communities in a soil conservation approach. In *Integrated Management and Biocontrol of Vegetable and Grain Crops Nematodes*, (Ciancio A., Mukerji K.G., eds.), *Integrated Management of Plant Pests and Diseases*, 2, Springer: 79-97.
- Mathews C.R., Bottrell D.G., Brown M.W. (2002). A comparison of conventional and alternative understory management practices for apple production: multi-trophic effects. *Applied Soil Ecology* 21(3): 221-231.
- Mathews C.R., Bottrell D.G., Brown M.W. (2004). Habitat manipulation of the apple orchard floor to increase ground-dwelling predators and predation of *Cydia pomonella* (L.) (Lepidoptera : Tortricidae). *Biological Control* 30(2): 265-273.
- Matile L. (1993). Les Diptères d'Europe occidentale. Tome 1, (Boubée S.N.d.E., ed.), Paris, 439 p.
- Matsumoto T., Itoika T., Nishida T. (2003). Cascading effects of a specialist parasitoid on plant biomass in a Citrus agroecosystem. *Ecological Research* 18(6): 651-659.
- Mattingly W.B., Hewlate R., Reynolds H.L. (2007). Species evenness and invasion resistance of experimental grassland communities. *Oikos* 116: 1164-1170.
- Mattson W.J.J. (1980). Herbivory in relation to plant nitrogen content. *Annual Review of Ecology and Systematics* 11: 119-161.
- Mauchline A.L., Watson S.J., Brown V.K., FroudWilliams R.J. (2005). Post-dispersal seed predation of non-target weeds in arable crops. *Weed Research* 45(2): 157-164.
- Maudsley M., Seeley B., Lewis O. (2002). Spatial distribution patterns of predatory arthropods within an English hedgerow in early winter in relation to habitat variables. *Agriculture, Ecosystems & Environment* 89(1/2): 77-89.
- Maudsley M.J. (2000). A review of the ecology and conservation of hedgerow invertebrates in Britain. *Journal of Environmental Management* 60(1): 65-76.
- Mauz I. (2002). Gens, Cornes et Crocs. Relations hommes - animaux et conceptions du monde, en Vanoise, au moment de l'arrivée des loups. PhD thesis, Ecole Nationale Supérieure du Génie Rural, des Eaux et Forêts, Paris, France.
- Mazzola M. (2004). Assessment and management of soil microbial community structure for disease suppression. *Annual Review of Phytopathology* 42: 35-59.
- Mboukou-Kimbatsa I., Bernhard-Reversat F. (2001). Effect of exotic tree plantations on invertebrates soil macrofauna and population changes with Eucalyptus hybrids and plantation age. In *Effect of exotic tree plantations on plant diversity and biological soil fertility in the Congo savanna: a reference to eucalypts*, (Bernhard-Reversat F., ed.), CIFOR, Bogor: 49-55.
- Mc Dowall F.H., Mc Gillivray A. (1963). Studies on the properties of New Zealand butterfat - VII. Effect of the stage of maturity of ryegrass fed to cows on the characteristics of butterfat and its carotene and vitamin A contents. *Journal of Dairy Science* 30: 59-66.
- McClure M.S., Andreadis T.G., Lacy G.H. (1982). Manipulating orchard ground cover to reduce invasion by leafhopper vectors of peach X-disease. *Journal of Economic Entomology* 75: 64-68.
- McDonald G., K. (2003). Competitiveness against grass weeds in field pea genotypes. *Weed Research* 43(1): 48-58.
- McGillivray C.W. (1995). Testing predictions of the resistance and resilience of vegetation subjected to extreme events. *Functional Ecology* 9: 640-649.
- McGovern R.J., Polston J.E., Danyluk G., Hiebert E., Abouzid A.M., Stansly P.A. (1994). Identification of a natural weed host of tomato mottle geminivirus in Florida. *Plant Disease* 78: 1102-1106.
- McGregor S.E. (1976). Insect pollination of cultivated crop plants, *Agricultural Handbook*, 496, Agricultural Research Services, 411 p.
- McIntyre N.E., Thompson T.R. (2003). A comparison of conservation reserve program habitat plantings with respect to arthropod prey for grassland birds. *American Midland Naturalist* 150(2): 291-301.
- McKemey A.R., Symondson W.O.C., Glen D.M., Brain P. (2001). Effects of slug size on predation by *Pterostichus melanarius* (Coleoptera: Carabidae). *Biocontrol Science and Technology* 11(1): 81-91.
- McMurtry J.A., Morse J.G., Johnson H.G. (1992). Studies of the impact of *Euseius* species (Acari: Phytoseiidae) on citrus mites using predator exclusion and predator release experiments. *Experimental & Applied Acarology* 15(4): 233-248.
- McNaughton S.J., Banyikwa F.F., McNaughton M.M. (1997). Promotion of the Cycling of Diet-Enhancing Nutrients by African Grazers. *Science* 278(5344): 1798-1800.
- Meagher R.L., Meyer J.R. (1990). Influence of Ground Cover and Herbicide Treatments on Tetranychus-Urticae Populations in Peach Orchards. *Experimental & Applied Acarology* 9(3-4): 149-158.
- Meddich A., Oihabi A., Abbas Y., Bizid E. (2000). Effect of arbuscular mycorrhizal fungi on drought resistance of clover. *Agronomie* 20(3): 283-295.
- Meek B., Loxton D., Sparks T., Pywell R., Pickett H., Nowakowski M. (2002). The effect of arable field margin composition on invertebrate biodiversity. *Biological Conservation* 106(2): 259-271.
- Meitner M.J. (2004). Scenic beauty of river views in the Grand Canyon: relating perceptual judgments to locations. *Landscape and Urban Planning* 68: 3-13.
- Mela T. (2003). Red clover grown in a mixture with grasses / yield, persistence and dynamics of quality characteristics. *Agricultural and Food Science in Finland* 12(3-4): 195-212.
- Memmott J. (1999). The structure of a plant-pollinator food web. *Ecology Letters* 2: 276-280.
- Memmott J., Craze P.G., Waser N.M., Price M.V. (2007). Global warming and the disruption of plant-pollinator interactions. *Ecology Letters* 10: 710-717.
- Memmott J., Waser N.M. (2002). Integration of alien plants into a native flower-pollinator visitation web. *Proceedings of the Royal Society B: Biological Sciences* 269(1508): 2395-2399.
- Memmott J., Waser N.M., Price M.V. (2004). Tolerance of pollination networks to species extinctions. *Proceedings of the Royal Society B: Biological Sciences* 271: 2605-2611.
- Menalled F.D., Landis D.A., Dyer L.E. (2004). Research and extension supporting ecologically based IPM systems. *Journal of Crop Improvement* 11(1/2): 153-174.

- Menalled F.D., Lee J.C., Landis D.A. (1999). Manipulating carabid beetle abundance alters prey removal rates in corn fields. *Biocontrol* 43(4): 441-456.
- Menalled F.D., Marino P.C., Renner K.A., Landis D.A. (2000). Post-dispersal weed seed predation in Michigan crop fields as a function of agricultural landscape structure. *Agriculture, Ecosystems & Environment* 77(3): 193-202.
- Menalled F.D., Smith R.G., Dauer J.T., Fox T.B. (2007). Impact of agricultural management on carabid communities and weed seed predation. *Agriculture, Ecosystems & Environment* 118(1-4): 49-54.
- Mercier J., Lindow S.E. (2001). Field performance of antagonistic bacteria identified in a novel laboratory assay for biological control of fire blight of pear. *Biological Control* 22(1): 66-71.
- Mercier V., Combe F., Fauvel G., Marboutie G., Lespinasse Y. (2000). A four year experiment on the development of mite populations on scab resistant apples, *International conference on integrated fruit production*, Leuven, Belgium, 1998/07/26-1998/08/01, Acta Horticulturae, ISHS: 325-330.
- Merry R.J., Lee M.R.F., Davies D.R., Dewhurst R.J., Moorby J.M., Scollan N.D., Theodorou M.K. (2006). Effects of high-sugar ryegrass silage and mixtures with red clover silage on ruminant digestion. 1. *In vitro* and *in vivo* studies of nitrogen utilization. *Journal of animal science* 84: 3049-3060.
- Meuret M. (1993). Piloter l'ingestion au pâturage. In *Pratiques d'élevage extensif. Identifier, modéliser, évaluer*, (Landais E., ed.), Nathan, Paris, France: 161-198.
- Meuret M., Bruchou C. (1994). Modélisation de l'ingestion selon la diversité des choix alimentaires réalisés par la chèvre au pâturage sur parcours, 1. *Recontres Recherches Ruminants*: 225-228.
- Meyer J.R., Zehr E.I., Meagher R.L., Jr., Salvo S.K. (1992). Survival and growth of peach trees and pest populations in orchard plots managed with experimental ground covers. *Agriculture, Ecosystems & Environment* 41(3-4): 353-363.
- Meyhofer R., Hindayana D. (2000). Effects of intraguild predation on aphid parasitoid survival. *Entomologia Experimentalis et Applicata* 97(1): 115-122.
- Michaud J.P. (2002). Classical biological control: a critical review of recent programs against citrus pests in Florida. *Annals of the Entomological Society of America* 95(5): 531-540.
- Michaud J.P. (2002). Invasion of the Florida citrus ecosystem by *Harmonia axyridis* (Coleoptera: Coccinellidae) and asymmetric competition with a native species, *Cycloneda sanguinea*. *Environmental Entomology* 31(5): 827-835.
- Michaud J.P. (2004). Natural mortality of Asian citrus psyllid (Homoptera: Psyllidae) in central Florida. *Biological Control* 29(2): 260-269.
- Michener C.D. (2000). *The bees of the world*, Johns Hopkins University Press, Baltimore USA, 952 p.
- Michon G. (2007). *Domesticating forest*, Editions IRD.
- Mikola J., Setälä H. (1998). Productivity and trophic-level biomasses in a microbial-based soil food web. *Oikos* 82: 158-168.
- Mikola J., Setälä H. (1998). Relating species diversity to ecosystem functioning - Mechanistic backgrounds and experimental approach with a decomposer food web. *Oikos* 83: 180-194.
- Mikola J., Yeates G.W., Barker G.M., Wardle D.A., Bonner K.I. (2001). Effects of defoliation intensity on soil food-web properties in an experimental grassland community. *Oikos* 92(2): 333-343.
- Milbau A., Nijs I., De Raedemaeker F., Reheul D., De Cauwer B. (2005). Invasion in grassland gaps: the role of neighbourhood richness, light availability and species complementarity during two successive years. *Functional Ecology* 19: 27-37.
- Millan de la Pena N.M., Butet A., Delettre Y., Morant P., Burel F. (2003). Landscape context and carabid beetles (Coleoptera : Carabidae) communities of hedgerows in western France. *Agriculture, Ecosystems & Environment* 94(1): 59-72.
- Millennium Ecosystem Assessment (2005). *Ecosystems and Human Well-Being*. Synthesis, Island Press: 155 p. <http://www.millenniumassessment.org/documents/document.356.aspx.pdf>
- Miller A. (1961). The Mouth Parts and Digestive Tract of Adult Dung Beetles (Coleoptera: Scarabaeidae), with Reference to the Ingestion of Helminth. *Journal of Parasitology* 47(5): 735-744.
- Miller B.L., Perry H.B., M.A. F., de Gregorio R.M. (1993). The effect of vitamin E addition on calves fed milk replacer. *Journal of Dairy Science* 76(Suppl 1): 221.
- Miller R.M., Jastrow J.D. (2000). Mycorrhizal fungi influence soil structure. In *Arbuscular mycorrhizas: Molecular biology and Physiology*, (Kapulnik Y., Douds D.D., eds.), Kluwer Academic, Dordrecht, The Netherlands: 3-18.
- Min B.R., Hart S.P. (2003). Tannins for suppression of internal parasites. *Journal of animal science* 81(14): E102-109.
- Min B.R., McNabb W.C., Barry T.N., Peters J.S. (2000). Solubilization and degradation of ribulose-1,5-bisphosphate carboxylase/oxygenase (EC 4.1.1.39; Rubisco) protein from white clover (*Trifolium repens*) and *Lotus corniculatus* by rumen microorganisms and the effect of condensed tannins on these processes. *Journal of Agricultural Science* 34: 305-317.
- Minarro M., Hemptinne J.L., Dapena E. (2005). Colonization of apple orchards by predators of *Dysaphis plantaginea*: sequential arrival, response to prey abundance and consequences for biological control. *Biocontrol* 50(3): 403-414.
- Misgav A. (2000). Visual preferences of the public for vegetation groups in Israel. *Landscape and Urban Planning* 48: 143-159.
- Mitchell C.E., Tilman D., Groth J.V. (2002). Effects of grassland plant species diversity, abundance, and composition on foliar fungal disease. *Ecology* 83(6): 1713-1726.
- Mochizuki A., Mitsunaga T. (2004). Non-target impact assessment of the introduced green lacewing, *Chrysoperla carnea* (Stephens) (Neuroptera : Chrysopidae) on the indigenous sibling species, *C-nipponensis* (Okamoto) through interspecific predation. *Applied Entomology and Zoology* 39(2): 217-219.
- Mohamed A.H., Lester P.J., Holtzer T.O. (2000). Abundance and effects of predators and parasitoids on the Russian wheat aphid (Homoptera: Aphididae) under organic farming conditions in Colorado. *Environmental Entomology* 29(2): 360-368.
- Mols C.M.M., Visser M.E. (2002). Great tits can reduce caterpillar damage in apple orchards. *Journal of Applied Ecology* 39: 888-899.
- Monclus R., Dreyer E., Villar M., Delmotte F.M., Delay D., Petit J.-M., Barbaroux C., Le Thiec D., Bréchet C., Brignolas F. (2006). Impact of drought on productivity and water use efficiency in 29 genotypes of *Populus deltoides* x *Populus nigra*. *New Phytologist* 169: 765-777.
- Monetti L.N., Fernandez N.A. (1995). Seasonal population dynamics of the European red mite (*Panonychus ulmi*) and its predator *Neoseiulus californicus* in a sprayed apple orchard in Argentina (Acari: Tetranychidae, Phytoseiidae). *Acarologia* 36(4): 325-331.

- Montané F., Rovira P., Casals P. (2007). Shrub encroachment into mesic mountain grasslands in the Iberian peninsula: Effects of plant quality and temperature on soil C and N stocks. *Global Biogeochemical Cycles* 21.
- Monzón V.H., Bosch J., Retana J. (2004). Foraging behavior and pollinating effectiveness of *Osmia cornuta* (Hymenoptera: Megachilidae) and *Apis mellifera* (Hymenoptera: Apidae) on 'Comice' pear. *Apidologie* 35: 575-585.
- Moore J.L., Mouquet N., Lawton J.H., Loreau M. (2001). Coexistence, saturation and invasion resistance in simulated plant assemblages. *Oikos* 94: 303-314.
- Morandin L., Winston M. (2005). Wild bee abundance and seed production in conventional, organic, and genetically modified canola. *Ecological Applications* 15(3): 871-881.
- Moreira F. (1999). Relationships between vegetation structure and breeding bird densities in fallow cereal steppes in Castro Verde, Portugal. *Bird Study* 46: 309-318.
- Morel I., Wyss U., Collomb M., Butikofer U. (2006). Influence de la composition botanique de l'herbe ou du foin sur la composition du lait. *Revue suisse Agriculture* 38: 9-15.
- Moretto A.S., Distel R.A., Didoné N.G. (2001). Decomposition and nutrient dynamic of leaf litter and roots from palatable and unpalatable grasses in a semi-arid grassland. *Applied Soil Ecology* 18(1): 31-37.
- Morishima H., Oka H.J. (1979). Genetic diversity in rice populations of Nigeria: influence on community structure. *Agro-ecosystems* 5: 263-269.
- Morlat R., Jacquet A., Asselin C. (1993). Principaux effets de l'enherbement permanent contrôlé du sol, dans un essai de longue durée en Anjou. *Progrès Agricole et Viticole* 110: 406-410.
- Morse R.A., Calderone N.W. (2000). The value of honeybees as pollinators of US crops in 2000. *Bee Culture* 128(3): 1-15.
- Mortensen D.A., Coble H.D. (1989). Influence of soil water content on Common Cocklebur (*Xanthium strumarium*) interference in soybean (*Glycine max*). *Weed Science* 37: 76-83.
- Mougel C., Offre P., Ranjard L., Corberand T., Gamalero E., Robin C., Lemanceau P. (2006). Dynamic of the genetic structure of bacterial and fungal communities at different developmental stages of Medicago truncatula Gaertn. cv. Jemalong line J5. *New Phytologist* 170(1): 165-175.
- Moulin C.H., Girard N., Dedieu B. (2001). Apport de l'analyse fonctionnelle des systèmes d'alimentation. *Fourrages* 167: 337-363.
- Mouquet N., Leadley P., Mériguet J., Loreau M. (2004). Immigration and local competition in herbaceous plant communities: a three-year seed-sowing experiment. *Oikos* 104: 77-90.
- Mouquet N., Moore J.C., Loreau M. (2002). Plant species richness and community productivity: why the mechanism that promotes coexistence matters. *Ecology Letters* 5: 56-65.
- Moyer-Henry K.A., Burton J.W., Israel D.W., Ruffy T.W. (2006). Nitrogen transfer between plants: a 15N natural abundance study with crop and weed species. *Plant and Soil* 282: 7-20.
- Mulder C.P.H., Bazeley-White E., Dimitrakopoulos P.G., Hector A., Scherer-Lorenzen M., Schmid B. (2004). Species evenness and productivity in experimental plant communities. *Oikos* 107(1): 50-63.
- Mulder C.P.H., Jumpponen A., Högberg P., Huss-Danell K. (2002). How plant diversity and legumes affect nitrogen dynamics in experimental grassland communities. *Oecologia* 133: 412-421.
- Mulder C.P.H., Koricheva J., Huss-Danell K., Högberg P., Joshi J. (1999). Insects affect relationships between plant species richness and ecosystem processes. *Ecology Letters* 2: 237-246.
- Mulia R., Dupraz C. (2005). Unusual fine root distributions of two deciduous tree species observed in Southern France: what consequences for root dynamics modelling? *Plant and Soil* 281: 71-85.
- Muller C.B., Fellowes M.D.E., Godfray H.C.J. (2005). Relative importance of fertiliser addition to plants and exclusion of predators for aphid growth in the field. *Oecologia* 143(3): 419-427.
- Muller C.B., Godfray H.C.J. (1999). Predators and mutualists influence the exclusion of aphid species from natural communities. *Oecologia* 119(1): 120-125.
- Müller-Schärer H., Scheepens P.C., Greaves M.P. (2000). Biological control of weeds in European crops: recent achievements and future work. *Weed Research* 40(1): 83-98.
- Mundt C.C. (2002). Use of multiline cultivars and cultivar mixtures for disease management. *Annual Review of Phytopathology* 40: 381-410.
- Mundt C.C., Cowger C., Garrett K.A. (2002). Relevance of integrated disease management to resistance durability. *Euphytica* 124(2): 245-252.
- Munkvold L., Kjoller R., Vestberg M., Rosendahl S., Jakobsen I. (2004). High functional diversity within species of arbuscular mycorrhizal fungi. *New Phytologist* 164: 357-364.
- Muradian R. (2001). Ecological thresholds: a survey. *Ecological Economics* 38: 7-24.
- Mwangi P.N., Schmitz M., Scherber C., Roscher C., Schumacher J., Scherer-Lorenzen M., Weisser W.W., Schmid B. (2007). Niche pre-emption increases with species richness in experimental plant communities. *Journal of Ecology* 95: 65-78.
- Myers N. (1996). Environmental services of biodiversity. *Proceedings of the National Academy of Sciences of the United States of America* 93(7): 2764-2769.
- Naeem M., Compton S.G., Incoll L.D., Wright C., Corry D.T. (1997). Responses of aphids to a silvoarable agroforestry landscape. *Agroforestry Forum* 8(3): 18-20.
- Naeem S. (2002). Ecosystem consequences of biodiversity loss: the evolution of a paradigm. *Ecology* 83: 1537-1552.
- Naeem S., Thompson L.J., Lawler S.P., Lawton J.H., Woodfin R.M. (1994). Declining biodiversity can alter the performance of ecosystems. *Nature* 368: 734-737.
- Naeem S., Thompson L.J., Lawler S.P., Lawton J.H., Woodfin R.M. (1995). Biodiversity and ecosystem functioning: empirical evidence from experimental microcosms. *Philosophical Transactions of the Royal Society, London, B* 347: 249-262.
- Nakashima Y., Birkett M.A., Pye B.J., Powell W. (2006). Chemically mediated intraguild predator avoidance by aphid parasitoids: Interspecific variability in sensitivity to semiochemical trails of ladybird predators. *Journal of Chemical Ecology* 32(9): 1989-1998.
- Nauleau F. (1995). Nouvelles techniques d'entretien des sols viticoles - conséquences œnologiques. *Phytoma, La Défense des Végétaux* 478: 47-48.

- Nazzi F., Paoletti M.G., Lorenzoni G.G. (1989). Soil invertebrate dynamics of soybean agroecosystems encircled by hedgerows or not in Friuli, Italy. First data. *Agriculture, Ecosystems & Environment* 27(1-4): 163-176.
- Neff J.C., Asner G.P. (2001). Dissolved organic carbon in terrestrial ecosystems: synthesis and a model. *Ecosystems* 4: 29-48.
- Nentwig W. (1988). Augmentation of beneficial arthropods by strip-management. 1. Succession of predacious arthropods and long-term change in the ratio of phytophagous and predacious arthropods in a meadow. *Oecologia* 76(4): 597-606.
- Nentwig W. (1992). The promotive effect of weeds in sown strips on beneficial arthropods. *Zeitschrift Fur Pflanzenkrankheiten Und Pflanzenschutz* 13: 33-40.
- Nentwig W. (1998). Weedy plant species and their beneficial arthropods: potential for manipulation in field crops. In *Conservation Biological Control*, (Barbosa P., ed.), Academic Press, San Diego: 49-71.
- Nentwig W., Frank T., Lethmayer C. (1998). Sown weed strips: artificial ecological compensation areas as an important tool in conservation biological control. In *Conservation Biological Control*, (Barbosa P., ed.), Academic Press, San Diego: 133-153.
- Neuens F., Reheul D. (2001). Crop rotation versus monoculture; yield, N yield and ear fraction of silage maize at different levels of mineral N fertilization. *Njas-Wageningen Journal of Life Sciences* 49(4): 405-425
- Nicholls C.I., Altieri M.A. (2004). The agroecological bases of ecological engineering for pest management. In *Ecological engineering for pest management - Advances in habitat manipulation for Arthropods*, (Gurr G.M. et al., eds.), CSIRO Publishing - CABI Publishing, Collingwood - Oxon: 33-54.
- Nicodemo D., Nogueira-Couto R.H. (2004). Use of repellents for honeybees (*Apis mellifera* L.) in vitro in the yellow passion-fruit (*Passiflora edulis* Deg) crop and in confined beef cattle feeders. *Journal of Venomous Animals and Toxins including Tropical Diseases* 10(1): 77-85.
- Nicolaisen M.H., Risgaard-Petersen N., Revsbech N.P., Reichardt W., Ramsing N.B. (2004). Nitrification-denitrification dynamics and community structure of ammonia oxidizing bacteria in a high yield irrigated Philippine rice field. *FEMS Microbiology Ecology* 49(3): 359-369.
- Niehoff B., Poehling H.M. (1995). Population dynamics of aphids and syrphid larvae in winter wheat treated with different rates of pirimicarb. *Agriculture, Ecosystems & Environment* 52(1): 51-55.
- Niemala P., Tahvanainen J., Sorjonen J., Hokkanen T., Neuvonen S. (1982). The influence of host plant growth form and phenology on the life strategies of Finnish microlepidopterous larvae. *Oikos* 39: 164-170.
- Niezen J.H., Waghorn G.C., Graham T., Carter J.L., Leathwick D.M. (2002). The effect of diet fed to lambs on subsequent development of *Trichostrongylus colubriformis* larvae in vitro and on pasture. *Veterinary Parasitology* 105(4): 269-283.
- Niezen J.H., Waghorn T.S., Charleston W.A.G., Waghorn G.C. (1995). Growth and gastrointestinal nematode parasitism in lambs grazing either lucerne (*Medicago sativa*) or sulla (*Hedysarum coronarium*) which contains condensed tannins. *Journal of Agricultural Science* 125(2): 281-289.
- Nijs I., Roy J. (2000). How important are species richness, species evenness and interspecific differences to productivity? A mathematical model. *Oikos* 88: 57-66.
- Niklaus P.A., Leadley P.W., Schmid B., Körner C. (2001). A long-term field study on biodiversity x elevated CO2 interactions in grassland. *Ecological Monographs* 71(3): 341-356.
- Nilsson C. (1985). Impact of ploughing on emergence of pollen beetle parasitoids after hibernation. *Journal of Applied Entomology* 100: 302-308.
- Nilsson C. (2003). Parasitoids of pollen beetles. In *Biocontrol of Oilseed Rape Pests*, (Alford D.V., ed.), Blackwell Science, Oxford, UK: 73-85.
- Nocera J.J., Parsons G.J., Milton G.R., Fredeen A.H. (2005). Compatibility of delayed cutting regime with bird breeding and hay nutritional quality. *Agriculture, Ecosystems & Environment* 107(2-3): 245-253.
- Nohl W. (2001). Sustainable landscape use and aesthetic perception – preliminary reflections on future landscape aesthetics. *Landscape and Urban Planning* 54: 223-237.
- Norberg J., Swaney D.P., Dushoff J., Lin J., Casagrandi R., Levin S.A. (2001). Phenotypic diversity and ecosystem functioning in changing environments: a theoretical framework. *PNAS* 98(20): 11376-11381.
- Nordlund D.A., Chalfant R.B., Lewis W.J. (1984). Arthropod populations, yield and damage in monocultures and polycultures of corn, beans and tomatoes. *Agriculture, Ecosystems & Environment* 11(4): 353-367.
- Norris R., F. (2005). Ecological bases of interactions between weeds and organisms in other pest categories. *Weed Science* 53: 909-913.
- Norris R.F., Kogan M. (2005). Ecology of interactions between weeds and arthropods. *Annual Review of Entomology* 50: 479-503.
- Noss R.F. (1990). Indicators for monitoring biodiversity: a hierarchical approach. *Conservation Biology* 4(4): 355-364.
- Nozières M.O., Dulphy J.P., Peyraud J.L., Baumont R. (2006). Prediction of forage rumen protein degradability from an *in situ* degradability database, 21. *General Meeting of the European Grassland Federation*, Badajoz, Spain, Lloveras J., ed.: 531-533.
- Nuernberg K., Dannenberger D., Nuernberg G., Ender K., Voigt J., Scollan N.D., Wood J.D., Nute G.R., Richardson R.I. (2005). Effect of a grass-based and a concentrate feeding system on meat quality characteristics and fatty acid composition of longissimus muscle in different cattle breeds. *Livestock Production Science* 94: 137-147.
- Oades J.M. (1993). The Role of Biology in the Formation, Stabilization and Degradation of Soil Structure. *Geoderma* 56(1-4): 377-400.
- Oberholzer F., Frank T. (2003). Predation by the carabid beetles *Pterostichus melanarius* and *Poecilus cupreus* on slugs and slug eggs. *Biocontrol Science and Technology* 13(1): 99-110.
- Ockinger E., Smith H.G. (2007). Semi-natural grasslands as population sources for pollinating insects in agricultural landscapes. *Journal of Applied Ecology* 44(1): 50-59.
- O'Connors P.J., Smith S.E., Smith F.A. (2002). Arbuscular mycorrhizas influence plant diversity and community structure in a semiarid herbland. *New Phytologist* 154: 209-218.
- Odat N., Jetschke G., Hellwig F.H. (2004). Genetic diversity of *Ranunculus acris* L. (Ranunculaceae) populations in relation to species diversity and habitat type in grassland communities. *Molecular Ecology* 13: 1251-1257.
- Odhambo J.J.O., Bomke A.A. (2000). Short term nitrogen availability following overwinter cereal/grass and legume cover crop monocultures and mixtures in south coastal British Columbia. *Journal of Soil and Water Conservation* 55(3): 347-354.

- Oelmann Y., Wilcke W., Temperton V.M., Buchmann N., Roscher C., Schumacher J., Schulze E.-D., Weisser W.W. (2007). Soil and Plant Nitrogen Pools as Related to Plant Diversity in an Experimental Grassland. *Soil Science Society of America Journal* 71(3): 720-729.
- Offre P., Pivato B., Siblot S., Gamalero E., Corberand T., Lemanceau P., Mougél C. (2007). Identification of bacterial groups preferentially associated with mycorrhizal roots of *Medicago truncatula*. *Applied and Environmental Microbiology* 73(3): 913-921.
- Ogren R.E., Kawakatsu M. (1998). American Nearctic and Neotropical land planarian (Tricladida : Terricola) faunas. *Pedobiologia* 42(5-6): 441-451.
- Ohgushi T., Sawada H. (1985). Arthropod predation limits the population density of an herbivorous lady beetle, *Henosepilachna niponica* (Lewis). *Researches on Population Ecology* 27(2): 351-359.
- Oksanen L., Oksanen T. (2000). The logic and realism of the hypothesis of exploitation ecosystems. *American Naturalist* 155(6): 703-723.
- Oldroyd G.E.D. (2001). Dissecting symbiosis: Developments in Nod factor signal transduction. *Annals of Botany* 87(6): 709-718.
- Olesen J.M., Bascompte J., Dupont Y.L., Jordano P. (2007). The modularity of pollination networks. *PNAS* 104(50): 19891-19896.
- Olkowski W., Zhang A. (1998). Habitat management for biological control, examples from China. In *Enhancing biological control - Habitat management to promote natural enemies of agricultural pests*, (Pickett C.H., Bugg R.L., eds.), University of California Press, Berkeley, Los Angeles, London: 255-270.
- Ollerton J. (1996). Reconciling ecological processes with phylogenetic patterns: the apparent paradox of plant-pollinator systems. *Journal of Ecology* 84: 767-769.
- Olofsson J., Oksanen L. (2002). Role of litter decomposition for increased primary productivity in areas heavily grazed by reindeer: a litterbag experiment *Oikos* 96: 507-515.
- Olsson G.A., Ronningen K. (1999). Environmental values in traditional agricultural landscapes in Norway. *Landbruksökonomisk Forum* 16(4).
- O'Neal M.E., Zontek E.L., Szendrei Z., Landis D.A., Isaacs R. (2005). Ground predator abundance affects prey removal in highbush blueberry (*Vaccinium corymbosum*) fields and can be altered by aisle ground covers. *BioControl* 50(2): 205-222.
- Oreszczytn S. (2000). A systems approach to the research of people's relationships with English hedgerows. *Landscape and Urban Planning* 50: 107-117.
- Orth D., Loiseau P., Loisel A., Perrin O., Balay C. (2004). Un cas d'évaluation physionomique de la biodiversité: options et questions. *Fourrages* 179: 335-352.
- Osoro K., Mateos-Sanz A., Frutos P., Garcia U., Ortega-Mora L.M., Ferreira L.M.M., Celaya R., Ferre I. (2007). Anthelmintic and nutritional effects of heather supplementation on Cashmere goats grazing perennial ryegrass-white clover pastures. *Journal of animal science* 85(3): 861-870.
- Ostman O. (2004). The relative effects of natural enemy abundance and alternative prey abundance on aphid predation rates. *Biological Control* 30(2): 281-287.
- Ostman O., Ekbom B., Bengtsson J. (2001). Landscape heterogeneity and farming practice influence biological control. *Basic and Applied Ecology* 2(4): 365-371.
- Ostman O., Ekbom B., Bengtsson J., Weibull A.C. (2001). Landscape complexity and farming practice influence the condition of polyphagous carabid beetles. *Ecological Applications* 11(2): 480-488.
- Otte A., Simmering D., Wolters V. (2007). Biodiversity at the landscape level: recent concepts and perspectives for multifunctional land use. *Landscape Ecology* 22: 639-642.
- Otway S., Hector A., Lawton J.H. (2005). Resource dilution effects on specialist insect herbivores in a grassland biodiversity experiment. *Journal of Animal Ecology* 74: 234-240.
- Paine T.D., Millar J.G. (2005). Invasion biology and management of insects feeding on eucalyptus in California. In *Entomological research in Mediterranean forest ecosystems*, (Lieutier F., Ghaïoule D., eds.), Versailles France: 221-228.
- Palmberg C., Scherer-Lorenzen M., Jumpponen A., Carlsson G., Huss-Danell K., Högberg P. (2005). Inorganic soil nitrogen under grassland plant communities of different species composition and diversity. *Oikos* 110: 271-282.
- Palmer M., Maurer T.A. (1997). Does diversity beget diversity: a case study of crops and weeds. *Journal of Vegetation Science* 8: 235-240.
- Panter K.E., James L.F. (1989). Death camas-early grazing can be hazardous. *Rangelands* 11: 147-149.
- Paoletti M.G., Hu D.X., Marc P., Huang N.X., Wu W.L., Han C.R., He J.H., Cai L.W. (1999). Arthropods as bioindicators in agroecosystems of Jiang Han Plain, Qianjiang City, Hubei China. *Critical Reviews in Plant Sciences* 18(3): 457-465.
- Paolini V., De La Farge F., Prevot F., Dorchie P., Hoste H. (2005). Effects of the repeated distribution of sainfoin hay on the resistance and the resilience of goats naturally infected with gastrointestinal nematodes. *Veterinary Parasitology* 127(3-4): 277-283.
- Paolini V., Fouraste I., Hoste H. (2004). In vitro of tree woody plant and sainfoin extracts on 3rd- stage larvae and adult worms of three gastrointestinal nematodes. *Parasitology* 129: 69-77.
- Parisi L., Didelot F., Brun L. (2003). Raisonner la lutte contre la tavelure du pommier, un enjeu majeur pour une arboriculture durable, *Septième Conférence Internationale sur les Maladies des Plantes*, Tours, France, 3-4-5 décembre 2003, Plantes A.F.d.P.d., ed.
- Park R.J. (1972). Flavour differences in meat from lambs grazed on lucerne (*Medicago sativa*) or phalaris (*Phalaris tuberosa*) pastures. *Journal of Agricultural Science* 78: 47-52.
- Parkin T.B., Kaspar T.C. (2006). Nitrous oxide emissions from corn-soybean systems in the Midwest. *Journal of Environmental Quality* 35(4): 1496-1506.
- Pastor J., Cohen Y. (1997). Herbivores, the Functional Diversity of Plant Species, and the Cycling of Nutrients in Ecosystems. *Theoretical Population Biology* 51: 165-179.
- Pastor J., Dewey B., Naiman R.J., McInnes P.F., Cohen Y. (1993). Moose Browsing and Soil Fertility in the Boreal Forests of Isle Royale National Park *Ecology* 74: 467-480.
- Patra A.K., Abbadie L., Clays-Josserand A., Degrange V., Grayston S.J., Guillaumaud N., Loiseau P., Louault F., Mahmood S., Nazaret S., Philippot L., Poly F., Prosser J.I., Le Roux X. (2006). Effects of management regime and plant species on the enzyme activity and genetic structure of N-fixing, denitrifying and nitrifying bacterial communities in grassland soils. *Environmental Microbiology* 8(6): 1005-1016.

- Paulsen H.M., Schochow M., Ulber B., Kuhne S., Rahmann G. (2006). Mixed cropping systems for biological control of weeds and pests in organic oilseed crops. *Aspects of Applied Biology* 79: 215-219.
- Pell J.K., Vandenberg J.D. (2002). Interactions among the aphid *Diuraphis noxia*, the entomopathogenic fungus *Paecilomyces fumosoroseus* and the coccinellid *Hippodamia convergens*. *Biocontrol Science and Technology* 12(2): 217-224.
- Peng R.K., Christian K., Gibb K. (1999). The effect of colony isolation of the predacious ant, *Oecophylla smaragdina* (F.) (Hymenoptera: Formicidae), on protection of cashew plantations from insect pests. *International Journal of Pest Management* 45(3): 189-194.
- Peng R.K., Incoll L.D., Sutton S., Wright L.C., Chadwick A. (1993). Diversity of airborne arthropods in a silvoarable agroforestry system. *Journal of Applied Ecology* 30(3): 551-562.
- Peoples M.B., Herridge D.F., Ladha J.K. (1995). Biological dinitrogen fixation: an efficient source of nitrogen for sustainable agriculture production. *Plant and Soil* 174: 3-28.
- Pères G., Cluzeau D., Curmi P., Hallaire V. (1998). Earthworm activity and soil structure changes due to organic enrichments in vineyard systems. *Biology and Fertility of Soils* 27(4): 417-424.
- Pérez-de-Luque A.J., Sillero C., Moral A., Cubero J.I., Rubiales D. (2004). Effect of sowing date and host resistance on the establishment and development of *Orobanche crenata* in faba bean and common vetch. *Weed Research* 44(4): 282-288.
- Perez-Piqueres A., Edel-Hermann W., Alabouvette C., Steinberg C. (2006). Response of soil microbial communities to compost amendments. *Soil Biology & Biochemistry* 38(3): 460-470.
- Perfecto I., Vandermeer J.H., Lopez Bautista G., Ibarra Nunez G., Greenberg R., Bichier P., Langridge S. (2004). Greater predation in shaded coffee farms: the role of resident neotropical birds. *Ecology* 85(10): 2677-2681.
- Personeni E., Loiseau P. (2005). Species strategy and N fluxes in grassland soil - A question of root litter quality or rhizosphere activity? *European Journal of Agronomy* 22(2): 217-229.
- Pesson P., Louveaux J. (1984). Pollinisation et productions végétales, Editions Inra Paris, 663 p.
- Petanidou T. (2003). Introducing plants for bee-keeping at any cost? – Assessment of *Phacelia tanacetifolia* as nectar source plant under xeric Mediterranean conditions. *Plant Systematics and Evolution* 238(1-4): 155-168.
- Peters D.P.C., Pielke R.A.S., Bestelmeyer B.T., Allen C.D., Munson-McGee S., Havstad K.M. (2004). Cross-scale interactions, nonlinearities, and forecasting catastrophic events. *Proceedings of the National Academy of Sciences (U.S.A.)* 101: 15130-15153.
- Peterson T.A., Shapiro C.A., Flowerday A.D. (1990). Rainfall and previous crop effects on crop yields. *American journal of alternative agriculture* 5: 33-37.
- Petrie G.A., Seguin-Swartz G., Gugel R.K. (1995). Latent infection of Brassicaceae in the field by *Leptosphaeria maculans* (blackleg). *Canadian journal of plant pathology* 17(1): 75-81.
- Petsikos C., Dalias P., Troumbis A.Y. (2007). Effects of *Oxalis pes-caprae* L. invasion in olive groves *Agriculture, Ecosystems & Environment* 120(2-4): 325-329.
- Pfammatter W., Vuignier R. (1998). Amélioration de la lutte biologique dans les cultures fruitières au moyen de bandes de plantes sauvages, *Colloque transnational sur Les luites biologique, intégrée et raisonnée. Bilan et perspectives au regard d'expériences régionales et européennes*, Lille, France, 21-23 janv. 98, Nord-Pas-de-Calais R., ed.: 71-72.
- Pfiffner L., Luka H. (2000). Overwintering of arthropods in soils of arable fields and adjacent semi-natural habitats. *Agriculture, Ecosystems & Environment* 78(3): 215-222.
- Pfiffner L., Wyss E. (2004). Use of sown wildflower strips to enhance natural enemies of agricultural pests. In *Ecological engineering for pest management - Advances in habitat manipulation for Arthropods*, (Gurr G.M. et al., eds.), CSIRO Publishing - CABI Publishing, Collingwood - Oxon: 165-186.
- Pfisterer A.B., Balvanera P., Buchmann N., He J.-S., Nakashizuka T., Raffaelli D., Schmid B. (2005). The Role of Biodiversity for Ecosystem Services: Current Knowledge, SCOPE, SCNAT: 33 p. http://www.biodiversity.ch/about_us/activities/synthesis_project/documents/PfistererEtalWhitePaper2005.pdf
- Pfisterer A.B., Diemer M., Schmid B. (2003). Dietary shift and lowered biomass gain of a generalist herbivore in species poor experimental plant communities. *Oecologia* 135: 234-241.
- Pfisterer A.B., Joshi J., Schmid B., Fischer M. (2004). Rapid decay of diversity-productivity relationships after invasion of experimental plant communities *Basic and Applied Ecology* 5: 5-14.
- Pfisterer A.B., Schmid B. (2002). Diversity-dependent production can decrease the stability of ecosystem functioning. *Nature* 416: 84-86.
- Pfimplin A. (1993). Conduite et utilisation des associations graminées - trèfle blanc. *Fourrages* 135: 407-428.
- Philippot L., Kuffner M., Cheneby D., Depret G., Laguerre G., Martin-Laurent F. (2006). Genetic structure and activity of the nitrate-reducers community in the rhizosphere of different cultivars of maize. *Plant and Soil* 287(1-2): 177-186.
- Pichot A. (1991). Evolution de la flore adventice et de son contrôle en Beauce depuis 50 ans. *Compte Rendus de l'Académie Françaises* 77(2): 117-127.
- Pickett C.H., Bugg R.L. (1998). Enhancing biological control - Habitat management to promote natural enemies of agricultural pests, University of California Press, Berkeley - Los Angeles - London, 422 p.
- Pielke Sr. R.A. (2001). Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall. *Reviews of Geophysics* 39: 151-177.
- Pielke Sr. R.A., Avissar R. (1990). Influence of landscape structure on local and regional climate. *Landscape Ecology* 4(2/3): 133-155.
- Piggin C.M. (1982). The biology of Australian weeds. 8. *Echium plantagineum* L. *Journal of the Australian Institute of Agricultural Science* 48: 3-16.
- Pimentel D. (1961). Species diversity and insect population outbreaks. *Annals of the Entomological Society of America* 54: 76-86.
- Pimentel D., Hepperly P., Hanson J., Douds D., Seidel R. (2005). Environmental, energetic, and economic comparisons of organic and conventional farming systems. *BioScience* 55(7): 573-582.
- Pimentel D., Wilson C., McCullum C., Huang R., Dwen P., Flack J., Tran Q., Saltman T., Cliff B. (1997). Economic and Environmental Benefits of Biodiversity. *BioScience* 47: 747-757.
- Pimentel D., Zuniga R., Morrison D. (2005). Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics* 52: 273-288.

- Pineda A., Morales I., Marcos-Garcia M.A., Fereres A. (2007). Oviposition avoidance of parasitized aphid colonies by the syrphid predator *Epiplatys balteatus* mediated by different cues. *Biological Control* 42: 274-280.
- Piotrowski J.S., Denich T., Klironomos J.N., Graham J.M., Rillig M.C. (2004). The effects of arbuscular mycorrhizas on soil aggregation depend on the interaction between plant and fungal species. *New Phytologist* 164(2): 365-373.
- Piper J.K. (1999). Natural systems agriculture. In *Biodiversity in agroecosystems*, (Collins W.W., Qualset C.O., eds.), CRC Press, Boca Raton: 167-196.
- Piutti S., Hallet S., Rousseaux S., Philippot L., Soulas G., Martin-Laurent F. (2002). Accelerated mineralisation of atrazine in maize rhizosphere soil. *Biology and Fertility of Soils* 36(6): 434-441.
- Plancquaert P. (1990). Création des prairies temporaires et artificielles. In *Encyclopédie des techniques agricoles*, Techniques Agricoles, Paris: fasc. 2220, 1-19.
- Platt J.O., Caldwell J.S., Kok L.T. (1999). Effect of buckwheat as a flowering border on populations of cucumber beetles and their natural enemies in cucumber and squash. *Crop Protection* 18(5): 305-313.
- Pointereau P., Bazile D. (1995). Arbres des champs : haies, alignements, prés, vergers ou l'art du bocage, Solagro, Toulouse, 150 p.
- Polis G.A., Holt R.D. (1992). Intraguild predation: the dynamics of complex trophic interactions. *Trends in Ecology & Evolution* 7: 151-154.
- Pollard K.A., Holland J.M. (2006). Arthropods within the woody element of hedgerows and their distribution pattern. *Agricultural and Forest Entomology* 8(3): 203-211.
- Polley H.W., Wilsey B.J., Derner J.D. (2003). Do species evenness and plant density influence the magnitude of selection and complementarity effects in annual plant mixtures. *Ecology Letters* 6: 248-256.
- Pontes L.D.S., Soussana J.F., Louault F., Andueza D., Carrere P. (2007). Leaf traits affect the above-ground productivity and quality of pasture grasses. *Functional ecology* 21: 844-853.
- Pontin D.R., Wade M.R., Kehrl P., Wratten S.D. (2006). Attractiveness of single and multiple species flower patches to beneficial insects in agroecosystems. *Annals of Applied Biology* 148(1): 39-47.
- Porter P.M., Huggins D.R., Perillo C.A., Quiring S.R., Crookston R.K. (2003). Organic and other management strategies with two and four-year crop rotations in Minnesota. *Agronomy Journal* 95: 233-244.
- Porter P.M., Lauer J.G., Lueschen W.E., Ford J.H., Hoverstad T.R., Oplinger E.S., Crookston R.K. (1997). Environment affects the corn and soybean rotation effect. *Agronomy Journal* 89: 441-448.
- Postma-Blaauw M.B., Bloem J., Faber J.H., van Groenigen J.W., de Goede R.G.M., Brussaard L. (2006). Earthworm species composition affects the soil bacterial community and net nitrogen mineralization. *Pedobiologia* 50(3): 243-256.
- Potts S., Petanidou T., Roberts S., O'Toole C., Hulbert A., Willmer P.G. (2006). Plant-pollinator biodiversity and pollination services in a complex mediterranean landscape. *Biological Conservation* 129(4): 519-529.
- Potts S.G., Vulliamy B., Dafni A., Ne'eman G., O'Toole C., Roberts S., Willmer P.G. (2003). Response of plant-pollinator communities following fire: changes in diversity, abundance and reward structure. *Oikos* 101: 103-112.
- Potts S.G., Vulliamy B., Roberts S., O'Toole C., Dafni A., Ne'eman G., Willmer P.G. (2004). Nectar resource diversity organises flower-visitor community structure. *Entomologia Experimentalis et Applicata* 113: 103-107.
- Potvin C., Chapin F.S.I., Gonzalez A., Leadley P., Reich P., Roy J. (2007). Plant Biodiversity and Responses to Elevated Carbon Dioxide. In *Terrestrial Ecosystems in a Changing World*, (Canadell J. et al., eds.), Springer-Verlag: 125-134.
- Poudel D.D., Horwath W.R., Lanini W.T., Temple S.R., van Bruggen A.H.C. (2002). Comparison of soil N availability and leaching potential, crop yields and weeds in organic, low-input and conventional farming systems in northern California. *Agriculture, Ecosystem and Environment* 90(2): 125-137.
- Powell W., Pickett J.A. (2003). Manipulation of parasitoids for aphid pest management: progress and prospects. *Pest Management Science* 59(2): 149-155.
- Prache S., Gordon I.J., Rook A.J. (1998). Foraging behaviour and diet selection in domestic herbivores. *Annales de Zootechnie* 47: 335-345.
- Prache S., Priolo A., Grolier P. (2003). Persistence of carotenoid pigments in the blood of concentrate-finished grazing sheep: Its significance for the traceability of grass-feeding. *Journal of animal science* 81: 360-367.
- Prasad R.P., Snyder W.E. (2004). Predator interference limits fly egg biological control by a guild of ground-active beetles. *Biological Control* 31(3): 428-437.
- Prasad R.P., Snyder W.E. (2006). Diverse trait-mediated interactions in a multi-predator, multi-prey community. *Ecology* 87(5): 1131-1137.
- Prasad R.P., Snyder W.E. (2006). Polyphagy complicates conservation biological control that targets generalist predators. *Journal of Applied Ecology* 43(2): 343-352.
- Prasifka J.R., Schmidt N.P., Kohler K.A., O'Neal M.E., Hellmich R.L., Singer J.W. (2006). Effects of living mulches on predator abundance and sentinel prey in a corn-soybean-forage rotation. *Environmental Entomology* 35(5): 1423-1431.
- Prévosto B., Dambrine E., Coquillard P., Robert A. (2006). Broom (*Cytisus scoparius*) colonization after grazing abandonment in the French Massif Central: impact on vegetation composition and resource availability. *Acta Oecologica* 30(2): 258-268.
- Prew R.D., Dyke G.V. (1979). Experiments comparing "break crops" as a preparation for winter wheat followed by spring barley. *Journal of Agricultural Science* 92: 189-201.
- Prieur-Richard A.-H., Lavorel S. (2000). Invasions : the perspective of diverse plant communities - a review. *Austral Ecology* 25(1): 1-7.
- Prieur-Richard A.-H., Lavorel S., Grigulis K., Dos Santos A. (2000). Plant community diversity and invasibility by exotics: invasion of Mediterranean old fields by *Conyza bonariensis* and *Conyza canadensis*. *Ecology Letters* 3(5): 412-422.
- Prieur-Richard A.-H., Lavorel S., Linhardt Y., Dossantos A. (2002). Plant diversity, herbivory and resistance of plant community to invasion in Mediterranean annual communities. *Oecologia* 130: 93-104.
- Priolo A., Vasta V. (2007). Effects of tannin containing diets on small ruminant meat quality. *Italian Journal of animal science* 6(suppl 1): 527-530.
- Proctor M., Yeo P., Lack A. (1996). The natural history of pollination, Timber Press, Portland, OR, 487 p.
- Prokopy R.J., Mason J.L., Christie M., Wright S.E. (1995). Arthropod pest and natural enemy abundance under second-level versus first-level integrated pest management practices in apple orchards: a 4-year study. *Agriculture, Ecosystems & Environment* 57(1): 35-47.

- Propp G.D., Morgan P.B. (1985). Effect of host distribution on parasitoidism of house-fly (Diptera: Muscidae) pupae by *Spalangia* spp. and *Muscidifurax raptor* (Hymenoptera: Pteromalidae). *Canadian Entomologist* 117(5): 515-524.
- Pullaro T.C., Marino P.C., Jackson D.M., Harrison H.F., Keinath A.P. (2006). Effects of killed cover crop mulch on weeds, weed seeds, and herbivores. *Agriculture, Ecosystems & Environment* 115(1-4): 97-104.
- Pulleman M.M., Six J., Uyl A., Marinissen J.C.Y., Jongmans A.G. (2005). Earthworms and management affect organic matter incorporation and microaggregate formation in agricultural soils. *Applied Soil Ecology* 29(1): 1-15.
- Putman R.J., Moore N.P. (1998). Impact of deer in lowland Britain on agriculture, forestry and conservation habitats. *Mammal Review* 28(4): 141-163.
- Pyšek P., Müllerová J., Jarošík V. (2007). Historical dynamics of *Heracleum mantegazzianum* invasion at regional and local scales. In *Ecology and Management of Giant Hogweed (Heracleum mantegazzianum)*, (Pyšek P. et al., eds.), CAB International: 42-54.
- Pywell R.F., Bullock J.M., Walker K.J., Coulson S.J., Gregory S.J., Stevenson M.J. (2004). Facilitating grassland diversification using the hemiparasitic plant *Rhinanthus minor*. *Journal of Applied Ecology* 41(5): 880-887.
- Pywell R.F., James K.L., Herbert I., Meek W.R., Carvell C., Bell D., Sparks T.H. (2005). Determinants of overwintering habitat quality for beetles and spiders on arable farmland. *Biological Conservation* 123(1): 79-90.
- Pywell R.F., Warman E.A., Hulmes L., Hulmes S., Nuttall P., Sparks T.H., Critchley C.N.R., Sherwood A. (2006). Effectiveness of new agri-environment schemes in providing foraging resources for bumblebees in intensively farmed landscapes. *Biological Conservation* 129(2): 192-206.
- Quested H.M., Callaghan T.V., Cornelissen J.H.C., Press M.C. (2005). The impact of hemiparasitic plant litter on decomposition: direct, seasonal and litter mixing effects. *Journal of Ecology* 93(1): 87-98.
- Quested H.M., Eriksson O., Fortunel C., Garnier E. (2007). Plant traits relate to whole-community litter quality and decomposition following land use change. *Functional Ecology* 21(6): 1016 - 1026.
- Quétiér F., Lavorel S., Thuiller W., Davies I.D. (2007). Plant trait-based assessment of ecosystem service sensitivity to land-use change in mountain grasslands. *Ecological Applications* 17: 2377-2386.
- Quétiér F., Rivoal F., Marty P., de Chazal J., Lavorel S. (soumis). Descriptions of a mountain grassland landscape by locals and visitors: an inquiry into narratives on ecosystem services. *Regional Environmental Change*.
- Quimby P.C., Bruckart W.L., Deloach C., J., Knutson L., Ralphs M.H. (1991). Biological control of rangeland weeds. In *Noxious Range Weeds*, Westview Press, Boulder: 83-102.
- Raaijmakers J.M., Weller D.M. (1998). Natural plant protection by 2,4-diacetylphloroglucinol - producing *Pseudomonas* spp. in take-all decline soils. *Molecular Plant-Microbes Interactions* 11: 144-152.
- Radosevich S.R., Holt J.S. (1984). *Weed Ecology: implications for vegetation management*, Wiley & Sons, New York, 265 p.
- Raghua S., Dhileepan K., Scanlan J.C. (2007). Predicting risk and benefit a priori in biological control of invasive plant species: A systems modelling approach. *Ecological Modelling* 208(2-4): 247-262.
- Rajcan I., Swanton C. (2001). Understanding maize-weed competition: resource competition, light quality and the whole plant. *Field Crops Research* 71: 139-150.
- Ramdhani N., Bux F. (2007). Functional characterization of heterotrophic denitrifying bacteria in activated sludge. *South African Journal of Science* 103(3-4): 113-116.
- Ranjard L., Lejon D.P.H., Mougel C., Schehrer L., Merdinoglu D., Chaussod R. (2003). Sampling strategy in molecular microbial ecology: influence of soil sample size on DNA fingerprinting analysis of fungal and bacterial communities. *Environmental Microbiology* 5(11): 1111-1120.
- Rao M.R., Singh M.P., Day R. (2000). Insect pest problems in tropical agroforestry systems: contributory factors and strategies for management. *Agroforestry Systems* 50(3): 243-277.
- Rasmont P., Ebmer P.A., Banaszak J., van der Zanden G. (1995). Hymenoptera Apoidea Gallica. Liste taxonomique des abeilles de France, de Belgique, de Suisse et du Grand-Duché de Luxembourg. *Bulletin de la Société entomologique de France* 100(HS): 98 p.
- Rasolo F., Raunet M. (1998). Gestion agrobiologique des sols et des systèmes de culture, *Gestion agrobiologique des sols et des systèmes de culture. Atelier international, Antsirabe, 1998/03/23-28*, CIRAD: 658 p.
- Raworth D.A., Robertson M.C., Bittman S. (2004). Effects of dairy slurry application on carabid beetles in tall fescue, British Columbia, Canada. *Agriculture, Ecosystems & Environment* 103(3): 527-534.
- Raymond B., Darby A.C., Douglas A.E. (2000). Intraguild predators and the spatial distribution of a parasitoid. *Oecologia* 124(3): 367-372.
- Rea J.H., Wratten S.D., Sedcole R., Cameron P.J., Davis S.I., Chapman R.B. (2002). Trap cropping to manage green vegetable bug *Nezara viridula* (L.) (Heteroptera: Pentatomidae) in sweet corn in New Zealand. *Agricultural and Forest Entomology* 4(2): 101-107.
- Reau R., Bodet J.M., Bordes J.P., Dore T., Ennaifar S., Moussart A., Nicolardot B., Pellerin S., Plenchette C., Quinsac A., Sausse C., Seguin B., Tivoli B. (2005). Effets allélopathiques des Brassicacées via leurs actions sur les agents pathogènes telluriques et les mycorhizes : analyse bibliographique. *OCL - Oleagineux, Corps Gras, Lipides* 12(3): 314-319.
- Rebek E.J., Sadof C.S., Hanks L.M. (2005). Manipulating the abundance of natural enemies in ornamental landscapes with floral resource plants. *Biological Control* 33(2): 203-216.
- Recous S. (2005). Soil microbial biomass: its role in nitrogen cycling and efficiency, Cambridge, 2005/12/14-16, IFS Proceedings 566: 20 p.
- Reeleder R.D. (2003). Fungal plant pathogens and soil biodiversity. *Canadian Journal of Soil Science* 83(3): 331-336.
- Reich P.B., Knops J., Tilman D., Craine J., Ellsworth D., Tjoelker M.G., Lee T.D., Wedin D., Naeem S., Bahaeddin D., Hendrey G., Jose S., Wrage K., Goth J., Bengtson W. (2001). Plant diversity enhances ecosystem responses to elevated CO₂ and nitrogen deposition. *Nature* 410: 809-812.
- Rengel Z., Ross G., Hirsch P. (1998). Plant genotype and micronutrient status influence colonization of wheat roots by soil bacteria. *Journal of Plant Nutrition* 21(1): 99-113.
- Restrepo-Saenz C., Muhammad I., Harvey C., Harmand J.M., Morales J. (2004). Relationships between tree cover in pastures and cattle productivity in livestock farms in the dry tropics, Canas, Costa Rica. *Agroforestería en las Américas* 41/42: 29-36.

- Reznik S.Y. (1991). The effects of feeding damage in ragweed *Ambrosia artemisiifolia* (Asteraceae) on populations of *Zygogramma suturalis* (Coleoptera, Chrysomelidae). *Oecologia* 88(2): 204-210.
- Rhainds M., English-Loeb G. (2003). Testing the resource concentration hypothesis with tarnished plant bug on strawberry: density of hosts and patch size influence the interaction between abundance of nymphs and incidence of damage. *Ecological Entomology* 28(3): 348-358.
- Rhine E.D., Fuhrmann J.J., Radosevich M. (2003). Microbial community responses to atrazine exposure and nutrient availability: Linking degradation capacity to community structure. *Microbial Ecology* 46(2): 145-160.
- Ricketts T.H., Daily G.C., Ehrlich P.R., Michener C.D. (2004). Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences of the United States of America* 101: 12579-12582.
- Ricklefs R.E., Miller T.E. (2005). Ecology, De Boeck University, Paris, 821 p.
- Ridsdill-Smith T.J. (1981). Some effects of three species of dung beetles (Coleoptera: Scarabaeidae) in south-western Australia on the survival of the bush fly *Musca vetustissima* Walker (Diptera: Muscidae) in dung pads. *Bulletin of Entomological Research* 71(3): 425-433.
- Ridsdill-Smith T.J., Matthiessen J.N. (1988). Bush fly *Musca vetustissima* (Diptera: Muscidae) control in relation to seasonal abundance of scarabaeine dung beetles (Coleoptera: Scarabaeidae) in south-western Australia. *Bulletin of Entomological Research* 78(4): 633-639.
- Riechert S.E., Bishop L. (1990). Prey control by an assemblage of generalist predators: spiders in garden test systems. *Ecology* 71(4): 1441-1450.
- Rieux R., Simon S., Defrance H. (1999). Role of hedgerows and ground cover management on arthropod populations in pear orchards. *Agriculture, Ecosystems & Environment* 73(2): 119-127.
- Riley R.H., Brick M.A., Conniff K.L., Wood D.R., Pearson C.H., Chapman P.L. (1993). Response of Seed Yield and Yield Components in Mixtures of Dry Edible Bean. *Field Crops Research* 33(3): 283-292.
- Rillig M.C. (2004). Arbuscular mycorrhizae and terrestrial ecosystem processes. *Ecology Letters* 7: 740-754.
- Rillig M.C., Lutgen E.R., Ramsey P.W., Klironomos J.N., Gannon J.E. (2005). Microbiota accompanying different arbuscular mycorrhizal fungal isoates influence soil aggregation. *Pedobiologia* 49: 251-259.
- Rillig M.C., Mummey D.L., Ramsey P.W., Klironomos J.N., Gannon J.E. (2006). Phylogeny of arbuscular mycorrhizal fungi predicts community composition of symbiosis-associated bacteria. *FEMS Microbiology Ecology* 57(3): 389-395.
- Rillig M.C., Wright S.F., Eviner V.T. (2002). The role of arbuscular mycorrhizal fungi and glomalin in soil aggregation: comparing effects of five plant species. *Plant and Soil* 238: 325-333.
- Risch S.J. (1981). Insect herbivore abundance in tropical monocultures and polycultures: an experimental test of two hypotheses. *Ecology* 62: 1325-1340.
- Risch S.J., Andow D.A., Altieri M.A. (1983). Agroecosystem diversity and pest control: data, tentative conclusions and new research directions. *Environmental Entomology* 12: 625-629.
- Ritchie M.E., Tilman D., Knops J.H.M. (1998). Herbivore Effects on Plant and Nitrogen Dynamics in Oak Savanna *Ecology* 79: 165-177.
- Rixen C., Huovinen C., Huovinen K., Stöckli V., Schmid B. (2008). A plant diversity x water chemistry experiment in subalpine grassland. *Perspectives in Plant Ecology, Evolution and Systematics* 10(1): 51-61.
- Rizvi S.J., Rizvi V. (1992). Allelopathy: basic and applied aspects Chapman & Hall, 504 p.
- Robertson L.N. (1986). Experimental studies of predation on grassland populations of Australian soldier fly, *Inopus rubriceps* (Macquart) (Diptera: Stratiomyidae). *New Zealand Journal of Zoology* 13(1): 75-81.
- Robson M.C., Fowler S.M., Lampkin N.H., Leifert C., Leitch M., Robinson D., Watson C.A., Litterick A.M. (2002). The agronomic and economic potential of break crops for ley/arable rotations in temperate organic agriculture. *Advances in Agronomy* 77: 369-427.
- Rochette P., Janzen H.H. (2005). Towards a revised coefficient for estimating N₂O emissions from legumes. *Nutrient cycling in agroecosystems* 73(2/3): 171-179.
- Rochon J.J., Doyle C.J., Greef J.M., Hopkins A., Molle G., Sitzia M., Scholefield D., Smith C.J. (2004). Grazing legumes in Europe: a review of their status, management, benefits, research needs and future prospects. *Grass & Forage Science* 59: 197-214.
- Roder W., Mason, S. C., Clegg M.D., Kniep K.R. (1989). Yield-soil water relationships in sorghum-soybean cropping systems with different fertilizer regimes. *Agronomy Journal* 81(3): 470-475.
- Rodrigues A., Andueza D., Picard F., Cecato U., Farruggia A., Baumont R. (2007). Valeur alimentaire et composition floristique des prairies permanentes : premiers résultats d'une étude conduite dans le Massif Central, 14. *Rencontres Recherches Ruminants*: 241.
- Rodriguez M.A., Hawkins B.A. (2000). Diversity, function and stability in parasitoid communities. *Ecology Letters* 3(1): 35-40.
- Roesch L.F.W., Camargo F.A.O., Bento F.M., Triplett E.W. (2008). Biodiversity of diazotrophic bacteria within the soil, root and stem of field-grown maize. *Plant and Soil* 302(1-2): 91-104.
- Roininen H., Price P.W., Tahvanainen J. (1996). Bottom-up and top-down influences in the trophic system of a willow, a galling sawfly, parasitoids and inquilines. *Oikos* 77(1): 44-50.
- Roose T., Fowler A.C. (2004). A mathematical model for water and nutrient uptake by plant root systems. *Journal of Theoretical Biology* 228: 173-184.
- Root R.B. (1973). Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecological Monographs* 43(1): 95-124.
- Roschewitz I., Hucker M., Tschamtké T., Thies C. (2005). The influence of landscape context and farming practices on parasitism of cereal aphids. *Agriculture, Ecosystems & Environment* 108(3): 218-227.
- Rosecrance R.C., McCarty G.W., Shelton D.R., Teasdale J.R. (2000). Denitrification and N mineralization from hairy vetch (*Vicia villosa* Roth) and rye (*Secale cereale* L.) cover crop monocultures and bicultures. *Plant and Soil* 227(1-2): 283-290.
- Rosenheim J.A., Wilhoit L.R., Armer C.A. (1993). Influence of intraguild predation among generalist insect predators on the suppression of an herbivore population. *Oecologia* 96(3): 439-449.
- Rosi-Marshall E.J., Tank J.L., Royer T.V., Whiles M.R., Evans-White M., Chambers C., Griffiths N.A., Pokelsek J., Stephen M.L. (2007). Toxins in transgenic crop byproducts may affect headwater stream ecosystems. *PNAS* 104(41): 16204-16208.

- Rossi M.N., Reigada C., Godoy W.A.C. (2006). The role of habitat heterogeneity for the functional response of the spider *Nesticodes rufipes* (Araneae : Theridiidae) to houseflies. *Applied Entomology and Zoology* 41(3): 419-427.
- Rossignol N., Bonis A., Bouzib J.B. (2006). Consequence of grazing pattern and vegetation structure on the spatial variations of net N mineralisation in a wet grassland. *Applied Soil Ecology* 31(1-2): 62-72.
- Rotheray G.E. (1993). Color guide to hoverfly larvae (Diptera, Syrphidae). *Dipterist Digest* 9: 1-158.
- Rothermel R.C. (1972). A mathematical model for predicting fire spread in wildland fuels., US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station Research Paper INT-115.
- Rowe E.C., Hairiah K., Giller K.E., van Noordwijk M., Cadish G. (1999). Testing the safety-net role of hedgerow tree roots by 15N placement at different soil depths. *Agroforestry Systems* 43: 81-93.
- Roy H.E., Pell J.K. (2000). Interactions between entomopathogenic fungi and other natural enemies: Implications for biological control. *Biocontrol Science and Technology* 10(6): 737-752.
- Rudolf V.H.W., Antonovics J. (2005). Species coexistence and pathogens with frequency-dependent transmission. *American Naturalist* 166(1): 112-118.
- Rukayadi Y., Suwanto A., Tjahjono B., Harling R. (2000). Survival and epiphytic fitness of a nonpathogenic mutant of *Xanthomonas campestris* pv. *glycines*. *Applied and Environmental Microbiology* 66(3): 1183-1189.
- Russell E.P. (1989). Enemies hypothesis: a review of the effect of vegetational diversity on predatory insects and parasitoids. *Environmental Entomology* 18(4): 590-599.
- Rustamani M.A., Kanehisa K., Tsumuki H., Shiraga T. (1992). Additional observations on aphid densities and gramine contents in barley lines. *Applied Entomology and Zoology* 27: 151-153.
- Rutledge C.E., Eigenbrode S.D. (2003). Epicuticular wax on pea plants decreases instantaneous search rate of *Hippodamia convergens* larvae and reduces attachment to leaf surfaces. *Canadian Entomologist* 135(1): 93-101.
- Ryan J., Ryan M.F., McNaedie F. (1980). The effect of interrow plant cover on populations of the cabbage root fly *Deltia brassicae* (Wiedemann). *Journal of Applied Ecology* 17: 31-40.
- Ryszkowski L., Kędziora A. (2007). Modification of water flows and nitrogen fluxes by shelterbelts *Ecological Engineering* 29(4): 388-400.
- Sacco D., Zavattaro L., Grignani C. (2003). Nitrogen balance in alfalfa and orchard grass meadows. *Rivista di Agronomia* 37(1): 69-74.
- Sadeghi A.M., Starr J.L., Teasdale J.R., Rosecrance R.C., Rowland R.A. (2007). Real-time soil profile water content as influenced by weed-corn competition. *Soil science* 172(10): 759-769.
- Sadler E.J., Turner N.C. (1994). Water relationships in a sustainable agricultural system. In *Sustainable Agriculture Systems*, (Hatfield J.L., Karlen D.L., eds.), Lewis Publishers, Boca Raton: 21-45.
- Sagarra L.A., Peterkin D.D. (1999). Invasion of the Caribbean by the hibiscus mealybug, *Maconellicoccus hirsutus* green [Homoptera: Pseudococcidae]. *Phytoprotection* 80(2): 103-113.
- San Emeterio L., Arroyo A., Canals R., M. (2004). Allelopathic potential of *Lolium rigidum* Gaud. on the early growth of three associated pasture species. *Grass & Forage Science* 59: 107-112.
- San Martin G., Adriaens T., Hautier L., Ottart N. (2005). la Coccinelle asiatique - *Harmonia axyridis*. *Insectes* 139(1): 7-11.
- San Miguel Ayanz A. (1994). La dehesa española – Origen, tipología, características y gestión, Fundación Conde del Valle de Salazar ed., Escuela Técnica Superior de Ingenieros de Montes, 96 p.
- Sanchez-Moreno S., Ferris H. (2007). Suppressive service of the soil food web : Effects of environmental management. *Agriculture, Ecosystems & Environment* 119(1-2): 75-87.
- Sanderson M.A., Goslee S.C., Soder K.J., Skinner R.H., Tracy B.F., Deak A. (2007). Plant species diversity, ecosystem function, and pasture management - A perspective. *Canadian Journal of Plant Science* 87(3): 479-487.
- Sanderson M.A., Soder K.J., Muller L.D., Klement K.D., Skinner R.H., Goslee S.C. (2005). Forage mixture productivity and botanical composition in pastures grazed by dairy cattle. *Agronomy Journal* 97(5): 1465-1471.
- Sandskar B. (2003). Apple scab (*Venturia inaequalis*) and pests in organic orchards. Doctoral Thesis, Department of Crop Science, Swedish University of Agricultural Sciences, Alnarp, 39.
- Sankaran M., McNaughton S.J. (1999). Determinants of biodiversity regulate compositional stability of communities. *Nature* 401: 691-693.
- Sarfraz M., Keddie A.B., Dossdall L.M. (2005). Biological control of the diamondback moth, *Plutella xylostella*: A review. *Biocontrol Science and Technology* 15(8): 763-789.
- Sarthou J.P. (2006). *Typha latifolia* Funck, une héliophyte hôte de Pucerons de substitution pour des Syrphidés, Cécidomyidés et Coccinellidés auxiliaires des cultures. *L'Entomologiste* 62(3-4): 117-118.
- Sarthou J.P., Ouin A., Arrignon F., Barreau G., Bouyjou B. (2005). Landscape parameters explain the distribution and abundance of *Episyrphus balteatus* (Diptera: Syrphidae). *European Journal of Entomology* 102(3): 539-545.
- Sarvari M., El Hallof N. (2006). Effect of crop rotation and nutrient supply on the yield of maize. *Cereal Research Communications* 34(1): 545-548.
- Sauvant D., Grolier P., Azais-Braesco V. (2002). Nutritional significance of vitamine A in dairy products. In *Encyclopaedia of Dairy Science*, Academic Press, London: 2657-2664.
- Scarratt S.L., Wratten S.D., Lavandero B., Irvin N.A. (2004). A hierarchy of research approaches to the successful use of "resource subsidies" to improve parasitoid performance. In *California Conference on Biological Control IV*, Berkeley, California, USA, 13-15 July, 2004, (Hoddle M.S., ed.), Berkeley USA: 88-94.
- Scehovic J. (1990). Tanins et autres polymères phénoliques dans les plantes de prairies : détermination de leur teneur et de leur activité biologique. *Revue Suisse d'Agriculture* 22(3): 179-184.
- Schausberger P. (2003). Cannibalism among phytoseiid mites: a review. *Experimental & Applied Acarology* 29(3-4): 173-191.
- Scheepens P.C., Müller-Schärer H., Kempenaar C. (2001). Opportunities for biological weed control in Europe. *BioControl* 46: 127-138.
- Schellhorn N.A., Sork V.L. (1997). The impact of weed diversity on insect population dynamics and crop yield in collards, *Brassica oleracea* (Brassicaceae). *Oecologia* 111(2): 233-240.
- Scherber C., Mwangi P.N., Temperton V.M., Roscher C., Schumacher J., Schmid B., Weisser W.W. (2006). Effects of plant diversity on invertebrate herbivory in experimental grassland. *Oecologia* 147(3): 489-500.

- Scherer-Lorenzen M., Palmberg C., Prinz A., Schulze E.-D. (2003). The role of plant diversity and composition for nitrate leaching in grasslands. *Ecology* 84: 1539-1552.
- Scheu S. (2001). Plants and generalist predators as links between the below-ground and above-ground system. *Basic and Applied Ecology* 2(1): 3-13.
- Schlöpfer F., Erickson J.D. (2001). A biotic control perspective on nitrate contamination of groundwater from agricultural production. *Agricultural and Resource Economic Review* 30: 113-126.
- Schlöpfer F., Pfisterer A.B., Schmid B. (2005). Non-random species extinction and plant production: implications for ecosystem functioning. *Journal of Applied Ecology* 42: 13-24.
- Schmaedick M.A., Shelton A.M. (2000). Arthropod predators in cabbage (Cruciferae) and their potential as naturally occurring biological control agents for *Pieris rapae* (Lepidoptera : Pieridae). *Canadian Entomologist* 132(5): 655-675.
- Schmid B. (2002). The species richness-productivity controversy. *Trends in Ecology & Evolution* 17: 113-114.
- Schmid B., Hector A. (2004). The value of biodiversity experiments. *Basic and Applied Ecology* 5(6): 535-542.
- Schmid B., Joshi J., Schlöpfer F. (2002). Empirical evidence for biodiversity-ecosystem functioning relationships. In *Linking biodiversity and ecosystem functioning*, (Kinzig A. et al., eds.), Monographs in Population Ecology, 33, Princeton University Press: 120-150.
- Schmidt M.H., Roschewitz I., Thies C., Tschamtko T. (2005). Differential effects of landscape and management on diversity and density of ground-dwelling farmland spiders. *Journal of Applied Ecology* 42(2): 281-287.
- Schmidt M.H., Thewes U., Thies C., Tschamtko T. (2004). Aphid suppression by natural enemies in mulched cereals. *Entomologia Experimentalis et Applicata* 113(2): 87-93.
- Schmidt M.H., Tschamtko T. (2005). The role of perennial habitats for Central European farmland spiders. *Agriculture, Ecosystems & Environment* 105(1-2): 235-242.
- Schmidt N.P., O'Neal M.E., Singer J.W. (2007). Alfalfa living mulch advances biological control of soybean aphid. *Environmental Entomology* 36(2): 416-424.
- Schmitz M., Mwangi P.N., Scherer-Lorenzen M., Weisser W.W., Schmid B. (in review). Density and evenness effects on biodiversity-ecosystem functioning relationships.
- Schmitz O.J., Hambäck P.A., Beckerman A.P. (2000). Trophic cascades in terrestrial systems: a review of the effects of carnivore removals on plants. *American Naturalist* 155(2): 141-153.
- Schroeder D. (1983). Biological control of weeds. In *Recent Advances in Weed Research*, (Fletcher W.W., ed.), Commonwealth Agricultural Bureaux: 41-78.
- Schroeder J., Thomas S.H., Murray L.W. (2005). Impacts of crop pests on weeds and weed-crop interactions. *Weed Science* 53(6): 918-922.
- Schroth G., Krauss U., Gasparotto L., Aguilar J., Duarte A., Vohland K. (2000). Pests and diseases in agroforestry systems of the humid tropics. *Agroforestry Systems* 50: 199-241.
- Schuld M., Schmuck R. (2000). Effects of thiacloprid, a new chloronicotinyl insecticide, on the egg parasitoid *Trichogramma cacoeciae*. *Ecotoxicology* 9(3): 197-205.
- Schwartz M.W., Brigham C.A., Hoeksema J.D., Lyons K.G., Hills M.H., van Mantgem P.J. (2000). Linking biodiversity to ecosystem functioning: implications for conservation ecology. *Oecologia* 122: 297-305.
- Schweitzer J.A., Bailey J.K., Hart S.C., Whitham T.G. (2005). Nonadditive effects of mixing cottonwood genotypes on litter decomposition and nutrient dynamics. *Ecology* 86(10): 2834-2840.
- Schweitzer J.A., Bailey J.K., Hart S.C., Wimp G.M., Chapman S.K., Whitham T.G. (2005). The interaction of plant genotype and herbivory decelerate leaf litter decomposition and alter nutrient dynamics. *Oikos* 110: 133-145.
- Schweitzer J.A., Bailey J.K., Rehill B.J., Martinsen G.D., Hart S.C., Lindroth R.L., Keim P., Whitham T.G. (2004). Genetically based trait in a dominant tree affects ecosystem processes. *Ecology Letters* 7: 127-134.
- Schweitzer P. (2004). Mauvaises herbes et apiculture. *L'abeille de France* 904: 293-295.
- Sebillotte M. (1990). Système de culture, un concept opératoire pour les agronomes. In *Les systèmes de culture*, (Combe L., Picard D., eds.), Inra: 165-196.
- Sebolt D.C., Landis D.A. (2004). Arthropod predators of *Galerucella californiensis* L. (Coleoptera: Chrysomelidae): an assessment of biotic interference. *Environmental Entomology* 33(2): 356-361.
- Seeber J., Seeber G.U.H., Kossler W., Langel R., Scheu S., Meyer E. (2005). Abundance and trophic structure of macro-decomposers on alpine pastureland (Central Alps, Tyrol): effects of abandonment of pasturing. *Pedobiologia* 49(3): 221-228.
- Seelmann L., Auer A., Hoffmann D., Schausberger P. (2007). Leaf pubescence mediates intraguild predation between predatory mites. *Oikos* 116(5): 807-817.
- Selje N., Hoffmann E.M., Muetzel S., Ningrat R., Wallace R.J., Becker K. (2007). Results of a screening programme to identify plants or plant extracts that inhibit ruminal protein degradation. *British Journal of Nutrition* 98: 45-53.
- Sellenet P. (2000). L'ivraie, l'ivresse et le bon grain. *La Garance Voyageuse* 52: 52-55.
- Senapati B.K., Lavelle P., Giri S., Pashanasi B., Alegre J., Decaëns T., Jiménez J.J., Albrecht A., Blanchart E., Mahieux M., Rousseaux L., Thomas R.J., Panigrahi P.K., Venkatachalam M. (1999). Soil earthworm technologies for tropical agroecosystems. In *Earthworm management in tropical agroecosystems*, (Lavelle P. et al., eds.), CAB International, Oxford: 199-237.
- Senmartin M., Aguiar M.R., Distel R., Moretto A.S., Ghersa C.M. (2004). Litter quality and nutrient cycling affected by grazing-induced species replacements along a precipitation gradient. *Oikos* 107: 148-160.
- Setälä H., Iiäksö J., Mikola J., Huhta V. (1998). Functional diversity of decomposer organisms in relation to primary production. *Applied Soil Ecology* 9(1-3): 25-31.
- Setälä H., McLean M.A. (2004). Decomposition rate of organic substrates in relation to the species diversity of soil saprophytic fungi. *Oecologia* 139(1): 98-107.
- Shaik S.A., Terrill T.H., Miller J.E., Kouakou B., Kannan G., Kaplan R.M., Burke J.M., Mosjidis J.A. (2006). Sericea lespedeza hay as a natural deworming agent against gastrointestinal nematode infection in goats. *Veterinary Parasitology* 139(1-3): 150-157.
- Shankaraiah C. (2007). Nitrogen management through biological process on nitrogen use efficiency in sugarcane and environmental protection. *Sugar Tech* 9(2/3): 132-136.

- Sharma S., Szele Z., Schilling R., Munch J.C., Schloter M. (2006). Influence of freeze-thaw stress on the structure and function of microbial communities and denitrifying populations in soil. *Applied and Environmental Microbiology* 72(3): 2148-2154.
- Shaw C., Pawluk S. (1986). The development of soil structure by *Ocotlasion tyraeum*, *Aporrectodea turgida* and *Lumbricus terrestris* and its relation to carbon budgets of three artificial soils. *Pedobiologia* 29: 327-339.
- Shea K., Chesson P.L. (2002). Community ecology theory as a framework for biological invasions. *Trends in Ecology and Evolution* 17(4): 170-176.
- Sheath G.W., Coulon J.B., Young O.A. (2001). Grassland management and animal product quality, *Grassland Ecosystems: an outlook into the 21st century*. 19. *International Grassland Congress*, Sao Paulo, Brazil, Gomide J.A. et al., eds.: 1019-1026.
- Sheehan C., Kirwan L., Connolly J., Bolger T. (2008). The effects of earthworm functional diversity on microbial biomass and the microbial community level physiological profile of soils. *European Journal of Soil Biology* 44(1): 65-70.
- Shelton M.D., Edwards C.R. (1983). Effects of weeds on the diversity and abundance of insects in soybeans. *Environmental Entomology* 12: 296-298.
- Shiel C.B., McAney C.M., Fairley J.S. (1991). Analysis of the diet of Natterer's bat *Myotis nattereri* and the common long-eared bat *Plecotus auritus* in the West of Ireland. *Journal of Zoology* 223: 299-305.
- Shiskoff N., Bruckart W.L. (1993). Evaluation of infection of target and non-target hosts by isolates of the potential biocontrol agent *Puccinia jaceae* that infect *Centaurea* spp. *Phytopathology* 83: 894-898.
- Shrewsbury P.M., Raupp M.J. (2006). Do top-down or bottom-up forces determine *Stephanitis pyrioides* abundance in urban landscapes? *Ecological Applications* 16(1): 262-272.
- Silvy C. (1999). Quantifications... le phytosanitaire III. *Dossier de l'environnement* 19: 201-212.
- Simon S. (1999). Incidence de l'environnement végétal sur les populations d'arthropodes du verger de poiriers. Thèse de doctorat, Biologie des populations et écologie, Université de Montpellier 2, Montpellier, 438 p.
- Six J., Elliot E.T., Paustian K. (2000). Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology & Biochemistry* 32: 2099-2103.
- Six J., Frey S.D., Thiet R.K., Batten K.M. (2006). Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Science Society of America Journal* 70(2): 555-569.
- Smart S.M., Bunce R.G.H., Marrs R.H., LeDuc M., Firbank L.G., Maskell L.C., Scott W.A., Thompson K., Walker K.J. (2005). Large-scale changes in the abundance of common higher plant species across Britain between 1978, 1990 and 1998 as a consequence of human activity: Tests of hypothesised changes in trait representation. *Biological Conservation* 124: 355-371.
- Smeding F.W., Joenje W. (1999). Farm-Nature Plan: landscape ecology based farm planning. *Landscape and Urban Planning* 46(1): 109-115.
- Smith J.G. (1976). Influence of crop background on aphids and other phytophagous insects on Brussels sprouts. *Annals of Applied Biology* 83: 1-13.
- Smith J.G. (1976). Influence of crop background on natural enemies of aphids on Brussels sprouts. *Annals of Applied Biology* 83: 15-29.
- Smith M.D., Hartnett D.C., Wilson G.W.T. (1999). Interacting influence of mycorrhizal symbiosis and competition on plant diversity in tallgrass prairie. *Oecologia* 121: 574-582.
- Smith M.D., Knapp A.K. (2003). Dominant species maintain ecosystem function with non-random species loss. *Ecology Letters* 6: 509-517.
- Smith P. (2008). Land use change and soil organic carbon dynamics. *Nutrient Cycling in Agroecosystems* 81(2): 1573-0867.
- Smith R.G., Gross K.L. (2006). Weed community and corn yield variability in diverse management systems. *Weed Science* 54: 106-113.
- Smith R.G., Gross K.L., Robertson G.P. (2008). Effects of Crop Diversity on Agroecosystem Function: Crop Yield Response. *Ecosystems* 11: 355-366.
- Smith S.E., Read D.J. (1997). *Mycorrhizal Symbiosis*, 2nd ed., Academic Press, London, 605 p.
- Snyder W.E., Clevenger G.M., Eigenbrode S.D. (2004). Intraguild predation and successful invasion by introduced ladybird beetles. *Oecologia* 140(4): 559-565.
- Snyder W.E., Wise D.H. (1999). Predator interference and the establishment of generalist predator populations for biocontrol. *Biological Control* 15(3): 283-292.
- Soder K.J., Rook A.J., Sanderson M.A., Goslee S.C. (2007). Interaction of plant species diversity on grazing behavior and performance of livestock grazing temperate region pastures. *Crop Science* 47(1): 416-425.
- Soder K.J., Sanderson M.A., Stack J.L., Muller L.D. (2006). Intake and performance of lactating cows grazing diverse forage mixtures. *Journal of Dairy Science* 89(6): 2158-2167.
- Sokol-Hessner L., Schmitz O.J. (2002). Aggregate Effects of Multiple Predator Species on a Shared Prey. *Ecology* 83(9): 2367-2372.
- Soliva R., Ronningen K., Bella I., Bezak P., Cooper T., Flo B.E., Marty P., Potter C. (2008). Envisioning upland futures: Stakeholder responses to scenarios for Europe's mountain landscapes. *Journal of Rural Studies* 24(1): 56-71.
- Soltner D. (1995). L'arbre et la haie. Pour la production agricole, pour l'équilibre écologique et le cadre de vie rurale, Collection Sciences et Techniques Agricoles, 208 p.
- Soon Y.K., Arshad M.A. (2004). Tillage, crop residue and crop sequence effects on nitrogen availability in a legume-based cropping system. *Canadian Journal of Soil Science* 84(4): 421-430
- Souchère V., Cerdan O., Dubreuil N., Le Bissonnais Y., King C. (2005). Title: Modelling the impact of agri-environmental scenarios on runoff in a cultivated catchment (Normandy, France). *Catena* 61: 229-240.
- Souchère V., King C., Dubreuil N., Lecomte-Morel V., Le Bissonnais Y., Chalal M. (2003). Grassland and crop trends: role of the European Union Common Agricultural Policy and consequences for runoff and soil erosion. *Environmental Science & Policy* 6(1): 7-16.
- Soussana J.F., Lüscher A. (2007). Temperate grasslands and global atmospheric change: a review. *Grass & Forage Science* 62: 127-134.
- Soussana J.-F., Teyssonere F., Picon-Cochard C., Dawson T.E. (2005). A trade-off between nitrogen uptake and use increases responsiveness to elevated CO₂ in infrequently cut mixed C3 grasses. *New Phytologist* 166: 217-230.
- Spahillari M., Hammer K., Gladis T., Diederichsen A. (1999). Weeds as part of agrobiodiversity. *Outlook on Agriculture* 28(4): 227-232.
- Spehn E.M., Hector A., Joshi J., Scherer-Lorenzen M., Schmid B., Bazeley-White E., Beierkuhnlein C., Caldeira M.C., Diemer M., Dimitrakopoulos P.G., Finn J.A., Freitas H., Giller P.S., Good J., Harris R., Höglberg P., Huss-Danell K., Jumpponen A., Koricheva J., Leadley P.W., Loreau M., Minns A., Mulder C.P.H., O'Donovan G., Otway S.J., Palmberg C., Pereira J.S., Pfisterer A.B., Prinz

- A., Read D.J., Schulze E.-D., Siamantziouras A.-S.D., Terry A.C., Troumbis A.Y., Woodward F.I., Yachi S., Lawton J.H. (2005). Ecosystem effects of biodiversity manipulations in European grasslands. *Ecological Monographs* 75: 37-63.
- Spehn E.M., Scherer-Lorenzen M., Schmid B., Hector A., Caldeira M.C., Dimitrakopoulos P.G., Finn J.A., Jumpponen A., O'Donovan G., Pereira J.S., Schulze E.-D., Troumbis A.Y., Körner C. (2002). The role of legumes as a component of biodiversity in a cross-European study of grassland biomass nitrogen. *Oikos* 98: 205-218.
- Speight M.C.D., Castella E., Sarthou J.-P., Monteil C. (2006). Syrph the Net on CD. In The database of European Syrphidae, Syrph the Net Publications, Dublin: Issue 5.
- Sprent J.I., t'Manneetje L. (1996). The role of legumes in sustainable farming systems: past, present and future, *Legumes in sustainable farming systems*, Occasional symposium No.30, Younie D., ed., British Grassland Society: 2-14.
- Spring J.L., Delabays N. (2006). Research experiment on weakly competitive cover cropping in vineyard: agronomical aspects. *Revue Suisse de Viticulture, Arboriculture et Horticulture* 38(6): 355-359.
- Stahl W., Sies H. (2005). Bioactivity and protective effects of natural carotenoids. *Biochimica et biophysica acta. Molecular basis of disease* 1740(2): 101-107.
- Stamps W.T., Linit M.J. (1997). Plant diversity and arthropod communities: Implications for temperate agroforestry. *Agroforestry Systems* 39(1): 73-89.
- Stark J.D., Banks J.E., Acheampong S. (2004). Estimating susceptibility of biological control agents to pesticides: influence of life history strategies and population structure. *Biological Control* 29(3): 392-398.
- Steffan-Dewenter I. (2003). Seed set of male-sterile and male-fertile oilseed rape (*Brassica napus*) in relation to pollinator density. *Apidologie* 34: 227-235.
- Steffan-Dewenter I., Munzenberg U., Burger C., Thies C., Tschamtker T. (2002). Scale-dependent effects of landscape context on three pollinator guilds. *Ecology* 83(5): 1421-1432.
- Steffan-Dewenter I., Tschamtker T. (1999). Effects of habitat isolation on pollinator communities and seed set. *Oecologia* 121(3): 432-440.
- Steffan-Dewenter I., Tschamtker T. (2000). Butterfly community structure in fragmented grasslands. *Ecology Letters* 3: 449-456.
- Steinberg C., Edel-Hermann V., Alabouvette C., Lemanceau P. (2007). Soil suppressiveness to plant diseases. In *Modern Soil Microbiology*, (van Elsas J.D. et al., eds.), CRC Press: 455-478.
- Stephan A., Meyer A.H., Schmid B. (2000). Plant diversity affects culturable soil bacteria in experimental grassland communities. *Journal of Ecology* 88: 988-998.
- Stephens M.J., France C.M., Wratten S.D., Frampton C. (1998). Enhancing biological control of leafrollers (Lepidoptera : Tortricidae) by sowing buckwheat (*Fagopyrum esculentum*) in an orchard. *Biocontrol Science and Technology* 8(4): 547-558.
- Stephens P.M., Davoren C.W. (1997). Influence of the earthworms *Aporrectodea trapezoides* and *A. rosea* on the disease severity of *Rhizoctonia solani* on subterranean clover and ryegrass. *Soil Biology & Biochemistry* 29(3-4): 511-516.
- Stevens M.H.H., Carson W.P. (2001). Phenological complementarity, species diversity, and ecosystem function. *Oikos* 92(2): 291-296.
- Stock W.D., Wienand K.T., Baker A.C. (1995). Impacts of invading N₂-fixing *Acacia* species on patterns of nutrient cycling in two Cape ecosystems: evidence from soil incubation studies and 15N natural abundance values. *Oecologia* 101: 375-382.
- Stockdale E.A., Lampkin N.H., Hovi M., Keatinge R., Lennartsson E.K.M., Macdonald D.W., Padel S., Tattersall F.H., Wolfe M.S., Watson C.A. (2001). Agronomic and environmental implications of organic farming systems. *Advances in Agronomy* 70: 261-325.
- Stocker R., Körner C.H., Schmid B., Niklaus P.A., Leadley P.W. (1999). A field study of the effects of elevated CO₂ and plant species diversity on ecosystem-level gas exchange in a planted calcareous grassland. *Global Change Biology* 5(1): 95-105.
- Stohlgren T.J., Binkley D., Chong G.W., Kalkhan M.A., Schell L.D., Bull K.A., Otsuki Y., Newman G., Bashkin M., Son Y. (1999). Exotic plant species invade hot spots of native plant diversity. *Ecological Monographs* 69: 25-46.
- Stohlgren T.J., Jarnevitch C., Chong G.W., Evangelista P.H. (2006). Scale and plant invasions: a theory of biotic acceptance. *Preslia* 78: 405-426.
- Stokstad E. (2007). The case of the empty hives. *Science* 316(5827): 970-972.
- Stolcova J. (2007). Insect injury and mortality of weed seedlings on early abandoned field, *14th EWRS Symposium*, Hamar (Norway), 18-21 June 2007: 233.
- Stone G.N., Sunnucks P. (1993). Genetic consequences of an invasion through a patchy environment - the cynipid gallwasp *Andricus quercuscalicis* (Hymenoptera: Cynipidae). *Molecular Ecology* 2(4): 251-268.
- Stopes C., Millington S., Woodward L. (1996). Dry matter and nitrogen accumulation by three leguminous green manure species and the yield of a following wheat crop in an organic production system. *Agriculture, Ecosystems & Environment* 57(2-3): 189-196.
- Storkey J. (2006). A functional group approach to the management of UK arable weeds to support biological diversity. *Weed Research* 46: 513-522.
- Storkey J., Westbury D.B. (2007). Managing arable weeds for biodiversity. *Pest Management Science* 63(6): 517-523.
- Stout J.C. (2007). Reproductive biology of the invasive exotic shrub, *Rhododendron ponticum* L. (Ericaceae). *Botanical Journal of the Linnean Society* 155(3): 373-381.
- Straub C.S., Snyder W.E. (2006). Species identity dominates the relationship between predator biodiversity and herbivore suppression. *Ecology* 87(2): 277-282.
- Stromberg K.D., Kinkel L.L., Leonard K.J. (2004). Quantifying the effect of bacterial antagonists on the relationship between phyllosphere population sizes of *Xanthomonas translucens* pv. *translucens* and subsequent bacterial leaf streak severity on wheat seedlings. *Biological Control* 29(1): 58-65.
- Sturz A.V., Christie B.R. (2003). Beneficial microbial allelopathies in the root zone: the management of soil quality and plant disease with rhizobacteria. *Soil & Tillage Research* 72(2): 107-123.
- Sturz A.V., Ryan D.A.J., Coffin A.D., Matheson B.G., Arsenault W.J., Kimpinski J., Christie B.R. (2004). Stimulating disease suppression in soils: sulphate fertilizers can increase biodiversity and antibiosis ability of root zone bacteria against *Streptomyces scabies*. *Soil Biology & Biochemistry* 36(2): 343-352.
- Suding K.N., Collins S.L., Gough L., Clark C., Cleland E.E., Gross K.L., Milchunas D.G., Pennings S. (2005). Functional- and abundance-based mechanisms explain diversity loss due to N fertilization. *Proceedings of the National Academy of Sciences (U.S.A.)* 102: 4387-4392.

- Sung D.G., Lim S.H., Ko J.W., Cho G.S. (2001). Scenic evaluation of landscapes for urban design purposes using GIS and ANN. *Landscape and Urban Planning* 56: 75-85.
- Sutherland J.P., Sullivan M.S., Poppy G.M. (2001). Distribution and abundance of aphidophagous hoverflies (Diptera: Syrphidae) in wildflower patches and field margin habitats. *Agricultural and Forest Entomology* 3(1): 57-64.
- Swift M.J., Andren O., Brussaard L., Briones M., Couteaux M.M., Ekschmitt K., Kjoller A., Loiseau P., Smith P. (1998). Global change, soil biodiversity, and nitrogen cycling in terrestrial ecosystems: three case studies. *Global Change Biology* 4(7): 729-743.
- Swift M.J., Izac A.M.N., van Noordwijk M. (2004). Biodiversity and ecosystem services in agricultural landscapes - are we asking the right questions? *Agriculture, Ecosystems & Environment* 104(1): 113-134.
- Symondson W.O.C., Cesarini S., Dodd P.W., Harper G.L., Bruford M.W., Glen D.M., Wiltshire C.W., Harwood J.D. (2006). Biodiversity vs. biocontrol: positive and negative effects of alternative prey on control of slugs by carabid beetles. *Bulletin of Entomological Research* 96(6): 637-645.
- Symondson W.O.C., Glen D.M., Ives A.R., Langdon C.J., Wiltshire C.W. (2002). Dynamics of the relationship between a generalist predator and slugs over five years. *Ecology* 83(1): 137-147.
- Symstad A.J. (2000). A test of the effects of functional group richness and composition on grassland invasibility. *Ecology* 81(1): 99-109.
- Symstad A.J., Chapin III F.S., Wall D.H., Gross K.L., Huenneke L.F., Mittelbach G.G., Peters D.P.C., Tilman D. (2003). Long-term and large-scale perspectives on the relationship between biodiversity and ecosystem functioning. *BioScience* 53: 89-98.
- Symstad A.J., Tilman D. (2001). Diversity loss, recruitment limitation, and ecosystem functioning: lessons learned from a removal experiment. *Oikos* 92: 424-435.
- Symstad A.J., Tilman D., Willson D., Knops J.M.H. (1998). Species loss and ecosystem functioning : effects of species identity and species composition. *Oikos* 81: 389-397.
- Szente K., Tuba Z., Nagy Z., Csintalan Z. (1993). Competition between *Chenopodium album* and *Helianthus annuus* as reflected in photosynthesis and transpiration. *Photosynthetica* 28(3): 465-472.
- Taghavi S., Barac T., Greenberg B., Borremans B., Vangronsveld J., van der Lelie D. (2005). Horizontal gene transfer to endogenous endophytic bacteria from poplar improves phytoremediation of toluene. *Applied and Environmental Microbiology* 71(12): 8500-8505.
- Tahvanainen J.O., Root R.B. (1972). The influence of vegetational diversity on the population ecology of a specialized herbivore, *Phyllotreta cruciferae* (Coleoptera: Chrysomelidae). *Oecologia* 10: 321-346.
- Tahvanainen L., Tyrväinen L., Ihalainen M., Vuorela N., Kolehmainen O. (2001). Forest management and public perceptions – visual versus verbal information. *Landscape and Urban Planning* 53: 53-70.
- Takayanagi H., Yamamoto Y., Takeda N. (1986). An ovary-stimulating factor in the shrimp *Paratyia compressa*. *Journal of experimental zoology* 240(2): 203-209.
- Tamene L., Vlek P.L.G. (2007). Assessing the potential of changing land use for reducing soil erosion and sediment yield of catchments: a case study in the highlands of northern Ethiopia. *Soil Use and Management* 23(1): 82-91.
- Tanaka S., Nishida T., Ohsaki N. (2007). Sequential rapid adaptation of indigenous parasitoid wasps to the invasive butterfly *Pieris brassicae*. *Evolution* 61(8): 1791-1802.
- Tasser E., Mader M., Tappeiner U. (2003). Effects of land use in alpine grasslands on the probability of landslides. *Basic and Applied Ecology* 4: 271-280.
- Tasser E., Tappeiner U. (2005). New model to predict rooting in diverse plant community compositions. *Ecological Modelling* 185(2-4): 195-211.
- Temperton V.M., Mwangi P.N., Scherer-Lorenzen M., Schmid B., Buchmann N. (2007). Positive interactions between nitrogen-fixing legumes and four different neighbouring species in a biodiversity experiment. *Oecologia* 151: 190-205.
- Theunissen J., Booij C.J.H., Lotz L.A.P. (1995). Effects of intercropping white cabbage with clovers on pest infestation and yield. *Entomologia Experimentalis et Applicata* 74(1): 7-16.
- Thibaudon M., Elias K., Besancenot J.P. (2004). Ambroisie et allergie : Le cas de la France *Environnement risques & santé* 3(6): 353-367.
- Thiébaud F., Cozic P., Véron F., Brau-Nogué C., Bornard A. (2001). Intérêts et limites des différents couverts fourragers et pratiques associées vis-à-vis de l'environnement. Analyse bibliographique. *Fourrages* 168: 449-475.
- Thiele J., Otte A. (2007). Impact of *Heracleum mantegazzianum* on invaded vegetation and human activities. In *Ecology and management of Giant Hogweed (Heracleum mantegazzianum)*, (Pysek P. et al., eds.), CABI, Wallingford: 144-156.
- Thies C., Roschewitz I., Tschamtké T. (2005). The landscape context of cereal aphid-parasitoid interactions. *Proceedings of the Royal Society B-Biological Sciences* 272(1559): 203-210.
- Thies C., Steffan Dewenter I., Tschamtké T. (2003). Effects of landscape context on herbivory and parasitism at different spatial scales. *Oikos* 101(1): 18-25.
- Thies C., Tschamtké T. (1999). Landscape structure and biological control in agroecosystems. *Science* 285: 893-895.
- Thiet R.K., Frey S.D., Six D.L. (2006). Do growth yield efficiencies differ between soil microbial communities differing in fungal:bacterial ratios? Reality check and methodological issues. *Soil Biology & Biochemistry* 38(4): 837-844.
- Thirup L., Johnsen K., Torsvik V., Spliid N.H., Jacobsen C.S. (2001). Effects of fenpropimorph on bacteria and fungi during decomposition of barley roots. *Soil Biology & Biochemistry* 33(11): 1517-1524.
- Thomas M.B., Wratten S.D., Sotherton N.W. (1992). Creation of 'island' habitats in farmland to manipulate populations of beneficial arthropods: predator densities and species composition. *Journal of Applied Ecology* 29(2): 524-531.
- Thomas S., H, Schroeder, J, Murray, L, W, (2005). The role of weeds in nematode management. *Weed Science* 53: 923-928.
- Thomas S.R., Goulson D., Holland J.M. (2000). Spatial and temporal distributions of predatory Carabidae in a winter wheat field. *Aspects of Applied Biology* 62: 55-60.
- Thomas S.R., Goulson D., Holland J.M. (2001). Resource provision for farmland gamebirds: the value of beetle banks. *Annals of Applied Biology* 139(1): 111-118.
- Thomas S.R., Noordhuis R., Holland J.M., Goulson D. (2002). Botanical diversity of beetle banks: Effects of age and comparison with conventional arable field margins in southern UK. *Agriculture, Ecosystems & Environment* 93(1-3): 403-412.

- Thompson I.P., Bailey M.J., Fenlon J.S., Fermor T.R., Lilley A.K., Lynch J.M., McCormack P.J., McQuilken M.P., Purdy K.J., Rainey P.B., Whipps J.M. (1993). Quantitative and qualitative seasonal changes in the microbial community from the phyllosphere of sugar beet (*Beta vulgaris*). *Plant and Soil* 150: 177-191.
- Thompson K., Askew A.P., Grime J.P., Dunnett N.P., Willis A.J. (2005). Biodiversity, ecosystem function and plant traits in mature and immature plant communities. *Functional Ecology* 19: 355-358.
- Thompson K., Jones A. (1999). Human population density and prediction of local plant extinction in Britain. *Conservation Biology* 13: 185-189.
- Thomson J.D. (2001). Using pollination deficits to infer pollinator declines: Can theory guide us? *Conservation Ecology* 5(1): U199-U209.
- Thomson L.J., Hoffmann A.A. (2006). Field validation of laboratory-derived IOBC toxicity ratings for natural enemies in commercial vineyards. *Biological Control* 39(3): 507-515.
- Tichit M., Durant D., Kernéis E. (2005). The role of grazing in creating suitable sward structures for breeding waders in agricultural landscapes. *Livestock Production Science* 96(1): 119-128.
- Tichit M., Havet A., Renault O., Potter T. (2008). Gérer l'hétérogénéité des prairies à différentes échelles : une clé pour la conception d'un système d'élevage performant sur le plan environnemental *In* Les exploitations d'élevage en mouvement. Flexibilités et dynamiques face aux aléas et aux nouveaux enjeux des filières et territoires (Dedieu B., Chia, E., Moulin, C.H., Leclerc, B., Tichit, M., ed.), QUAE, Paris: 253-264.
- Tietjen W.J., Cady A.B. (2007). Sublethal exposure to a neurotoxic pesticide affects activity rhythms and patterns of four spider species. *Journal of Arachnology* 35(2): 396-406.
- Tietjen W.J., Cady A.B. (2007). Sublethal insecticide exposure influences behaviors of web-building spiders, *Oral presentation at the American Arachnology Society 31st annual meeting, July 2007*, Susquehanna University (Pennsylvania).
- Tilman D., Downing J.A. (1994). Biodiversity and stability in grasslands. *Nature* 367: 363-365.
- Tilman D., Knops J. (1996). Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* 379(6567): 718-720.
- Tilman D., Lehman C.L., Bristow C.E. (1998). Diversity - stability relationships: statistical inevitability or ecological consequence? *American Naturalist* 151: 277-282.
- Tilman D., Lehman C.L., Thomson K.T. (1997). Plant diversity and ecosystem productivity : Theoretical considerations. *Proceedings of the National Academy of Science of the United States of America* 94: 1857-1861.
- Tilman D., May R.M., Lehman C.L., Nowak M.A. (1994). Habitat destruction and the extinction debt. *Nature* 371: 65-66.
- Tilman D., Reich P.B., Knops J.M.H. (2006). Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* 441: 629-632.
- Timms-Wilson T.M., Kilshaw K., Bailey M.J. (2004). Risk assessment for engineered bacteria used in biocontrol of fungal disease in agricultural crops. *Plant and Soil* 266(1-2): 57-67.
- Tisdall J.M., Odes J.M. (1982). Organic matter and water-stable aggregates in soils. *Journal of Soil Sciences* 33: 141-163.
- Tiunov A.V., Scheu S. (2005). Facilitative interactions rather than resource partitioning drive diversity-functioning relationships in laboratory fungal communities. *Ecology Letters* 8(6): 618-625.
- Toljander J.F., Artursson V., Paul L.R., Jansson J.K., Finlay R.D. (2005). Attachment of different soil bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal vitality and fungal species. *FEMS Microbiology Letters* 254(1): 34-40.
- Toljander J.F., Lindahl B.D., Paul L.R., Elfstrand M., Finlay R. (2007). Influence of arbuscular mycorrhizal mycelial exudates on soil bacterial growth and community structure. *FEMS Microbiology Ecology* 61: 295-304.
- Tonhasca A. (1993). Effects of agroecosystem diversification on natural enemies of soybean herbivores. *Entomologia Experimentalis et Applicata* 69(1): 83-90.
- Tonhasca A., Byrne D.N. (1994). The effects of crop diversification on herbivorous insects: a meta-analysis approach. *Ecological Entomology* 19(3): 239-244.
- Tomambe G., Cornu A., Verdier-Metz I., Pradel P., Kondjoyan N., Figueredo G., Hulin S., Bonanno A., Martin B. (2007). Effet de l'addition dans le lait d'une huile essentielle de plantes de prairie naturelle sur les caractéristiques sensorielles de fromage de type Cantal, 14. *Rencontres Recherches Ruminants*: 100.
- Tomambe G., Ferlay A., Farruggia A., Chilliard Y., Loiseau P., Garel J.P., Martin B. (2007). Effet de la diversité floristique des pâturages de montagne sur le profil en acides gras et les caractéristiques sensorielles des laits, 14. *Rencontres Recherches Ruminants*.
- Trana A., Fedele V., Cifuni G.F., Impemba G., Claps S., Rubino R. (2005). Relationships among diet botanical composition, milk fatty acid and herbage fatty acid content in grazing goats. *Options Méditerranéennes, Série A, Séminaires méditerranéens* 67: 269-273.
- Traveset A., Richardson D.M. (2006). Biological invasions as disruptors of plant reproductive mutualisms. *Trends in Ecology & Evolution* 21(4): 208-216.
- Tremblay A., Mineau P., Stewart R.K. (2001). Effects of bird predation on some pest insect populations in corn. *Agriculture, Ecosystems and Environment* 83(1): 143-152.
- Tscharntke T. (2000). Parasitoid populations in the agricultural landscape. *In* Parasitoid Population Biology, (Hochberg M.E., Ives A.R., eds.), Princeton University Press, Princeton: 235-253.
- Tscharntke T., Bommarco R., Clough Y., Crist T.O., Kleijn D., Rand T.A., Tylianakis J.M., van Nouhuys S., Vidal S. (2007). Conservation biological control and enemy diversity on a landscape scale. *Biological Control* 43(3): 294-309.
- Tscharntke T., Rand T.A., Bianchi F. (2005). The landscape context of trophic interactions: insect spillover across the crop-noncrop interface. *Annales Zoologici Fennici* 42(4): 421-432.
- Tscharntke T., Steffan Dewenter I., Kruess A., Thies C. (2002). Contribution of small habitat fragments to conservation of insect communities of grassland-cropland landscapes. *Ecological Applications* 12(2): 354-363.
- Tsitsilas A., Stuckey S., Hoffmann A.A., Weeks A.R., Thomson L.J. (2006). Shelterbelts in agricultural landscapes suppress invertebrate pests. *Australian Journal of Experimental Agriculture* 46(10): 1379-1388.
- Turner M.G., Gardner R.H., Dale V.H., O'Neill R.V. (1989). Predicting the spread of disturbance across heterogeneous landscapes. *Oikos* 55(1): 121-129.
- Turner M.G., Gardner R.H., O'Neill R.V. (2001). *Landscape Ecology in theory and practice: Pattern and Process*, Springer-Verlag, New York, 424 p.

- Umoru P.A., Powell W. (2002). Sub-lethal effects of the insecticides pirimicarb and dimethoate on the aphid parasitoid *Diaeretiella rapae* (Hymenoptera : Braconidae) when attacking and developing in insecticide-resistant hosts. *Biocontrol Science and Technology* 12(5): 605-614.
- Unsicker S.B., Baer N., Kahmen A., Wagner M., Buchmann N., Weisser W.W. (2006). Invertebrate herbivory along a gradient of plant species diversity in extensively managed grasslands. *Oecologia* 150(2): 233-246.
- Urano S., Abe J., Matsumura M., Nagasaka K., Takabayasi J. (2006). The number of *Cotesia plutellae* needed to suppress the population growth of diamondback moth, *Plutella xylostella*, in a greenhouse. *Bulletin OILB/SROP* 29(4): 187-190.
- Urbach G. (1990). Effect of feed on flavor in dairy foods. *Journal of Dairy Science* 73: 3639-3650.
- Urbaneja A., Garcia Mari F., Tortosa D., Navarro C., Vanaclocha P., Bagues L., Castanera P. (2006). Influence of ground predators on the survival of the Mediterranean fruit fly pupae, *Ceratitis capitata*, in Spanish citrus orchards. *BioControl* 51(5): 611-626.
- Vaast P., Van Kanten R., Siles P., Dzib B., Franck N., Harmand J.M., Génard M. (2004). Shade: A key factor for coffee sustainability and quality, *20th International Conference on Coffee Science, ASIC*, Bangalore, India: 887-896.
- Vallis I., Haydock K.P., Ross P.J., Henzell E.F. (1967). Isotopic studies on the uptake of nitrogen by pasture plants. III. The uptake of small additions of ¹⁵N-labelled fertilizer by Rhodes grass and Townsville lucerne. *Australian Journal of Agricultural Research* 18: 865-877.
- Valone T.J., Hoffman C.D. (2003). A mechanistic examination of diversity-stability relationships in annual plant communities. *Oikos* 103(3): 518-527.
- Vamosi J.C., Knight K.M., Steets J.A., Mazer S.J., Burd M., Ashman T.I. (2006). Pollination decays in biodiversity hotspots. *PNAS* 103: 956-961.
- van Bruggen A.H.C., Semenov A.M., van Diepeningen A.D., de Vos O.J., Blok W.J. (2006). Relation between soil health, wave-like fluctuations in microbial populations, and soil-borne plant disease management. *European Journal of Plant Pathology* 115(1): 105-122.
- van der Heijden M.G.A., Bardgett R.D., van Straalen N.M. (2008). The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* 11: 296-310.
- van der Heijden M.G.A., Boller T., Wiemken A., Sanders I.R. (1998). Different arbuscular mycorrhizal fungal species are potential determinants of plant community structure. *Ecology* 79(6): 2082-2091.
- van der Heijden M.G.A., Klironomos J.N., Ursic M., Moutoglou P., Streitwolf-Engel R., Boller T., Wiemken A., Sanders I.R. (1998). Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396(6706): 69-72.
- van der Heijden M.G.A., Scheublin T.R., Brader A. (2004). Taxonomic and functional diversity in arbuscular mycorrhizal fungi - is there any relationship? *New Phytologist* 164(2): 201-204.
- van der Putten W.H., Klironomos J.N., Wardle D.A. (2007). Microbial ecology of biological invasions. *ISME Journal* 1(1): 28-37.
- van der Putten W.H., Mortimer S.R., Hedlund K., Van Dijk C., Brown V.K., Leps J., Rodriguez-Barrueco C., Roy J., FDiaz Len T.A., Gormsen D., Korthals G.W., Lavorel S., Santa Regina I., Smilauer P. (2000). Plant species diversity as a driver of early succession in abandoned fields: a multi-site approach. *Oecologia* 124(1): 91-99.
- van Elsas J.D. (1997). *Modern Soil Microbiology* CRC Press, 708 p.
- van Elsas J.D., Costa R. (2007). Molecular assessment of soil microbial communities with potential for plant disease suppression. *In* Biotechnology and plant disease management, (Punja Z.K. *et al.*, eds.), CAB International, Wallingford UK: 498-517.
- van Elsas J.D., Garbeva P., Salles J. (2002). Effects of agronomical measures on the microbial diversity of soils as related to the suppression of soil-borne plant pathogens. *Biodegradation* 13(1): 29-40.
- van Elsen T. (2000). Species diversity as a task for organic agriculture in Europe. *Agriculture, Ecosystems & Environment* 77(1/2): 101-109.
- Van Emden H.F., Williams G.F. (1974). Insect stability and diversity in agro-ecosystems. *Annual Review of Entomology* 19: 455-475.
- Van Peer L., Nijs I., Reheul D., De Cauwer B. (2004). Species richness and susceptibility to heat and drought extremes in synthesized grassland ecosystems: compositional vs physiological effects. *Functional ecology* 18: 769-778.
- van Ruijven J., Berendse F. (2003). Positive effects of plant species diversity on productivity in the absence of legumes. *Ecology Letters* 6: 170-175.
- Van Ruijven J., Berendse F. (2007). Contrasting effects of diversity on the temporal stability of plant populations. *Oikos* 116: 1323-1330.
- van Wijnen H.J., van der Wal R., Bakker J.P. (1999). The impact of herbivores on nitrogen mineralization rate: consequences for salt-marsh succession. *Oecologia* 118(2): 225-231.
- Vanderhoeven S., Dassonville N., Chapuis-Lardy L., Hayez M., Meerts P. (2006). Impact of the invasive alien plant *Solidago gigantea* on primary productivity, plant nutrient content and soil mineral nutrient concentrations *Plant and Soil* 286(1-2): 259-268.
- Vandermeer J. (1995). The ecological basis of alternative agriculture. *Annual Review of Ecology and Systematics* 26: 201-224.
- Vandermeer J., Perfecto I. (1995). Breakfast of biodiversity: the truth about rain forest destruction, Food First Books, Oakland, 185 p.
- Vandermeer J., Perfecto Y., Philpott S.M. (2007). Clusters of ant colonies and robust criticality in a tropical agroecosystem. *Nature* 451: 457-459.
- Vandermeer J., Van Noordwijk M., Anderson J., Ong C., Perfecto I. (1998). Global change and multi-species agroecosystems: Concepts and issues *Agriculture, Ecosystems and Environment* 67: 1-22.
- Vandermeer J.H. (1989). *The Ecology of Intercropping*, Cambridge University Press, Cambridge, UK, 237 p.
- Vanni M.J., Findlay D.L. (1990). Trophic cascades and phytoplankton community structure. *Ecology* 71: 921-937.
- Varchola J.M., Dunn J.P. (2001). Influence of hedgerow and grassy field borders on ground beetle (Coleoptera : Carabidae) activity in fields of corn. *Agriculture, Ecosystems & Environment* 83(1-2): 153-163.
- Vavilov N.I. (1951). Origin, variation, immunity and breeding of cultivated plants: selected writings, *Chronica botanica*, Waltham, Massachusetts.
- Vazquez A., Pérez B., Fernandez-Gonzalez F., Moreno J.M. (2002). Recent fire regime characteristics and potential natural vegetation relationships in Spain. *Journal of Vegetation Science* 13: 663-676.
- Vellend M. (2005). Species diversity and genetic diversity: parallel processes and correlated patterns. *American Naturalist* 166(2): 199-215.

- Vellend M., Geber M.A. (2005). Connections between species diversity and genetic diversity. *Ecology Letters* 8: 767-781.
- Velthuis H.H.W., van Doorn A. (2006). A century of advances in bumblebee domestication and the economic and environmental aspects of its commercialization for pollination. *Apidologie* 37(4): 421-451.
- Verdier-Metz I., Coulon J.B., Pradel P., Viallon C., Albouy H., Berdagué J.L. (2000). Effect of the botanical composition of hay and casein genetic variants on the chemical and sensory characteristics of ripened Saint-Nectaire type cheeses. *Lait* 80: 361-370.
- Verdier-Metz I., Pradel P., Coulon J.B. (2002). Influence of the forage type and conservation on the cheese sensory properties, *Multi-function grasslands: quality forages, animal products and landscapes. 19. General Meeting of the European Grassland Federation, 2002/05/27-30, La Rochelle, France, Durand J.L., ed.: 604-605.*
- Verma M., Brar S.K., Tyagi R.D., Surampalli R.Y., Valero J.R. (2007). Antagonistic fungi, *Trichoderma* spp.: Panoply of biological control. *Biochemical Engineering Journal* 37(1): 1-20.
- Verschwele A., Niemann, P. (1993). Indirect weed control by selection of wheat cultivars, *Maitrise des adventices par voie non chimique IFOAM*, Dijon, Thomas J.-M., ed., IFOAM: 269-273.
- Vertuani S., Angusti A., Manfredini S. (2004). The antioxidants and pro-antioxidants network: an overview. *Current pharmaceutical design* 10(14): 1677-1694.
- Vessey J.K., Pawlowski K., Bergman B. (2005). Root-based N-2-fixing symbioses: Legumes, actinorhizal plants, Parasponia sp and cycads. *Plant and Soil* 274(1-2): 51-78.
- Vetter S., Fox O., Ekschmitt K., Wolters V. (2004). Limitations of faunal effects on soil carbon flow: density dependence, biotic regulation and mutual inhibition. *Soil Biology & Biochemistry* 36(3): 387-397.
- Viaud V., Mérot P. (2002). Nitrate and Dissolved Organic Carbon In Stream Water: Assessing The Buffer Capacities of Hedgerow Systems. Study Case In Brittany, France, *EGS XXVII General Assembly, Nice, 21-26 April 2002: Abstract #3055.*
- Vicens N., Bosch J. (2000). Pollinating efficacy of *Osmia cornuta* and *Apis mellifera* (Hymenoptera: Megachilidae, Apidae) on 'Red Delicious' apple. *Environmental Entomology* 29: 235-240.
- Vicens N., Bosch J. (2000). Weather-dependent pollinator activity in an apple orchard, with special reference to *Osmia cornuta* and *Apis mellifera* (Hymenoptera: Megachilidae and Apidae). *Environmental Entomology* 29: 413-420.
- Vilà M., Pino F., Font X. (2007). Regional assessment of plant invasions across different habitat types. *Journal of Vegetation Science* 18: 35-42.
- Vilà M., Weiner J. (2004). Are invasive plant species better competitors than native plant species? - Evidence from pairwise experiments. *Oikos* 105: 229-238.
- Vile D., Shipley B., Garnier E. (2006). A structural equation model to integrate changes in functional strategies during old-field succession. *Ecology* 87(2): 504-517.
- Villeneuve C., Cadet P. (1998). Interactions of *Helicotylenchus dihystra*, *Pratylenchus sefaensis* and *Tylenchorhynchus gladiolatus* on two plants from the soudano-sahelian zone of West Africa. *Nematropica* 28: 31-39.
- Villeneuve C., Cadet P. (1999). Influence of soil organic matter and ion concentration on some Senegalese plant-parasitic nematodes. *European Journal of Soil Biology* 35: 189-197.
- Violle C., Navas M.L., Vile D., Kazakou E., Fortunel C., Hummel I., Garnier E. (2007). Let the concept of trait be functional! *Oikos* 116: 882-892.
- Vockeroth J.R. (1992). The flower flies of the subfamily Syrphinae of Canada, Alaska and Greenland, *The Insects and Arachnids of Canada*, 18, Agriculture Canada, Ottawa, 456 p.
- Volkl W., Stechmann D.H. (1998). Parasitism of the black bean aphid (*Aphis fabae*) by *Lysiphlebus fabarum* (Hym., Aphidiidae): the influence of host plant and habitat. *Journal of Applied Entomology* 122(5): 201-206.
- von Tiedemann A. (2002). The suppressiveness potential of plants - An approach for "increasing the richness of the soil". *Berichte Über Landwirtschaft*: 57-66.
- Vreeken-Buijs M.J., Hassink J., Brussaard L. (1998). Relationships of soil microarthropod biomass with organic matter and pore size distribution in soils under different land use. *Soil Biology & Biochemistry* 30(1): 97-106.
- Vulliamy B., Potts S.G., Willmer P.G. (2006). The effects of cattle grazing on plant-pollinator communities in a fragmented Mediterranean landscape. *Oikos* 114(3): 529-543.
- Wacker L., Baudois O., Eicheberger-Glinz S., Schmid B. (2008). Environmental heterogeneity increases complementarity in experimental grassland communities. *Basic and Applied Ecology* 9(5): 467-474.
- Wackers F.L. (2004). Assessing the suitability of flowering herbs as parasitoid food sources: flower attractiveness and nectar accessibility. *Biological Control* 29(3): 307-314.
- Waldrop M.P., Firestone M.K. (2006). Response of microbial community composition and function to soil climate change. *Microbial Ecology* 52(4): 716-724.
- Waldrop M.P., Zak D.R., Blackwood C.B., Curtis C.D., Tilman D. (2006). Resource availability controls fungal diversity across a plant diversity gradient. *Ecology Letters* 9(10): 1127-1135.
- Walker M.P., Dover J.W., Sparks T.H., Hinsley S.A. (2006). Hedges and green lanes: Vegetation composition and structure. *Biodiversity and Conservation* 15(8): 2595-2610.
- Wallace H.M., Vithanage V., Exley E.M. (1996). The Effect of Supplementary Pollination on Nut Set of Macadamia (Proteaceae) *Annals of Botany* 78: 765-773.
- Waller P.J., Ljungstrom B.L., Schwan O., Martin L.R., Morrison D.A., Rydzik A. (2006). Biological control of sheep parasites using *Duddingtonia flagrans*: Trials on commercial farms in Sweden. *Acta Veterinaria Scandinavica* 47(1): 23-32.
- Wallis de Vries M.F., Tallowin J.R.B., Dulphy J.P., Sayer M., Diana E. (2004). Effects of livestock breed and stocking rate on sustainable grazing systems: 5. Short-term effects on fauna. *In Land use systems in grassland dominated regions. Proceedings of the 20th General Meeting of the European Grassland Federation, Luzern, Switzerland, 21-24 June 2004, (Lüscher A. et al., eds.), Zurich Switzerland: 626-628.*
- Wang B., Qiu Y.L. (2006). Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 16(5): 299-363.
- Wardle D.A. (2002). *Communities and Ecosystems – Linking the Aboveground and Belowground Components*, University Press, Princeton, NJ, Princeton, 400.

- Wardle D.A. (2006). The influence of biotic interactions on soil biodiversity. *Ecology Letters* 9(7): 870-886.
- Wardle D.A., Bardgett R.D., Klironomos J.N., Setälä H., Van der Putten W.H., Wall D.H. (2004). Ecological linkage between aboveground and belowground biota. *Science* 304: 1634-1637.
- Wardle D.A., Bonner K.I., Barker G.M. (2000). Stability of ecosystem properties in response to above-ground functional group richness and composition. *Oikos* 89: 11-23.
- Wardle D.A., Bonner K.I., Barker G.M., Yeates G.W., Nicholson K.S., Bardgett R.D., Watson R.N., Ghani A. (1999). Plant removals in perennial grassland: vegetation dynamics, decomposers, soil biodiversity and ecosystem properties. *Ecological Monographs* 69: 535-568.
- Wardle D.A., Bonner K.I., Nicholson K.S. (1997). Biodiversity and plant litter : experimental evidence which does not support the view that enhanced species richness improves ecosystem function. *Oikos* 79: 247-258.
- Wardle D.A., Grime J.P. (2003). Biodiversity and stability of grassland ecosystem functioning. *Oikos* 100(3): 622-623.
- Wardle D.A., Williamson W.M., Yeates G.W., Bonner K.I. (2005). Trickle-down effects of aboveground trophic cascades on the soil food web. *Oikos* 111(2): 348-358.
- Wardle D.A., Yeates G.W., Williamson W., Bonner K.I. (2003). The response of a three trophic level soil food web to the identity and diversity of plant species and functional groups. *Oikos* 102(1): 45-56.
- Warren M.S., Hill J.K., Thomas J.A., Asher J., Fox R., Huntley B., Roy D.B., Telfer M.G., Jeffcoate S., Harding P., Jeffcoate G., Willis S.G., Greatorex-Davies J.N., Moss D., Thomas C.D. (2001). Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature* 414: 65-69.
- Waser N.M., Chittka L., Price M.V., Williams N.M., Ollerton J. (1996). Generalization in pollination systems, and why it matters. *Ecology* 77(4): 1043-1060.
- Wasilewska L. (1995). Differences in development of soil nematode communities in single-species and multi-species grass experimental treatments. *Applied Soil Ecology* 2(1): 53-64.
- Waterhouse D.F. (1974). The biological control of dung. *Scientific American* 230: 100-109.
- Way M.J., Cammell M.E. (1981). Effects of weeds and weed control on invertebrate pest ecology. In Proc. Univ. York and Br. Ecol. Soc. and Feder. Br. Plant Pathol., Pitman, London: 443-458.
- Weller D.M., Raaijmakers J.M., Gardener B.B.M., Thomashow L.S. (2002). Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annual Review of Phytopathology* 40: 309-348.
- Wells W.C., Isom W.H., Waines J.C. (1988). Outcrossing rates of six common bean lines. *Crop Science* 28: 177-178.
- Weltzin J.F., Muth N.Z., Von Holle B., Cole P.G. (2003). Genetic diversity and invasibility: a test using a model system with a novel experiment design. *Oikos* 103(3): 505-518.
- Wertz S., Degrange V., Prosser J.I., Poly F., Commeaux C., Freitag T., Guillaumaud N., Le Roux X. (2006). Maintenance of soil functioning following erosion of microbial diversity. *Environmental Microbiology* 8(12): 2162-2169.
- Wertz S., Degrange V., Prosser J.I., Poly F., Commeaux C., Guillaumaud N., Le Roux X. (2007). Decline of soil microbial diversity does not influence the resistance and resilience of key soil microbial functional groups following a model disturbance. *Environmental Microbiology* 9(9): 2211-2219.
- West B.C., Messmer T.A. (2006). Effects of livestock grazing on duck nesting habitat in Utah. *Rangeland Ecology & Management* 59(2): 208-211.
- Westerkamp C. (1991). Honeybees are poor pollinators – why? *Plant Systematics and Evolution* 177: 71-75.
- Westerkamp C., Gottsberger G. (2000). Diversity pays in crop pollination. *Crop Science* 40: 1209-1222.
- Westerman P.R., Wes J.S., Kropff M.J., Van der Werf W. (2003). Annual losses of weed seeds due to predation in organic cereal fields. *Journal of Applied Ecology* 40(5): 824-836.
- Westgard P.H., Flexner L.J., Vanburskirk P., Gonzalves P., Hilton R. (1990). Dispersal pattern of the twospotted spider mite from orchard groundcover into pear. *Bulletin OILB/SROP* 13(2): 53-57.
- Weston L.A. (2005). History and current trends in the use of allelopathy for weed management. *HortTechnology* 15(3): 529-534.
- Westphal A., Becker J.O. (2001). Soil suppressiveness to *Heterodera schachtii* under different cropping sequences. *Nematology* 3: 551-558.
- Westphal C., Steffan-Dewenter I., Tschamtké T. (2003). Mass flowering crops enhance pollinator densities at a landscape scale. *Ecology Letters* 6(11): 961-965.
- Westphal C., Steffan-Dewenter I., Tschamtké T. (2006). Bumblebees experience landscapes at different spatial scales: possible implications for coexistence. *Oecologia* 149: 289-300.
- White A.J., Wratten S.D., Berry N.A., Weigmann U. (1995). Habitat manipulation to enhance biological control of Brassica pests by hover flies (Diptera: Syrphidae). *Journal of Economic Entomology* 88(5): 1171-1176.
- White J., Hodgson J. (2000). New Zealand Pasture and Crop Science, OUP Australia & New Zealand, 336p.
- White P.C.L., Hassall M. (1994). Effects of Management on Spider Communities of Headlands in Cereal Fields. *Pedobiologia* 38(2): 169-184.
- Whitham T.G., Bailey J.K., Schweitzer J.A., Shuster S.M., Bangert R.K., LeRoy C.J., Lonsdorf E.V., Allan G.J., DiFazio S.P., Potts B.M., Fischer D.G., Gehring C.A., Lindroth R.L., Marks J.C., Hart S.C., Wimp G.M., Wooley S.C. (2006). A framework for community and ecosystem genetics: from genes to ecosystems. *Nature Review Genetics* 7: 510-523.
- Whitman W.B., Coleman D.C., Wiebe W.J. (1998). Prokaryotes: The unseen majority. *Proceedings of the National Academy of Sciences of the United States of America* 95(12): 6578-6583.
- Whitmore A.P., Schroder J.J. (2007). Intercropping reduces nitrate leaching from under field crops without loss of yield: A modelling study. *European Journal of Agronomy* 27(1): 81-88.
- Whittington F.M., Dunn R., Nute G.R., Richardson R.I., Wood J.D. (2006). Effect of pasture type on lamb product quality. New developments in Sheepmeat quality, *New Developments in Sheepmeat quality*, 2006/05/24-25, Bristol, British Society of Animal Science: 27-31.
- Whittington R., Winston M.L. (2003). Are bumble bee colonies in tomato greenhouses obtaining adequate nutrition? *Canadian Entomologist* 135(6): 883-892.

- Wiggers M.S., Pratt P.D., Tipping P.W., Welbourn C., Cuda J.P. (2005). Within-plant distribution and diversity of mites associated with the invasive plant *Schinus terebinthifolius* (Sapindales : Anacardiaceae) in Florida. *Environmental Entomology* 34(4): 953-962.
- Wijesinghe D.K., John E.A., Hutchings M.J. (2005). Does pattern of soil resource heterogeneity determine plant community structure? An experimental investigation. *Journal of Ecology* 93: 99-112.
- Wilby A., Thomas M.B. (2002). Natural enemy diversity and pest control: patterns of pest emergence with agricultural intensification. *Ecology Letters* 5(3): 353-360.
- Wilby A., Villareal S.C., Lan L.P., Heong K.L., Thomas M.B. (2005). Functional benefits of predator species diversity depend on prey identity. *Ecological Entomology* 30(5): 497-501.
- Wilhelm W., Wortmann C.S. (2004). Tillage and rotation interactions for corn and soybean grain yield as affected by precipitation and air temperature. *Agronomy Journal* 96: 425-432.
- Wilhoit L.R. (1991). Modelling the population dynamics of different aphid genotypes in plant variety mixtures. *Ecological Modelling* 55: 257-283.
- Williams I.H. (1994). The dependence of crop production within the European Union on pollination by honeybees. *Agricultural Zoology Reviews* 6: 229-257.
- Wilsey B.J., Polley H.W. (2002). Reductions in grassland species evenness increase dicot seedling invasion and spittle bug infestation. *Ecology Letters* 5: 676-684.
- Wilsey B.J., Polley H.W. (2004). Realistically low species evenness does not alter grassland species richness-productivity relationships. *Ecology* 85: 2693-2700.
- Wilsey B.J., Potvin C. (2000). Biodiversity and ecosystem functioning: The importance of species evenness and identity in a Quebec old field. *Ecology* 81(4): 887-893.
- Wilson B.J., Peters N.C.B. (1982). Some studies of competition between *Avena fatua* L. and spring barley. I. The influence of *A. fatua* on yield of barley. *Weed Research* 22: 143-148.
- Winfield A.L. (1963). A study on the effects of insecticides on parasites of larvae of blossom beetles. *Entomologia Experimentalis et Applicata* 6: 309-318.
- Winfree R., Williams N.M., Dushoff J., Kremen C. (2007). Native bees provide insurance against ongoing honey bee losses. *Ecology Letters* 10(11): 1105-1113.
- Wisler G.C., Norris R.F. (2005). Interactions between weeds and cultivated plants as related to management of plant pathogens. *Weed Science* 53: 914-917.
- Wolfe B.E., Husband B.C., Klironomos J.N. (2005). Effects of a belowground mutualism on an aboveground mutualism. *Ecology Letters* 8: 218-223.
- Wolfe L.M. (2002). Why alien invaders succeed: support for the escape-from-enemy hypothesis. *American Naturalist* 160(6): 705-711.
- Wolfe L.M., Elzinga J.A., Biere A. (2004). Increased susceptibility to enemies following introduction in the invasive plant *Silene latifolia*. *Ecology Letters* 7(9): 813-820.
- Wolfe M.S. (2000). Crop strength through diversity. *Nature* 406(6797): 681-682.
- Wolff A., Paul J.-P., Martin J.-L., Bretagnolle V. (2001). The benefits of extensive agriculture to birds : the case of the Little Bustard. . *Journal of Applied Ecology* 38: 963-975
- Wolsing M., Prieme A. (2004). Observation of high seasonal variation in community structure of denitrifying bacteria in arable soil receiving artificial fertilizer and cattle manure by determining T-RFLP of nir gene fragments. *FEMS Microbiology Ecology* 48(2): 261-271.
- Woodcock B.A., Potts S.G., Westburry D.B., Ramsay A.J., Lambert M., Harris S.J., Brown V.K. (2008). The importance of sward architectural complexity in structuring predatory and phytophagous invertebrate assemblages. *Ecological Entomology* 32(3): 302-311.
- Wright D.H. (1983). Species-energy theory: an extension of species-area theory. *Oikos* 41: 496-506.
- Wright J.P., Naeem S., Hector A., Lehman C.L., Reich P.B., Schmid B., Tilman D. (2006). Conventional functional classification schemes underestimate the relationship with ecosystem functioning. *Ecology Letters* 9: 111-120.
- Wyss E. (1995). The effects of weed strips on aphids and aphidophagous predators in an apple orchard. *Entomologia Experimentalis et Applicata* 75(1): 43-49.
- Xiong Y.G. (1990). Final Technical Report of Paulownia Project (Phase II), Académie chinoise d'Agroforesterie.
- Xiong Z.Q., Xing G.X., Tsuruta H., Shen G.Y., Shi S.L., Du L.J. (2002). Field study on nitrous oxide emissions from upland cropping systems in China. *Soil Science and Plant Nutrition* 48(4): 539-546.
- Yarrow N.H., Penning P.D. (2001). The liveweight gain of Limousine x Friesian heifers grazing perennial ryegrass/white clover swards of different clover content and the effects of their grazing on sward botanical composition. *Grass & Forage Science* 56: 238-248.
- Yeates G.W., Bardgett R.D., Cook R., Hobbs P.J., Bowling P.J., Potter J.F. (1997). Faunal and microbial diversity in three Welsh grassland soils under conventional and organic management regimes. *Journal of Applied Ecology* 34(2): 453-470.
- Yergeau E., Bokhorst S., Huiskes A.H.L., Boschker H.T.S., Aerts R., Kowalchuk G.A. (2007). Size and structure of bacterial, fungal and nematode communities along an Antarctic environmental gradient. *FEMS Microbiology Ecology* 59(2): 436-451.
- Young O.A., Cruickshank G.J., Maclean K.S., Muir P.D. (1994). Quality of meat from lambs grazed on seven pasture species in Hawkes Bay. *New Zealand Journal of Agricultural Research* 37(2): 177-186.
- Young O.A., Lane G.A., Priolo A., Fraser K. (2003). Pastoral and species flavour in lambs raised on pasture, lucerne or maize. *Journal of the science of food and agriculture* 83(2): 93-104.
- Yuan F., Ran W., Shen Q.R., Wang D.Z. (2005). Characterization of nitrifying bacteria communities of soils from different ecological regions of China by molecular and conventional methods. *Biology and Fertility of Soils* 41(1): 22-27.
- Zak D.R., Holmes W.E., White D.C., Peacock A.D., Tilman D. (2003). Plant diversity: soil microbial communities, and ecosystem function: are there any links? *Ecology* 84: 2042-2050.
- Zamora D.S., Jose S., Napolitano K. (2007). Competition for nitrogen in a loblolly pine (*Pinus taeda*) - cotton (*Gossypium hirsutum*) alley cropping system in the Southeastern United States, *When Trees and Crops Get Together, Proceedings of the 10th North American Agroforestry Conference*, Québec City, Canada, Campeau A.O.a.S., ed.

- Zang L.S., Liu T.X. (2007). Intraguild interactions between an oligophagous predator, *Delphastus catalinae* (Coleoptera: Coccinellidae), and a parasitoid, *Encarsia sophia* (Hymenoptera: Aphelinidae), of *Bemisia tabaci* (Homoptera: Aleyrodidae). *Biological Control* 41(1): 142-150.
- Zannone L., Assemat L., Rotili P., Jacquard P. (1983). An experimental study of intraspecific competition within several forage crops. *Agronomie* 3(5): 451-459.
- Zavaleta E.S., Hulvey K.B. (2004). Realistic species losses disproportionately reduce grassland resistance to biological invaders. *Science* 306: 1175-1177.
- Zehnder G.W., Hough-Goldstein J. (1990). Colorado potato beetle (Coleoptera: Chrysomelidae) population development and effects on yield of potatoes with and without straw mulch. *Journal of Economic Entomology* 83(5): 1982-1987.
- Zeiter M., Stampfli A., Newbery D.M. (2006). Recruitment limitation constrains local species richness and productivity in dry grassland. *Ecology* 87: 942-951.
- Zeppa G., Giordano M., Gerbi V., Arlorio M. (2003). Fatty acid composition of Piedmont "Ossolano" cheese. *Lait* 83: 167-173.
- Zhang A., Olkowski W. (1989). Ageratum cover crop aids Citrus biocontrol in China. *IPM Practitioner* 11: 8-10.
- Zhang W., Ricketts T.H., Kremen C., Carney K., Swinton S.M. (2007). Ecosystem services and dis-services to agriculture. *Ecological Economics* 64(2): 253-260.
- Zimdahl R.L. (1980). Weed-crop competition : a review, International Plant Protection Center, Oregon State University, 195 p.
- Zobel M., Öpik M., Moora M., Pärtel M. (2006). Biodiversity and ecosystem functioning: It is time for dispersal experiments. *Journal of Vegetation Science* 17: 543-547.
- Zurbrugg C., Frank T. (2006). Factors influencing bug diversity (Insecta: Heteroptera) in semi-natural habitats. *Biodiversity and Conservation* 15(1): 275-294.